



Selections of minimal conditions for a simple intensification and scale up of w-transaminase reactions

Deslauriers, Maria Gundersen; Lloyd, Richard C; Woodley, John

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):


Gundersen, M. T., Lloyd, R. C., & Woodley, J. (2014). Selections of minimal conditions for a simple intensification and scale up of w-transaminase reactions. Poster session presented at Gordon Research Conference: Biocatalysis, Smithfield, RI, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Selections of minimal conditions for a simple intensification and scale up of ω -transaminase reactions

Maria T Gundersen¹, Richard C Lloyd² and John M Woodley¹



1) Department of Chemical and Biochemical Engineering, DTU, DK

2) Dr. Reddy's Chirotech Technology Centre, Cambridge, UK

INTRODUCTION

A step wise decision matrix is presented to quickly evaluate ω -transaminase for a 'simple scale up' in the synthetic direction. Here a 'simple scale up' is defined as a system without specialized equipment or process development, thus a rapid implementation. The three step method consists of: 1. thermodynamic evaluation, 2. biocatalyst screen and 3. inhibition characterization. Each step of the method has a cut off value for easy implementation. Demonstrated by a case study which eliminated reaction pair candidates based on the cut off criteria. Finally, the most promising candidate was intensified.

CASE STUDY



CONCLUSION

We have here outlined and successfully demonstrated a simple method for selecting reaction candidates suitable for a simple and rapid intensification and scale up. This can be valuable in processes where time is of the essence, such as small singular batches and pharmaceutical applications. Although, the ω -transaminase reaction is used here, we nonetheless argue that this method can be applied for other biocatalytic reactions, for well developed enzymes, where the limitations of the implementation is known.

INTENSIFICATION

The best reaction candidate pair 1-Boc-3-piperidone and isopropylamine was successfully and intensified up to 75 g/L and ran at a 25 mL scale. The best isolated product yields were obtained with the 50 g/L reaction where 70% (S)-(+)-3-amino-1-Boc-piperidine was isolated.