



Off-shore Wind Atlas of the Central Aegean Sea: A simple comparison of NCEP/NCAR RE-analysis data, QuickSCAT and ENVISAT Synthetic Aperture Radar (SAR) by use of Wind Atlas Method

Bingöl, Ferhat; Hasager, Charlotte Bay; Karagali, Ioanna; Badger, Jake; Badger, Merete; Foussekis, Dimitri ; Nielsen, Morten; Topouzelis, Konstantinos

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Bingöl, F., Hasager, C. B., Karagali, I., Badger, J., Badger, M., Foussekis, D., ... Topouzelis, K. (2012). Off-shore Wind Atlas of the Central Aegean Sea: A simple comparison of NCEP/NCAR RE-analysis data, QuickSCAT and ENVISAT Synthetic Aperture Radar (SAR) by use of Wind Atlas Method. Poster session presented at EWEA 2012 - European Wind Energy Conference & Exhibition, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Off shore Wind Atlas of Aegean Sea

A simple comparison of RE-analysis data, QuickSCAT and SAR

Ferhat Bingöl(1) Charlotte Hasager(1) Ioanna Karagali(1) Jake Badger(1) Merete Badger(1)
Dimitris Foussekis (2) Morten Nielsen (1) Konstantinos Topouzelis(3)
(1) DTU Wind Energy, Roskilde, DENMARK (2) CRES, Athens, GREECE (3) University of the Aegean, Mitilene, GREECE



Abstract

Offshore wind energy is progressing rapidly in many parts of the world. Among them, the Aegean Sea is a unique area. Until now, several offshore locations (e.g. North Sea [1], Baltic Sea [2]) have been investigated by means of a new technique based on ENVISAT Synthetic Aperture Radar (SAR) data.


In the current study we try to investigate the positive and the negative sides of SAR results at the Central Aegean Sea and compare with two other parallel developing techniques; NCEP/NCAR re-analysis based Ocean wind Atlas scatterometer winds QuickSCAT [3]. WASP software and/or method is used in all steps of the study.

Data Sources

NCEP/NCAR : Re-analysis data for the period of 1971 to 2001 is used. The data has approximately 2° resolution and it consists of 6-hourly model results. Therefore each wind atlas is a product of nearly 60000 samples. Wind Atlases at 4 locations in Central Aegean sea are chosen for the selected area.

QuickSCAT: With a daily global coverage of more than 90% of the world's oceans, QuickSCAT provides a valuable dataset with more than 10 years, between 1999 and 2009, of two daily observations at 10m a.s.l. Data are available at a 0.25° grid. After the source data is filtered for rain, the grid points of the selected area include almost 5000 valid wind speed and wind direction measurements during the 10 years of measurement [3].

ENVISAT SAR: The mission is running since 2001, where high resolution wind fields can be measured at 10m height a.s.l. DTU Wind Energy processes several SAR images a day. The Central Aegean dataset currently includes 500 ENVISAT ASAR WMS scenes from March 2010 and March 2012. SAR data has been used in several other places where in-situ measurement comparisons were available. [1,2].



Conclusion

One can rapidly calculate the wind resource map of an offshore location including statistical parameters by using the WASP based tools we have created for different type of data sources.

A shortcoming of this study is the lack of validation with in-situ measurements which would greatly improve the usage of the datasets. It is intended that several on-shore and off-shore calculations are used in further steps of this study.

Nevertheless, obtained knowledge can be used to locate possible offshore masts for future campaigns, especially when the SAR results are used.

References

- [1] Wind class sampling of satellite SAR imagery for offshore wind resource mapping Badger, M.; Badger, J.; Nielsen, M.; Hasager, C.B.; Peña, A. J. Appl. Meteorol. Climatol. 2010, Vol. 49, p. 2474-2491
- [2] SAR-Based Wind Resource Statistics in the Baltic Sea, Hasager, C. B., Badger, M., Peña, A., Larsén, X.G., Bingöl, F. Remote Sens. 2011, 3, 117-144; doi:10.3390/rs3010117
- [3] Ioanna Karagali. Offshore Wind Energy: Wind and Sea Surface Temperature from Satellite Observations. PhD thesis, DTU Wind Energy, 2012.
- [4] I. Troen and E.L. Petersen. European wind atlas. Technical report, Risø A National Laboratory, Roskilde, 1989.

Method

Our final goal with our data sources was to create wind atlases to generate resource grids for certain areas. The subject in a 'wind atlas' is a systematic and comprehensive collection of regional wind climates (RWC) derived by the wind atlas methodology [4].


Standard wind resource grid calculations are made by using the three different data sets and these datasets are described in the Data Sources section. All applied steps to reach the resource grid are available in the conference proceedings.



Results

NCEP/NCAR derived ocean wind atlas captures only large-scale flow features. Flow features due to meso-scale phenomena are missing. This area provides a test bench for evaluating other wind atlas products derived from alternative reanalysis data set. This will be an important element to the evaluation of the global wind atlas project.

QuickSCAT has a higher than NCEP/NCAR and lower than ENVISAT SAR spatial detail. The averaging inherent in the wind retrieval process of QuickSCAT and NCEP/NCAR eliminates the small scale features seen by SAR.



Acknowledgements

- NCEP/NCAR reanalysis data are provided by the National (USA) Center for Environmental Prediction (NCEP) and the National (USA) Center for Atmospheric Research (NCAR).
- QuickScat data are produced by Remote Sensing Systems and sponsored by the NASA Ocean Vector Winds Science Team. Data are available at www.remss.com.