



## The effects of gliadin on urine metabolome in mice

Roager, Henrik Munch; Zhang, Li; Frandsen, Henrik Lauritz; Licht, Tine Rask

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Roager, H. M., Zhang, L., Frandsen, H. L., & Licht, T. R. (2015). The effects of gliadin on urine metabolome in mice. Abstract from Keystone Symposia - Gut Microbiota Modulation of Host Physiology: The Search for Mechanism, Keystone, Colorado, United States.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Abstract for Keystone Symposia - Gut Microbiota Modulation of Host Physiology: The Search for Mechanism**

March 1-6, 2015

Colorado, USA

### **The effects of gliadin on urine metabolome in mice**

Henrik Munch Roager<sup>1</sup>, Li Zhang<sup>1</sup>, Henrik Lauritz Frandsen<sup>2</sup>, Tine Rask Licht<sup>1</sup>

<sup>1</sup>Division of Food Microbiology, National Food Institute, Technical University of Denmark, Denmark.

<sup>2</sup>Division of Food Chemistry, National Food Institute, Technical University of Denmark, Denmark.

Gliadin, a proline-rich protein of gluten, is thought to modulate the gut microbiota and affect the intestinal permeability and immune system. However, little is known about the long-term effects of gliadin on the host and microbial metabolism. To study this, we compared the urine metabolome of two groups of mice, which were on a high fat diet with and without gliadin, respectively, for 23 weeks. Using liquid chromatography mass-spectrometry (MS) followed by multivariate analyses we were able to show a clear separation of the two groups of mice based on their urine metabolome. Discriminating urinary metabolites were identified by tandem MS and compared to MS libraries and authentic standards. Gliadin mice had higher levels of proline-containing dipeptides most likely originating from the gliadin itself. Furthermore, higher levels of tryptophan- and tyrosine-related metabolites were observed in the gliadin mice. Also, Maillard reaction products and  $\beta$ -oxidized tocopherols were observed in higher levels in the urine of gliadin mice, suggesting increased oxidative stress in the gliadin mice. Indisputably, gliadin affected the urine metabolome. However, the mechanisms behind the observed metabolite changes are yet to be elucidated.