



Growth inhibition of *Aeromonas salmonicida* and *Yersinia ruckeri* by disinfectants containing peracetic acid

Meinelt, Thomas; Phan, T.; Behrens, S.; Pedersen, Lars-Flemming; Wienke, A.; Liu, D.; Straus, David L.

Published in:
Diseases of Aquatic Organisms

Link to article, DOI:
[10.3354/dao02847](https://doi.org/10.3354/dao02847)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Meinelt, T., Phan, T., Behrens, S., Pedersen, L-F., Wienke, A., Liu, D., & Straus, D. L. (2015). Growth inhibition of *Aeromonas salmonicida* and *Yersinia ruckeri* by disinfectants containing peracetic acid. *Diseases of Aquatic Organisms*, 113(3), 207-213. DOI: 10.3354/dao02847

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Growth inhibition of *Aeromonas salmonicida* and *Yersinia ruckeri* by disinfectants containing peracetic acid

Thomas Meinelt^{1,*}, Thy-My Phan¹, Sascha Behrens¹, Andreas Wienke¹,
Lars-Flemming Pedersen², Dibo Liu^{1,3}, David L. Straus⁴

¹Department of Ecophysiology and Aquaculture, Leibniz Institute of Freshwater Ecology and Inland Fisheries, Müggelseedamm 301, 12587 Berlin, Germany

²Technical University of Denmark, DTU Aqua, Section for Aquaculture, Section for Aquaculture, North Sea Research Centre, PO Box 101, 9850 Hirtshals, Denmark

³Faculty of Agriculture and Horticulture, Humboldt-University-Berlin, Invalidenstraße 42, 10115 Berlin, Germany

⁴US Department of Agriculture, Agricultural Research Service, Harry K. Dupree — Stuttgart National Aquaculture Research Center, PO Box 1050, Stuttgart, AR 72160, USA

ABSTRACT: Peracetic acid (PAA) is a therapeutic agent used for disinfection in aquaculture, but it must be investigated thoroughly in order to mitigate diseases without harming the fish. Successful disinfectants (like PAA) should not leave dangerous residues in the environment in order to successfully contribute to sustainable aquaculture. The aim of our study was to compare the effectiveness of 6 commercial PAA products with different molecular PAA:H₂O₂ ratios to reduce bacterial growth of *Aeromonas salmonicida* and *Yersinia ruckeri* and to determine effective concentrations and exposure times. All products reduced colony-forming units (CFUs) of *A. salmonicida* and *Y. ruckeri*. Products with higher molecular PAA:H₂O₂ ratios inhibited growth better than products with lower molecular PAA:H₂O₂ ratios at the same PAA concentration; this indicates that H₂O₂ is not the driving force in the reduction of *A. salmonicida* and *Y. ruckeri* growth by PAA *in vitro*. The practical application of the products with high molecular PAA:H₂O₂ ratios should be prioritized if these pathogens are diagnosed.

KEY WORDS: PAA · Peroxide · H₂O₂ · Furunculosis · Yersiniosis · Fish pathogen · Aquaculture · Bacteria · Water treatment

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

For intensive aquaculture, high animal density and handling can increase the susceptibility of fish to infections. Therefore, therapeutic agents are used for effective fish farm biosecurity and to inactivate potentially pathogenic micro-organisms in aquaculture (Verner-Jeffreys et al. 2004, 2009). However, many therapeutic agents have been banned in the European Union, and elsewhere, because of harmful effects to the environment. Therefore, Schlotfeldt (1998) stated that there is a crisis in the treatment of

diseases in aquaculture, since only a few therapeutic agents are allowed in Europe.

Alternative therapeutic agents, such as hydrogen peroxide (H₂O₂), peracetic acid (PAA) and ozone are able to mitigate diseases without harming the fish. Moreover, these agents leave no dangerous residues in the environment in order to successfully contribute to sustainable aquaculture. The oxidizing agents H₂O₂, PAA and ozone are low-molecular weight compounds and can pass through the bacterial membrane to react with internal cellular components, thereby causing damage to cell structure and the

release of intracellular components (Finnegan et al. 2010).

Freer & Novy (1902) first reported the germicidal activities of PAA, citing the 'excellent disinfection and cold sterilization actions of PAA'. The antimicrobial properties of PAA have been recognized for many years (Baldry 1983, Alasri et al. 1993, Stampi et al. 2001, Crebelli et al. 2005). Consequently, PAA has been routinely used in agriculture, food processing, wastewater treatment and hospitals (Baldry & French 1989a,b, Baldry et al. 1991, 1995, Lefevre et al. 1992, Stampi et al. 2001, Caretti & Lubello 2003, Vinnerås et al. 2003).

Fish pathogens such as *Vibrio anguillarum*, *Flexibacter ovoliticus* and *Aeromonas salmonicida* cause a significant decrease in halibut larval survival (Bergh et al. 1992, 1997). The concentrations of bacteria in halibut larvae were higher overall in recirculated water than in flow-through water (Verner-Jeffreys et al. 2004). As a promising therapeutic agent in aquaculture, 7 mg PAA l⁻¹ decreased colony-forming units (CFUs) of *Salmonella typhimurium* by at least 5 log units after treatment for 1 h (Jolivet-Gougeon et al. 1996). Marchand et al. (2012) tested different PAA products at different concentrations against the fungus *Saprolegnia parasitica* and the bacterium *Flavobacterium columnare*; they found a significant reduction of bacterial growth at 1 mg l⁻¹ for 2 of the substances tested. Other PAA products showed significant bacterial inhibition at concentrations of 2 or 4 mg l⁻¹. Products with a lower molecular PAA:H₂O₂ ratio showed better growth inhibition at lower PAA concentrations in their study.

A. salmonicida is the causative agent of furunculosis (ulcer disease, goldfish ulcer disease, carp erythrodermatitis) in cyprinids (Hiney & Olivier 1999) and motile aeromonad infection in salmonids (Noga 2010). *Yersinia ruckeri* is the causative agent of redmouth disease (enteric redmouth disease [ERM], redmouth, yersiniosis, blood spot, *Y. ruckeri* infection) in

different fish species (Tobback et al. 2007, Noga 2010). Both diseases are costly to the aquaculture industry. Verner-Jeffreys et al. (2009) tested a product containing 5% PAA against *A. salmonicida* ssp. *salmonicida*, *Y. ruckeri*, *Carnobacterium piscicola* and *Lactococcus garvieae* using concentrations of 0.0025, 0.005, 0.01, 0.0165 and 0.025 mg PAA l⁻¹; they found the PAA product was toxic to the bacteria at concentrations of 0.0165 and 0.025 mg l⁻¹.

There are very limited published data on the activity of PAA disinfectants against *A. salmonicida* and *Y. ruckeri*. The aim of the present study was to compare the effectiveness of PAA products with different molecular PAA:H₂O₂ ratios to reduce bacterial growth of *A. salmonicida* and *Y. ruckeri* and to determine effective concentrations and exposure times *in vitro*.

MATERIALS AND METHODS

Six commercially available PAA products with different molecular PAA:H₂O₂ concentrations were tested (Table 1). The product with the highest molecular PAA:H₂O₂ ratio (E400) was set as the reference substance. *Aeromonas salmonicida* and *Yersinia ruckeri* monocultures were obtained from the Chemisches und Veterinäruntersuchungsamt Stuttgart (Stuttgart, Germany).

The bacteria were cultured on CASO agar (Carl Roth) at 22°C in plastic petri dishes. After visual growth was observed, the plates were stored at 4°C and sub-cultured every 2 wk. The transfer of the bacteria to a new plate was done with an inoculating loop by spreading the inoculum. All plates were checked for contaminations via stereo-microscopy. At 72 h after inoculum, bacteria were transferred from an agar plate with an inoculating loop to CASO Bouillon (Carl Roth), vortexed and cultured for 24 h. A 0.1 ml bacterial broth was mixed with 10 ml of each

Table 1. Concentrations of peracetic acid (PAA), hydrogen peroxide (H₂O₂) and the molar ratio of PAA:H₂O₂ in each commercial product used in the present study of products for reduction of bacterial growth. Suppliers: KESLA PHARMA WOLFEN GmbH (Greppin, Germany), AppliChem GmbH (Darmstadt, Germany) and Sigma-Aldrich Co. (St. Louis, Missouri, USA)

ID	Product	Supplier	PAA (%)	H ₂ O ₂ (%)	PAA:H ₂ O ₂
Lspez	Wofasteril L. Spez	KESLA PHARMA WOLFEN	3	40	0.034
E35	Wofasteril 035	KESLA PHARMA WOLFEN	3.5	10	0.156
AC150	Peracetic acid 15%	AppliChem	15	24	0.28
E250	Wofasteril E250	KESLA PHARMA WOLFEN	25	30	0.37
SI400	Peracetic acid solution	Sigma-Aldrich	39	6	2.91
E400	Wofasteril E400	KESLA PHARMA WOLFEN	40	12	1.49

PAA solution to give nominal concentrations of 0.1, 0.5, 1.0, 1.5 and 2.0 mg l⁻¹ in deionized water; these concentrations were chosen based on preliminary experiments. By diluting the PAA, its disinfectant effect is greatly diminished and there was no need for a neutralizer. A geometric serial dilution was prepared with each of these PAA solutions from 10⁻¹ to 10⁻⁷ (*Y. ruckeri*) and from 10⁻¹ to 10⁻⁶ (*A. salmonicida*). At exposure times of 5, 10, 20 and 30 min, CFUs were determined by using the drop-plate method (ISO 2005). For this method, 100 µl from each dilution was distributed drop-wise on a 90 mm sterile CASO agar plate containing 10 ml agar. The reduction of CFUs was determined at different concentrations and exposure times.

An unexposed initial count was a control for the time experiment. After 24 h (*Y. ruckeri*) and 48 h (*A. salmonicida*) of incubation, the CFUs were enumerated by means of a stereo-microscope. The incubation time of *A. salmonicida* was increased due to slower growth. All test procedures were performed under sterile conditions and at 22°C.

The influences of exposure time, concentration and product on the number of CFUs were analyzed by multiple linear regression models. Additionally, box plots were used to illustrate the influence of exposure time and concentration on the number of CFUs for each product.

RESULTS

Aeromonas salmonicida

In general, *A. salmonicida* CFUs decreased with increasing PAA concentration for all products we tested (Table 2). A PAA concentration increase of 1 mg l⁻¹ reduced the CFUs by >2 × 10⁻⁶. The number

Table 2. Multiple linear regression analysis of *Aeromonas salmonicida* colony-forming units. The E400 product has the highest peracetic acid concentration and was set as the reference substance to compare other results

Variable	Regression coefficient	Standard error	p-value
Concentration	-2009046	91029	<0.001
Time	-33672	5742	<0.001
Lspez	1159586	214242	<0.001
E35	1372670	214242	<0.001
AC150	3863693	214242	<0.001
SI400	187615	214242	0.382
E250	152308	214242	0.478

of CFUs also decreased with increasing exposure time for all products. The increase of exposure time by 1 min decreased the CFUs by >30 000. However, the effect is negligible after 10 min exposure time. The effect of E400 in reducing CFUs was significantly different from that of the Lspez, E35 and AC150 products (p < 0.001), but similar to that of the SI400 and E250 products (Table 2).

Exposure time

Our reference substance, the E400 product, reduced the mean *A. salmonicida* CFUs to <2.5 × 10⁻⁶ after an exposure time of 5 min (Fig. 1); this was also true for the Lspez, SI400 and E250 products. The SI400 product showed the strongest effect on reducing *A. salmonicida* CFUs within the same exposure time. For the E35 product, mean *A. salmonicida* CFUs of <2.5 × 10⁻⁶ were realized after an exposure time of 10 min. In contrast, mean CFUs of <2.5 × 10⁻⁶ were not achieved at any exposure times tested with the AC150 product; exposure of AC150 up to 30 min did not lower the CFUs of *A. salmonicida*.

PAA concentration

Increasing PAA concentrations of all tested PAA products resulted in reduced *A. salmonicida* CFUs (Fig. 2). However, an increase of the PAA concentration from 1.5 to 2 mg l⁻¹ did not decrease CFUs further. The Lspez and E35 products effectively reduced the *A. salmonicida* CFUs to <2.5 × 10⁻⁶ at a PAA concentration of at least 1 mg l⁻¹. The same effect was achieved by SI400 and E250 at a PAA concentration of 0.5 mg l⁻¹. In contrast, the AC150 product did not reduce the CFUs to <2.5 × 10⁻⁶ at any tested PAA concentration.

Yersinia ruckeri

In general, the number of *Y. ruckeri* CFUs was lower with increasing concentrations for all products we tested. A PAA concentration increase of 1 mg l⁻¹ reduced the CFUs by >10 × 10⁻⁶ (Table 3). Concentrations of 2 mg PAA l⁻¹ reduced the CFUs to nearly zero. All products reduced the *Y. ruckeri* CFUs after 5 min exposure time. An increase of the exposure by 1 min lowered the CFUs by >1.8 × 10⁻⁶ (Table 3). The Lspez, E35, AC150, SI400 and E250 products were not significantly different from E400 (Table 3).




Fig. 1. Reduction of *Aeromonas salmonicida* colony-forming units (CFUs) by peracetic acid products tested over time. Line: median; box: 25th–75th percentiles (IQR); whiskers: min./max. observation within $1.5 \times$ IQR below/above 25th/75th percentile

Exposure time

All products except E250 reduced the mean CFUs of *Y. ruckeri* to $<15 \times 10^{-6}$ after an exposure time of 5 min (Fig. 3). The E250 product needed a longer exposure time to reduce CFUs.

Table 3. Multiple linear regression analysis of *Yersinia ruckeri* colony-forming units. The E400 product has the highest peracetic acid concentration and was set as the reference substance to compare other results

Variable	Regression coefficient	Standard error	p-value
Concentration	-10665968	2161497	<0.001
Time	-1804624	136019	<0.001
Lspez	1720500	5055315	0.734
E35	4662841	5165197	0.367
AC150	5036282	5081241	0.322
SI400	-863677	5055315	0.864
E250	7161469	5055315	0.158




Fig. 2. Reduction of *Aeromonas salmonicida* CFUs by peracetic acid products tested over concentration. Boxplot limits as in Fig. 1

Product concentration

Following exposure to the E400, E35 and SI400 products at PAA concentrations of 0.5 mg l^{-1} , the mean *Y. ruckeri* CFUs $<15 \times 10^{-6}$ were achieved (Fig. 4). The Lspez, AC150 and E250 products at concentrations of 1 mg PAA l^{-1} reduced CFUs.

DISCUSSION

Aeromonas salmonicida

For the experiment with *A. salmonicida*, we found that concentration, time and PAA product have significant influences on the growth of the bacteria; all products were able to reduce CFUs. Multiple linear regressions indicated that 1 concentration step reduced the CFUs by 2×10^{-6} and 1 time step reduced the CFUs by 33 000 (Table 2). Significant differences in CFUs were found with the E400 versus Lspez, E35 and AC150 products, while the E250 and SI400 products were not significantly different from the E400 product.




Fig. 3. Reduction of *Yersinia ruckeri* CFUs by peracetic acid products tested over time. Boxplot limits as in Fig. 1




Fig. 4. Reduction of *Yersinia ruckeri* CFUs by peracetic acid products tested over concentration. Boxplot limits as in Fig. 1

Compared with concentration, time seems to be of less influence in reducing the bacteria. This implies that the effect of the PAA:H₂O₂ mixture is very rapid, and only short exposure times are necessary to reduce pathogens in the water.

Considering time as well as concentrations, the E400, E250 and SI400 products seem to be the most effective against the Gram-negative bacteria *A. salmonicida*. The minimum inhibitory concentration to affect bacterial growth ranged from 0.5 mg l⁻¹ for E400, SI400 and E250 products to 1 mg l⁻¹ and higher for Lspez, E35 and AC150 products. The minimum exposure time to inhibit bacterial growth was 5 min for all products tested except the E35 product.

Marchand et al. (2012) found that the E250 product showed less inhibition of *Saprolegnia parasitica* and *Flavobacterium columnare*. They also found that the 2 products with the lowest PAA concentrations and very low PAA:H₂O₂ ratios (Lspez and E35) to be significantly more effective on the fungus and the bacterium tested. In contrast, we found the E35, Lspez and AC150 products were less effective than SI400 or E250 products. In the present study, E250 was one of the most efficient products.

Straus & Meinelt (2009) and Meinelt et al. (2009) used *Ichthyophthirius multifiliis* to demonstrate that various PAA products affect fish pathogens differently. Mean growth reductions in our investigations indicated the products with higher concentrations of PAA (and thus a higher molar PAA:H₂O₂ ratio) were more toxic to *A. salmonicida*. This is in contrast to Marchand et al. (2012) and Straus & Meinelt (2009), but in agreement with Wagner et al. (2002). The latter authors found that an additional amount of H₂O₂ did not cause additive toxic effects against *A. salmonicida*, and they confirmed the findings that larger doses of H₂O₂ are required compared to PAA for a similar level of disinfection of this bacterium.

Additionally, there seems to be a species-dependent sensitivity to the PAA products. Whereas *F. columnare* and *S. parasitica* were sensitive to the Lspez and E35 product exposures, *A. salmonicida* and *Y. ruckeri* were more sensitive to the SI400 and E250 products.

Because PAA has a low therapeutic index and concentrations higher than 1 mg l⁻¹ become toxic to fish, the E400, SI400 and E250 products are suitable as a prophylaxis against *A. salmonicida*. For all except

the E35 product, concentrations of 1 mg l⁻¹ with a 5 min exposure could significantly reduce the CFUs of *A. salmonicida* by >10⁻⁶. In our experiments, the E35 product was not very successful. However, in contrast to the findings of Marchand et al. (2012), the SI400 product, and not the Lspez product, had the highest toxicity to *A. salmonicida*. The SI400 product is characterized by the highest PAA:H₂O₂ ratio; thus, PAA might have a higher impact in this equilibrium than H₂O₂ against *A. salmonicida*.

Yersinia ruckeri

For *Y. ruckeri*, the exposure to PAA significantly reduced the CFUs of the bacterium by time and concentration (Table 3). No significant differences could be found between the E400 product and the other products tested.

Inhibition of *Y. ruckeri* was achieved with all PAA products dependent on concentration (Fig. 4). A concentration increase of 1 mg l⁻¹ reduced the CFUs by >10 × 10⁻⁶; prolonging the exposure by 5 min reduced the CFUs by >1.8 × 10⁻⁶ (Table 3). Compared with concentration, time seems to have less influence on reducing the bacteria and confirms the findings on *A. salmonicida*. However, the reduction of CFUs by time and concentration is much stronger than for *A. salmonicida*. Therefore, *Y. ruckeri* is more sensitive to PAA exposure than *A. salmonicida*. Exposure times longer than 5 min are necessary to reduce the growth of *Y. ruckeri*. Prolonging the exposure from 5 to 30 min does not reduce the CFUs any further.

For practical applications, a short exposure of 5 to 10 min is sufficient to reduce the population of bacteria and the stress on fish. An initial concentration of 1 mg l⁻¹ was necessary to reduce the CFUs to <15 × 10⁻⁶ for all products in this study, with the exception of the E35 product. For this product, concentrations of 1.5 mg l⁻¹ were necessary to reduce bacterial growth below 15 × 10⁻⁶. Fig. 4 shows that the efficiency of the products is very different. As found against *A. salmonicida*, the SI400 and E400 products more strongly affect bacterial growth than the other substances, and not all products are useful and manageable for the treatment of all microorganisms in aquaculture. In contrast to disinfecting waste water from sewage plants (Stampi et al. 2001, Kitis 2004) and keeping the low therapeutic index of PAA in mind, only low, non-toxic dosages are suitable in aquaculture. For *A. salmonicida* and *Y. ruckeri*, a strong product-dependent reduction of the CFUs can be realized at even lower PAA concentrations.

SUMMARY

In conclusion, all products reduced CFUs of *A. salmonicida* and *Y. ruckeri*. Products with a higher molecular PAA:H₂O₂ ratio inhibited growth better than products with a lower molecular PAA:H₂O₂ ratio. Therefore, H₂O₂ is not the driving force in the reduction of *A. salmonicida* and *Y. ruckeri* growth by PAA *in vitro*. The practical application of the products with a high molecular PAA:H₂O₂ ratio should be prioritized if these pathogens are diagnosed. Future research should investigate *in vitro* toxicity to pathogens in aquatic systems.

Acknowledgements. We thank the Schreiner-Foundation for Research and Education for financial support. We also thank Bradley Farmer and Cindy Ledbetter for their review of the manuscript. Mention of trade names or commercial products in this article is solely informative and does not imply recommendation or endorsement by the Leibniz Institute of Freshwater Ecology and Inland Fisheries, the DTU Aqua, or the US Department of Agriculture. The USDA is an equal opportunity provider and employer.

LITERATURE CITED

- Alasri A, Valverde M, Roques C, Michel G, Cabassud C, Aptel P (1993) Sporocidal properties of peracetic acid and hydrogen peroxide, alone and in combination, in comparison with chlorine and formaldehyde for ultrafiltration membrane disinfection. *Can J Microbiol* 39:52–60
- Baldry MGC (1983) The bactericidal, fungicidal and sporicidal properties of hydrogen peroxide and peracetic acid. *J Appl Bacteriol* 54:417–423
- Baldry MGC, French MS (1989a) Activity of peracetic acid against sewage indicator organisms. *Water Sci Technol* 21:1747–1749
- Baldry MGC, French MS (1989b) Disinfection of sewage effluent with peracetic acid. *Water Sci Technol* 21: 203–206
- Baldry MGC, French MS, Slater D (1991) The activity of peracetic acid on sewage indicator bacteria and viruses. *Water Sci Technol* 24:353–357
- Baldry MGC, Cavadore A, French MS, Massa G, Rodrigues LM, Schirch PFT (1995) Effluent disinfection in warm climates with peracetic acid. *Water Sci Technol* 31:161–164
- Bergh Ø, Hansen GH, Taxt RE (1992) Experimental infection of eggs and yolk sac larvae of halibut, *Hippoglossus hippoglossus* L. *J Fish Dis* 15:379–391
- Bergh Ø, Hjeltnes B, Skiftesvik AB (1997) Experimental infection of turbot *Scophthalmus maximus* and halibut *Hippoglossus hippoglossus* yolk sac larvae with *Aeromonas salmonicida* subsp. *salmonicida*. *Dis Aquat Org* 29:13–20
- Caretti C, Lubello C (2003) Wastewater disinfection with PAA and UV combined treatment: a pilot plant study. *Water Res* 37:2365–2371
- Crebelli R, Conti L, Monarca S, Feretti D and others (2005) Genotoxicity of the disinfection byproducts resulting from peracetic acid- or hypochlorite-disinfected sewage wastewater. *Water Res* 39:1105–1113

- Finnegan M, Linley E, Denyer SP, McDonnell G, Simons C, Maillard JY (2010) Mode of action of hydrogen peroxide and other oxidizing agents: differences between liquid and gas forms. *J Antimicrob Chemother* 65:2108–2115
- Freer PC, Novy FG (1902) On the formation, decomposition and germicidal action of benzoylacetyl and diacetyl peroxides. *Am Chem J* 27:161–193
- Hiney M, Olivier G (1999) Furunculosis (*Aeromonas salmonicida*). In: Woo PTK, Bruno D (eds) Fish diseases and disorders, Vol 3: Viral, bacterial and fungal infections. CAB International, Wallingford, p 341–425
- ISO (International Organization for Standardization) (2005) ISO 8199: Water quality—General guidance on the enumeration of micro-organisms by culture. ISO, London
- Jolivet-Gougeon A, Braux AS, Sauvager F, Arturo-Schaan M, Cormier M (1996) Influence of peracetic acid on *Escherichia coli* H10407 strain in laboratory microcosms. *Can J Microbiol* 42:60–65
- Kitis M (2004) Disinfection of wastewater with peracetic acid: a review. *Environ Int* 30:47–55
- Lefevre F, Audic JM, Ferrand F (1992) Peracetic acid disinfection of secondary effluents discharged off coastal seawater. *Water Sci Technol* 25:155–164
- Marchand PA, Phan TM, Straus DL, Farmer BD, Stüber A, Meinelt T (2012) Reduction of *in vitro* growth in *Flavobacterium columnare* and *Saprolegnia parasitica* by products containing peracetic acid. *Aquacult Res* 43: 1861–1866
- Meinelt T, Matzke S, Stüber A, Pietrock M, Wienke A, Mitchell AJ, Straus DL (2009) Toxicity of peracetic acid (PAA) to tomites of *Ichthyophthirius multifiliis*. *Dis Aquat Org* 86:51–56
- Noga EJ (2010) Fish disease, diagnosis and treatment, 2nd edn. John Wiley & Sons, London
- Schlotfeldt HJ (1998) Therapienotstand bei Fischen in der Bundesrepublik Deutschland. *Fischer Teichwirt* 49:42–47
- Stampi S, De Luca G, Zanetti F (2001) Evaluation of the efficiency of peracetic acid in the disinfection of sewage effluents. *J Appl Microbiol* 91:833–838
- Straus DL, Meinelt T (2009) Acute toxicity of peracetic acid (PAA) formulations to *Ichthyophthirius multifiliis* theronts. *Parasitol Res* 104:1237–1241
- Tobback E, Decostere A, Hermans K, Haesebrouck F, Chiers K (2007) *Yersinia ruckeri* infections in salmonid fish. *J Fish Dis* 30:257–268
- Verner-Jeffreys DW, Shields RJ, Bricknell IR, Birkbeck TH (2004) Effects of different water treatment methods and antibiotic addition on larval survival and gut microflora development in Atlantic halibut (*Hippoglossus hippoglossus* L.) yolk-sac larvae. *Aquaculture* 232: 129–143
- Verner-Jeffreys DW, Joiner CL, Bagwell NJ, Reese RA, Husby A, Dixon PF (2009) Development of bactericidal and virucidal testing standards for aquaculture disinfectants. *Aquaculture* 286:190–197
- Vinnerås B, Holmqvist A, Bagge E, Albiñ A, Jonsson H (2003) The potential for disinfection of separated faecal matter by urea and by peracetic acid for hygienic nutrient recycling. *Bioresour Technol* 89:155–161
- Wagner M, Brumelis D, Gehr R (2002) Disinfection of wastewater by hydrogen peroxide or peracetic acid: development of procedures for measurement of residual disinfectant and application to physicochemically treated municipal effluent. *Water Environ Res* 74:33–50

Editorial responsibility: David Bruno,
Aberdeen, UK

Submitted: September 25, 2014; Accepted: January 21, 2015
Proofs received from author(s): March 20, 2015