



Retinoic acid signalling is required for the efficient differentiation of CD4+ T cells into pathogenic effector cells during the development of intestinal inflammation

Rivollier, Aymeric Marie Christian; Pool, Lieneke; Frising, Ulrika; Wendland, Kerstin; Agace, William Winston

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Rivollier, A. M. C., Pool, L., Frising, U., Wendland, K., & Agace, W. W. (2015). *Retinoic acid signalling is required for the efficient differentiation of CD4+ T cells into pathogenic effector cells during the development of intestinal inflammation*. Abstract from 17th International Congress of Mucosal Immunology , Berlin, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Retinoic acid signalling is required for the efficient differentiation of CD4⁺ T cells into pathogenic effector cells during the development of intestinal inflammation

Aymeric Rivollier², Lieneke Pool², Ulrika Frising¹, Kerstin Wendland¹ & William Agace^{1,2}

¹ Department of Experimental Medical Sciences, Lund University, Lund, Sweden

² National Veterinary Institute, Danish Technical University, Frederiksberg C, Denmark

Epidemiological studies of vitamin A-deficient populations have illustrated the importance of the vitamin A metabolite retinoic acid (RA) in mucosal immune responses. However, RA seems to be a double-edge sword in CD4⁺ T cell biology. While it sustains the development of foxp3⁺ regulatory T cells, it was also very recently reported to be essential for the stability of the Th1 lineage and to prevent transition to a Th17 program.

Here we explored the role of RA signalling in CD4⁺ T cells during the development of intestinal inflammation in the T cell transfer colitis model. We found that RA signalling-deficient CD4⁺ T cells are less potent at inducing intestinal inflammation compared to their RA signalling-competent counterparts and exhibit a differentiation skewing towards more IFN γ ⁻ IL-17⁺, IL-17⁺IFN γ ⁺ and foxp3⁺ cells, while their capacity to differentiate into IL-17⁻ IFN γ ⁺ Th1 cells is compromised. *In vitro* studies confirm the inefficacy of RA signalling-deficient T cells to generate *bona fide* Th1 cells and demonstrate their aberrant increased ROR γ t expression while their differentiation into Th17 remains unaffected. Surprisingly, RA signalling-deficient CD45RB^{lo} regulatory T cells (Tregs) are however as efficient as their RA signalling-competent counterparts to inhibit colitis development.

Together our results indicate that RA, through its receptor RAR α , negatively regulates the early expansion of CD4⁺ T cells during colitis and is necessary for the generation of colitogenic Th1/Th17 cells, while it is dispensable for the protective function of Treg cells. We are currently deciphering the mechanisms of these effects of RA on CD4⁺ T cells.