



Modelling and predicting growth of psychrotolerant pseudomonads in milk and cottage cheese

Martinez Rios, Veronica; Østergaard, Nina Bjerre; Rosshaug, Per Sand; Dalgaard, Paw

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Martinez Rios, V., Østergaard, N. B., Rosshaug, P. S., & Dalgaard, P. (2015). Modelling and predicting growth of psychrotolerant pseudomonads in milk and cottage cheese. Poster session presented at 9th International Conference on Predictive Modelling in Food, Rio de Janeiro, Brazil.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Modelling and predicting growth of psychrotolerant pseudomonads in milk and cottage cheese

Veronica Martinez-Rios¹, Nina B. Østergaard¹, Per S. Rosshaug¹, Paw Dalgaard¹

¹Microbial Food Safety and Quality, Division of Microbiology and Production, National Food Institute (DTU Food), Technical University of Denmark, Kgs. Lyngby, Denmark

Objective

To develop and validate mathematical models for growth of psychrotolerant pseudomonads in chilled milk and in cottage cheese with cultured cream dressing.

Introduction

Psychrotolerant pseudomonads are associated with sensory spoilage, due to production of fruity and nauseous off-odours in e.g. raw milk, pasteurized milk and cottage cheese. Predictive food microbiology models have the potential to evaluate the effect of temperature on microbial growth during distribution as well as be used to determine how product characteristics can be modified to reduce growth to an acceptable level. However, it is challenging to model and predict growth of spoilage or pathogenic microorganisms in fermented foods, such as cottage cheese, due to inhibiting substances produced by lactic acid bacteria during fermentation or later during storage and distribution (Østergaard et al. 2014).

Methods

Growth kinetics of psychrotolerant pseudomonads were determined using Bioscreen C experiments, challenge tests, storage trial and literature data. The effect of storage temperature and product characteristics on growth rates (μ_{max}) were described using a simplified cardinal parameter model and the gamma concept (Østergaard et al., 2014; Zwietering et al., 1992). The developed broth /Bioscreen C model included the effect of temperature, pH, NaCl/ a_w , lactic-, sorbic acid and their interaction (Le Marc et al., 2002). Subsequently, the reference growth rate parameter (μ_{ref} at 25°C) was fitted to a total of 35 μ_{max} -values from cottage cheese with cultured cream dressing. Lag time models were developed from growth curves in products by calculation of relative lag times. μ_{max} -models for milk and cottage cheese were evaluated by calculation of bias (B_f)- and accuracy (A_f)- factors from literature and new experimental μ_{max} data (Table 1).

Table 1. Evaluation of growth rate models for psychrotolerant pseudomonads

Model	Data	n	B_f	A_f
Broth ($\mu_{ref} = 1.03 \text{ h}^{-1}$)	Pasteurized milk	33	1.08	1.32
Calibrated to cottage cheese ($\mu_{ref} = 0.62 \text{ h}^{-1}$)	Cottage cheese and raw milk	26	1.08	1.43

The acceptable simulation zone (ASZ) method was used to evaluate the performance of the combined lag time and μ_{max} models on growth (log CFU/g) at constant and dynamic temperatures (Fig.1).




Figure 1. ASZ method used to compare observed and predicted growth of psychrotolerant pseudomonads in cottage cheese with cultured cream dressing (log CFU/g)

Results

The new models were successfully validated for 59 growth curves of psychrotolerant pseudomonads in dairy products (Fig. 1; Table 1). The ASZ method showed the new models for pasteurized milk (ASZ = 96%) as well as cottage cheese and raw milk (ASZ = 78%) to successfully predict growth of psychrotolerant pseudomonads at constant and dynamic storage temperatures.

Conclusions

The present study developed and validated mathematical models to predict growth of psychrotolerant pseudomonads in chilled milk and cottage cheese with cultured cream dressing. The cottage cheese model can be used to evaluate the effect of product reformulations on growth.

