

## **A 10-days heatwave around flowering superimposed on climate change conditions significantly affects production of 22 barley accessions**

**Ingvordsen, Cathrine Heinz; Lyngkjær, Michael F.; Peltonen-Sainio, Pirjo; Mikkelsen, Teis Nørgaard; Bagger Jørgensen, Rikke**

*Published in:*  
Procedia Environmental Sciences

*Link to article, DOI:*  
[10.1016/j.proenv.2015.07.239](https://doi.org/10.1016/j.proenv.2015.07.239)

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Ingvordsen, C. H., Lyngkjær, M. F., Peltonen-Sainio, P., Mikkelsen, T. N., & Bagger Jørgensen, R. (2015). A 10-days heatwave around flowering superimposed on climate change conditions significantly affects production of 22 barley accessions. *Procedia Environmental Sciences*, 29, 160-161. DOI: 10.1016/j.proenv.2015.07.239

## **DTU Library** Technical Information Center of Denmark

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Agriculture and Climate Change - Adapting Crops to Increased Uncertainty (AGRI 2015)

## A 10-days heatwave around flowering superimposed on climate change conditions significantly affects production of 22 barley accessions

*C.H. Ingvordsen<sup>a\*</sup>, M.F. Lyngkjær<sup>b</sup>, P. Peltonen-Sainio<sup>c</sup>,*

*T.N. Mikkelsen<sup>a</sup>, R.B. Jørgensen<sup>a</sup>*

<sup>a</sup>*Technical University of Denmark, Frederiksborgvej 399, 4000 Roskilde, Denmark*

<sup>b</sup>*University of Copenhagen, Thorvaldsensvej 40 1871 Frederiksberg, Denmark*

<sup>c</sup>*MTT Agrifood Research Finland Plant Production Jokionen 31600 Finland*

### Abstract

Extreme climate events as heatwaves, floods and storms cause acute changes in season variability influencing primary production and are very likely to increase in magnitude and/or frequency (IPCC, AR5, WGI)<sup>1,2</sup>.

In the present study 22 primarily Nordic barley accessions were grown in four basic climate treatments of 1) 19/12°C (day/night) and 400 ppm carbon dioxide concentration [CO<sub>2</sub>] mimicking ambient South Scandinavian summer conditions, 2) elevated temperature (+5°C day/night), 3) elevated [CO<sub>2</sub>] at 700 ppm and 4) the combination of elevated temperature and [CO<sub>2</sub>]. Temperature and [CO<sub>2</sub>] were at levels representing a worst case scenario (~RCP8.5, IPCC) at the end of the 21<sup>st</sup> century. A 10 day-heatwave of 33/22°C (day/night) was superimposed around the time of flowering on the basic climate treatments.

The superimposed heatwave decreased overall grain yield in all combinations, however, vast variation in response was identified among accessions. In the two-factor treatment the decrease in grain yield varied from 2-80%. The heatwave caused the strongest overall effect in the treatment of elevated [CO<sub>2</sub>] decreasing grain yield by 48% and the least effect (35%) was observed under elevated temperature suggesting elevated temperature to have a priming effect. In all heatwave treatments allocation of biomass was changed, increasing aboveground vegetative biomass and decreasing grain yield as previously reported<sup>3</sup>.

The treatment with the combination of elevated temperature, [CO<sub>2</sub>] and the superimposed heatwave may best represent a future climate scenario since more than one climate factor most likely will change at a time. From the basic ambient treatment to the two-factor treatment including heatwave, grain yield decreased 52%.

Our study emphasizes the need for assessing the effects of extreme events under climate change conditions on numerous accessions in order to select appropriate genotypes for breeding future cultivars that can secure the primary production.

© 2015 Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the organizing committee of the Agriculture and Climate Change - Adapting Crops to Increased Uncertainty (AGRI 2015)

*Keywords:* biomass allocation, climate change, extreme events; genotype differences; heat exposure; production stability; temperature priming

---

## References

1. Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., et al. Long-term Climate Change: Projections, Commitments and Irreversibility. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M., editors. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press: 2013.
2. Fischer EM, Schär C. Consistent geographical patterns of changes in high-impact European heatwaves. *Nat Geosci* 2010; **3**:398–403.
3. Bencze S, Veisz O, Bedo Z. Effects of high atmospheric CO<sub>2</sub> and heat stress on phytomass, yield and grain quality of winter wheat. *Cereal Res Commun* 2004; **32**:75–82.