Chemotaxonomy of the genus *Stemphylium*

Olsen, Kresten Jon Kromphardt; Andersen, Birgitte

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Olsen, K. J. K., & Andersen, B. (2016). *Chemotaxonomy of the genus Stemphylium*. Poster session presented at 9th Joint Natural Products Conference 2016, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Email: krjko@dtu.dk

Chemotaxonomy of the genus *Stemphylium*

Kresten Jon Kromphardt Olsen and Birgitte Andersen

Department of Biotechnology and Biomedicine Technical university of Denmark, Kgs. Lyngby, Denmark

Abstract

The filamentous fungal genus *Stemphylium* (Anamorphic *Pleospora*) is often found on various crops, and especially the common animal feed plant *Medicago sativa* (alfalfa) is often infected by this plant pathogen. With this in mind it is important to consider what consequences such a contamination can have, e.g. production of mycotoxins. (Firsvad *et al.* (2009))

A clade of *Stemphylium* spp. i.e. *S. herbarum*, *S. alfalfae*, *S. sedicola*, *S. tomatonis* and *S. vesicarium* are troublesome to distinguish as they share both morphological and molecular characteristics. This study has focused on using chemotaxonomy via HPLC-UV/VIS-MS analysis as a third method for identification and to investigate the chemical potential of the genus *Stemphylium*.

A total of 253 unique compounds were used for chemotaxonomy and the majority of these were unknown compounds.

Morphological identification

The morphology of *Stemphylium* conidia. Some species require special knowledge of the morphology (I-VI) to identify, while others are easier to identify by this method (VII-IX).

Figure 1. Comparison of micromorphology within the *Stemphylium* genus. I – *S. eturmiunum*, II – *S. herbarum*, III – *S. astragali*, IV – *S. gracilareae*, V – *S. alfalfae*, VI – *S. majusculum*, VII – *S. loti*, VIII – *S. sarciniforme* and IX – *S. lancipes*. Same scale bar for all pictures.

Examples of compounds

The majority of known *Stemphylium* compounds are of polyketide origin.

Conclusion and perspectives

Chemotaxonomy with the genus of *Stemphylium* gave a new tool for identification of the individual species. Also, a list of detected compounds within the genus has been made. With this list it is possible to mine for novel bioactivities as it is possible to analyze vastly different metabolic profiles within the same genus.

Chemical identification

Group C

Chemotaxonomy gave some resolution of group C. Identification via phylogeny would give more resolution but a total of 8 chemical groups were identified. Some were of known isolates e.g. *S. gracilareae*, *S. majusculum* and *S. eturmiunum*.

Interestingly *S. lycopersici* (group A) and *S. callistephi* (group D) clustered within group C.

Group A, B, D, E, F

All species of this group are very easy to distinguish from group C. These also have solid separation and many unique compounds to differentiate them.

References

Firsvad JC, Andersen B, Thrane U. The use of secondary metabolite profiling in chemotaxonomy of filamentous fungi. Mycological research 2008; 112: 231-240