Solar UV Irradiation-Induced Production of Greenhouse Gases from Plant Surfaces: From Leaf to Earth

Mikkelsen, Teis Nørgaard; Bruhn, Dan; Ambus, Per

Published in:
Progress in Botany

Link to article, DOI:
[10.1007/124_2016_10](https://doi.org/10.1007/124_2016_10)

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Mikkelsen, T. N., Bruhn, D., & Ambus, P. (2016). Solar UV Irradiation-Induced Production of Greenhouse Gases from Plant Surfaces: From Leaf to Earth. In *Progress in Botany* Springer. Progress in Botany
https://doi.org/10.1007/124_2016_10

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Solar UV irradiation-induced production of greenhouse gases from plant** 2 **surfaces: From leaf to Earth**

3 Teis Nørgaard Mikkelsen¹⁾, Dan Bruhn²⁾ and Per Ambus³⁾

4 ¹⁾Department of Environmental Engineering, Technical University of Denmark, Building 115, DK - 2800 Kgs.
5 Lyngby, Denmark. e-mail: temi@env.dtu.dk

6 ²⁾Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H, DK- 9220 Aalborg
7 East, Denmark. e-mail: db@bio.aau.dk

8 ³⁾Department of Geosciences and Natural Resource Management, University of Copenhagen, Øster
9 Voldgade 10, DK-1350 Copenhagen K, Denmark. e-mail: peam@ign.ku.dk

10

11 **Abstract**

12 During the past few decades it has been documented that the ultra-violet (UV) component of natural
13 sunlight alone or in combination with visible light can instantaneously stimulate aerobic plant production of
14 a range of important trace gases: CH₄, CO₂, CO, short-chain hydrocarbons/ non-methane volatile organic
15 compounds (NMVOC), NO_x and N₂O. This gas production, near or at the plant surface, is a new discovery
16 and is normally not included in emission budgets (e.g. by the Intergovernmental Panel on Climate Change,
17 IPCC) due to a lack of information with respect to validation and upscaling. For CH₄ it is known that the light
18 dose controls emission under ambient and artificial light conditions, but the atmospheric gas composition
19 and other environmental factors can influence gas production as well. Several plant components, including
20 pectin and leaf wax, have been suggested as a precursor for CH₄ production, but underlying mechanisms
21 are not fully known. For other gases such generating processes have not been established yet and
22 mechanisms remain hypothetical. Field measurements of UV-induced emissions of the gases under natural
23 light conditions are scarce. Therefore, realistic upscaling to the ecosystem level is uncertain for all gases.
24 Nevertheless, based on empirical response curves, we propose the first global upscaling of UV induced N₂O
25 and CO to illustrate emission ranges from a global perspective and as a contribution to an ongoing
26 quantification process. When scaled to the global level, the UV-induced emission of CO by vegetation
27 surfaces amounts to up to 22 Tg yr⁻¹, which equals 11-44% of all the natural terrestrial plant sources
28 accounted for so far. The total light-driven N₂O emissions amount to 0.65-0.78 Tg yr⁻¹, which equals 7-24%
29 of the natural terrestrial source strength accounted for (range 3.3 - 9 Tg N yr⁻¹). In this review, we
30 summarize current knowledge, based on experimental work with sunlight and artificial light, and estimate

31 potential emission ranges and uncertainties, placing the available data into perspective. We discuss the
32 state of the art in proposed mechanisms, precursors, and environmental relationships, we consider the
33 relevance of measured emission rates, and we also suggest a range of future research topics. Furthermore
34 we propose and describe methods and techniques that can be used for future research.

35 *Keywords: Atmospheric chemistry, air pollution, UV-A, UV-B, stable isotopes techniques, N₂O, CH₄, NO_x,
36 NO_y, CO, CO₂, short-chain hydrocarbons, NMVOC, upscaling, vegetation, plants*

37

38 **Content**

39 1 Introduction

40 2 Light induced gases at the plant surface

41 2.1 Well-mixed greenhouse gases (CO₂, CH₄ and N₂O)

42 2.2 Short lived gases (CO, NO_x, NMVOC)

43 3 Methods & techniques

44 4 Mechanisms, sources and emissions

45 4.1 UV-Radiation penetration through the canopy

46 4.2 UV-Radiation penetration through the leaf

47 4.3 UV-Photochemistry mechanism in and on plant surfaces

48 4.3.1 Action spectra

49 4.3.2 UV-response functions

50 4.3.3 Temperature interactions

51 4.3.4 [O₂] or O-radicals dependency

52 4.4 Precursors

53 5 Upscaling

54 5.1 Upscaling of methane, CH₄

55 5.2 Upscaling of carbon monoxide, CO

56 5.3 Upscaling of nitrous oxide, N₂O

57 6 Perspectives and conclusions

58 6.1 Realistic emission rates

59 6.2 Future studies

60 6.3 Known gas emission stimulated by UV

61 6.4. Perspectives

62 Appendix A Methods & techniques

63 Appendix B Global drivers for CO and N₂O contribution

64 Appendix C Response functions for global CO and N₂O contribution

65

66

67

68 **1 Introduction**

69 For decades it has been recognized that sunlight plays significant roles in atmospheric chemistry and that
70 the UV component is the driving force for tropospheric photochemical processes. For instance, UV-
71 photolysis of ozone generates excited state oxygen atoms, which react with water vapour to constitute the
72 primary source of hydroxyl radical (OH). OH is the major component for the overall oxidising capacity of the
73 troposphere (Isaksen et al. 2009). Lately, a number of trace gases are reported to be released by plants in
74 instantaneous responses to UV-radiation, such as methane (CH₄) (Bruhn et al. 2007, 2009, 2012, 2014a;
75 Röckmann et al. 2007; Vigano et al. 2007, 2008, 2009; McLeod et al. 2008; McLeod and Keppler, 2010;
76 Messenger et al. 2009; Fraser et al. 2015), carbon monoxide (CO) (Derendorp et al. 2011a; Bruhn et al.
77 2013), short-chain hydrocarbons (Derendorp et al. 2011b; Fraser et al. 2015), mono-nitrogen oxides (NO_x
78 and NO_y) (Hari et al. 2003; Raivonen et al. 2006, 2009) and nitrous oxide (N₂O) (Bruhn et al. 2014b).

79 These UV-driven gas productions are in principle newly revealed terrestrial sources that remain to be
80 considered in greenhouse gas accounting. Production of all gases stimulated by UV involves emissions at
81 relatively low rates, and the group of gases are diverse and have different impacts and interactions with the
82 atmosphere. However, several of the gases are potent greenhouse gases or interact with the turnover of
83 atmospheric greenhouse gases.

84 We describe current knowledge of methods and techniques for measuring these gases, with a particular
85 focus on the special requirements needed to accomplish flux measurements under controlled UV-
86 exposures and at generally very low rates. The distinction between laboratory and field measurements is
87 addressed with respect to techniques and deductions. We also report on current understanding of the
88 possible mechanisms and sources behind these gas productions, comment on current upscaling attempts,
89 and present the first upscaling and quantification of UV stimulated CO and N₂O emissions. Finally, we
90 highlight the perspectives of the newly discovered UV stimulated gas sources with respect to research
91 needs and impact on current research.

92

93 **2 Light induced gases at the plant surface**

94 Here we summarize the important features of most of the gases that are known to be formed at the plant
95 surface during exposure to UV light. The gases can be divided into two groups: i) well-mixed greenhouse
96 gases (CO₂, CH₄ and N₂O), and ii) short-lived gases (CO, NO_x, non-methane volatile organic compounds
97 (NMVOC)).

98 **2.1 Well-mixed greenhouse gases (CO₂, CH₄ and N₂O)**

99 The atmospheric concentrations of the greenhouse gases carbon dioxide (CO₂), methane (CH₄), and nitrous
100 oxide (N₂O) have all increased since 1750 due to human activity. In 2011 the concentrations of these
101 greenhouse gases were 391 ppm, 1803 ppb, and 324 ppb, and exceeded pre-industrial levels by about 40%,
102 150%, and 20%, respectively (Table 2.1). Concentrations of CO₂, CH₄, and N₂O now substantially exceed the
103 highest concentrations recorded in ice cores during the past 800,000 years (Masson-Delmotte et al. 2013).
104 The mean rates of increase in atmospheric concentrations over the past century are unprecedented in the
105 last 22,000 years (IPCC, 2013). In Table 2.1 we present the Global Warming Potential (GWP) that integrates
106 radiative forcing (RF) out to a particular time horizon, in this case 100 years. The GWP can be interpreted as
107 an index of the total energy added to the climate system by a component in question relative to that added
108 by CO₂ (Myhre et al. 2013). There are multiple sources of all three gases, which can be divided into two
109 main groups, anthropogenic and natural (Table 2.1). Quantification of the various source strengths from
110 both groups remain uncertain and global budgets remain unclear (Ciais et al. 2013).

111 The main anthropogenic CO₂ sources are burning of fossil fuels (coal, oil and gas), deforestation and
112 production of cement (Ciais et al. 2013). The removal of anthropogenic CO₂ from the atmosphere by
113 natural processes will take a few hundred thousand years (Ciais et al. 2013). The natural CO₂ sources are
114 autotrophic and heterotrophic respiration, decomposition of plant tissues (litter and soil carbon that is
115 released back into the atmosphere) and additional disturbance processes (e.g., natural fires). The natural
116 source strength is 20 times higher than the anthropogenic emission, but is counterbalanced by natural CO₂
117 uptake from the atmosphere by plant photosynthesis (Beer et al. 2010).

118 Massive increases in the number of domestic ruminants, natural gas extraction and use, expansion of rice
119 paddy agriculture and establishment of urban landfills and waste dumps represent the dominant
120 anthropogenic CH₄ sources (Stocker et al. 2013). Wetlands are the dominant natural source of atmospheric
121 CH₄ (EPA 2010). During the last two decades, natural sources of CH₄ have accounted for 35 to 50% of the
122 decadal mean global emissions (Ciais et al. 2013).

123 The anthropogenic N₂O sources are primarily agricultural, and the anthropogenic part accounts for
124 approximately 40% of the total emission (Ciais et al. 2013). Natural sources are constituted by upland soils
125 and riparian areas together with oceans, estuaries, and rivers (EPA 2010). Human-induced perturbations of
126 the nitrogen cycle, in addition to interactions with CO₂ sources and sinks, affect emissions of N₂O both on
127 land and from the ocean (Stocker et al. 2013). It is likely that N₂O emissions from soils will increase due to

the increased demand for feed/food and the reliance of agriculture on nitrogen fertilizers. Climate warming will likely amplify agricultural and natural terrestrial N₂O sources (Ciais et al. 2013).

Table 2.1. Important features of gases that are known to be formed at the plant surface during exposure to UV light

GAS	Atm. Conc.	Ref	Lifetime (years)	Ref	GWP 100 year	Ref	Source		Ref	Units
							Antropogenic	Natural		
CH ₄	1803 ppb	3	9.1	3	28	5	354±45	202±35	1	Tg CH ₄ yr ⁻¹
CO ₂	391 ppm	3	n.a.	-	1	5	8.3 ± 0.7	n.a.	1	PgC yr ⁻¹
N ₂ O	324 ppb	3	131	3	265	5	6.9 (2.7–11.1)	11.0 (5.4–19.6)	1	TgN yr ⁻¹
CO	80 ppb	4	months	5	5.3** ± 2.3	7	608	50-200***	8	TgC yr ⁻¹
NO _x	5-999 ppt	4	hours	3	-159** ± 79	7	37.5	11.3	2	TgN yr ⁻¹
NM VOC	n.a.	-	hours - months	3	n.a.	-	126.9	440-720*	6	TgC yr ⁻¹

*) Only isoprene and monoterpenes, **) direct and indirect aerosol effects included, ***) only from plants.

References: 1) Ciais, et al. 2013; 2) Dentener et al. 2006; 3) Hartmann et al. 2013; 4) IPCC, 2001; 5) Myhre et al. 2013; 6) Boucher et al. 2013, 7) Shindell et al. 2009; 8) Tarr et al. 1995

2.2 Short lived gases (CO, NO_x, NM VOC)

Emissions of CO, NM VOCs and NO_x (NO + NO₂) do not have a direct effect on RF, but affect climate indirectly as precursors to tropospheric O₃ and aerosol formation, and their impacts on hydroxyl-concentrations and CH₄ lifetime. NM VOCs include aliphatic, aromatic and oxygenated hydrocarbons (e.g., aldehydes, alcohols and organic acids), and have atmospheric lifetimes ranging from hours to months. Global coverage of NM VOC measurements is poor, except for a few compounds (Hartmann et al. 2013). Emissions of CO and NM VOC are virtually certain to have induced a positive RF via production of the climatic drivers CO₂, CH₄ and O₃, while emissions of NO_x are likely to have induced a net negative RF. (Table 2.1; IPCC, 2013). With its lifetime of 2 to 3 months, the effect of CO emission is less dependent on location than is the case for NO_x (Myhre et al, 2013). Due to their short atmospheric lifetime (hours), NO_x concentrations are highly variable in time and space. Solomon et al. (2007) described the potential of satellite observations of NO₂ to verify and improve NO_x emission inventories and their trends, and reported NO₂ increases of 50% over the industrial areas of China from 1996 to 2004. An extension of this analysis reveals increases between of 1.7x and 3.2x over parts of China, while over Europe and the USA NO₂ has decreased by 30 to 50% between 1996 and 2010 (Hilboll et al. 2013).

152 The major sources of atmospheric CO are in situ production by oxidation of hydrocarbons (mostly CH₄ and
153 isoprene) and direct emission resulting from incomplete combustion of biomass and fossil fuels. The
154 anthropogenic CO emission is estimated to be 608 TgC yr⁻¹ (Table 2.1); natural sources have been
155 estimated to account for up to half of the global CO emissions (Khalil and Rasmussen, 1990), and direct
156 emissions from plants are estimated to be 50-200 TgC yr⁻¹ (Tarr et al. 1995). An analysis of MOPITT
157 (Measurements of Pollutants in the Troposphere) and AIRS (Atmospheric Infrared Sounder) satellite data
158 suggest a clear and consistent decline of CO columns for 2002–2010 over a number of polluted regions in
159 Europe, North America and Asia, with a global trend of about –1% yr⁻¹ (Yurganov et al. 2010; Fortems-
160 Cheiney et al. 2011; Worden et al. 2013; Hartmann et al. 2013).

161 Reports on trends in a range of NMVOCs generally indicate a decline over urban and rural regions of North
162 America and Europe, on the order of a few percent to more than 10% yr⁻¹ (Hartmann et al. 2013). The
163 anthropogenic emission is between 15 and 22% of the total NMVOC emissions (Table 2.1)

164

165 **3 Methods & techniques**

166 UV-induced gaseous emissions from specific substances, plant organs, whole plants or whole ecosystems
167 have classically been studied under controlled environmental conditions by employment of sealed
168 enclosures ranging in complexity from simple commercial test tubes to highly advanced plant cuvettes
169 (plant parts) or whole chamber enclosures (plants or plant and soil communities).

170 Generally the studied gas components are emitted at very low rates from plant surfaces, and in order to
171 achieve detectable levels of gas accumulation the use of enclosures is required. A static enclosure that is
172 operated by manual sampling or connected in a closed gas sampling loop to the analyzer (e.g. Bruhn et al.
173 2009; 2014b) provides high analytical sensitivity for determining changes in gas mixing ratios, but may
174 require appropriate meticulous techniques to control undesired changes in other gas components, e.g.
175 moisture, CO₂ and O₃. Alternatively an open flow-through design may be used whereby the enclosure is
176 continuously purged with ambient- or zero-air (e.g. Vigano et al. 2008). Meanwhile, deployment of
177 enclosures is associated with multiple challenges that may affect the experimental conditions in
178 uncontrolled and undesired directions, leading to experimental flaws and artefacts.

179 Environmental controls of temperature, humidity and air composition are crucial for work with biological
180 materials in enclosures and to reveal important abiotic controllers for the investigated processes.

181 Complications by uncontrolled changes in the environment may further be augmented by the fact that

182 relatively long enclosure times are needed in order to uncover low reaction constants. It is beyond the
183 scope of this article to provide a detailed protocol of principles and methodologies for the use of sealed
184 enclosures to study gas exchange from surfaces. Instead, the reader is referred to literature providing
185 detailed descriptions on flux-chamber design and applications (e.g. de Klein and Harvey, 2015; Altimir et al.
186 2002; Skiba et al. 1992). In appendix A, we expand methods & techniques with respect to temperature,
187 chamber material, surface reactions and reactive species, analysis of gas mixing ratios and light sources.

188

189 **4 Mechanisms, sources and emissions**

190 Sunlight can induce trace gas emission from plants by several mechanisms from different precursors. Here
191 we focus on direct effects of UV radiation on trace gas emission from terrestrial plant surfaces as these are
192 largely ignored in global budgets. Indirect effects of UV on trace gas emission and other physiological
193 functions are reviewed elsewhere (Caldwell et al. 1995, 1999; Björn et al. 1996; Rozema et al. 1997; Bruhn
194 et al. 2012).

195

196 **4.1 UV-Radiation penetration through the canopy**

197

198 Leaves throughout a canopy are affected not only by the PAR and IR-spectrum of solar radiation but also by
199 UV radiation. Leaves do not transmit UV radiation but reflect some (up to ca. 6%) UV radiation (Grant
200 1997), and the high energy light still penetrates the canopy. Canopy structure, leaf area index (LAI), the
201 extent of direct and diffuse radiation all influence UV penetration through the canopy (Brown et al. 1994;
202 Shulski et al. 2004) and leaves. The penetration by UV-B varies less than that of PAR with leaf inclination
203 due to the higher diffuse component of UV light than PAR (Caldwell 1981; Deckmyn et al 2001). For
204 example, canopy UV-B transmittance (τ) may vary with LAI between $\tau = \exp(-1.01 \text{ LAI})$ and $\tau = \exp(-$
205 $0.17(\text{LAI}-1))$ depending on species and degree of clear sky (Shulski et al. 2004).

206

207 **4.2 UV-Radiation penetration through the leaf**

208

209 UV-B absorption of adaxial leaf cuticles caused by pigments (chromophores, e.g. flavonoids and other
210 phenolic compounds covalently bound to cutin) ranges from very high in some species with <3%
211 transmittance to >64% transmittance in other species (Bauer et al. 1998). The highest absorption is typically
212 in evergreen species (Baur et al. 1998). Some evergreen species also contain fluorophores in the cuticular
213 wax, which may convert solar UV irradiation into blue light that can be harnessed for photosynthesis.

214 However, the epicuticular wax *per se* (fatty acyl chains) can also absorb significant amounts of UV-B and
215 thus protect against UV-B (Long et al. 2003). Further, trichome layers also protect against UV-B
216 (Karabourniotis & Bornman 1999). UV-A radiation penetrates deeper into the mesophyll than UV-B in all
217 examined species (Liakoura et al. 2003). Whereas pigment changes in leaves during seasonal changes result
218 in varying degrees of reflectance and transmittance of PAR and IR, there appear to be no seasonal changes
219 in degrees of reflectance and transmittance in the UV region (Yoshimura et al. 2010), despite strong
220 seasonal fluctuations in the leaf concentration of UV-absorbing compounds (Liakoura et al. 2001).

221

222 **4.3 UV-Photochemistry mechanism in and on plant surfaces**

223

224 Photochemical reactions are typically complex. UV radiation can excite various molecules and this may
225 result in a change in molecular orbital occupancy, an increase in energy, and changes in local bonding and
226 charge distribution. Upon return to a lower energy state of the molecules, the released energy or the
227 energy transfer to a neighbouring molecule triggers reactions almost instantaneously. The radiation energy
228 is inversely proportional to the wavelength. Thus, UV-B radiation causes the cleavage of more chemical
229 bonds than does for instance UV-A and PAR. Following this, a multitude of radical reactions may take place
230 and thus greatly increase the quantum yield.

231 From the lab there is plenty of evidence that artificial UV radiation induces an almost instantaneous, i.e.
232 photochemically induced, trace gas emission from plants or plant components, such as carbon-based
233 molecules including CH₄ (McLeod et al. 2008; Keppler et al. 2008; Vigano et al. 2008, 2009; Bruhn et al.
234 2009, 2014; Messenger et al. 2009; Fraser et al. 2015), CO (Tarr et al. 1995; Schade et al. 1999; Brandt et al.
235 2009; Derendorp et al. 2011a; Bruhn et al. 2013), CO₂ (McLeod et al. 2008), and hydrocarbons (McLeod et
236 al. 2008; Derendorp et al. 2011b; Fraser et al. 2015) and nitrogen-based molecules including N₂O (Bruhn et
237 al. 2014b) and NO_x/NO_y (Hari et al. 2003; Raivonen et al. 2006, 2009). Conversely, from the field distinct
238 evidence that natural UV radiation induces an almost instantaneous, i.e. photochemically induced, trace
239 gas emission from plants is far less common, but is documented for CO (Bruhn et al. 2013), N₂O (Bruhn et
240 al. 2014b) and NO_x/NO_y (Hari et al. 2003; Raivonen et al. 2006, 2009). With respect to CH₄ and isoprene,
241 however, there are only indirect indications that natural UV radiation induces an almost instantaneous
242 emission (Keppler et al. 2006; Tiiva et al. 2007).

243

244 **4.3.1 Action spectra**

245 Additional evidence pointing towards direct UV-induced trace gas emission from plants or plant
246 components being an abiotic rather than a biotic process is that in most cases the higher energy UV-B

247 results in higher emission rates than does UV-A at a given irradiance intensity. This is observed in the lab for
 248 plant emission of CH₄ (McLeod et al. 2008; Bruhn et al. 2009), CO (Tarr et al. 1995; Schade et al. 1999;
 249 Bruhn et al. 2013) and N₂O (Bruhn et al. 2014b). Only McLeod et al. (2008) have conducted a detailed
 250 analysis of the action spectrum of direct UV-induced trace gas emission, finding the CH₄ efflux rate from
 251 citrus pectin-impregnated glass fibre sheets to scale linearly with an idealized spectral UV weighting
 252 function. The function weighted CH₄ efflux is an order of magnitude lower for each 80 nm increase in
 253 wavelength. This spectral weighting function differs from other processes where metabolic activity
 254 intrinsically is involved, such as the erythema function (see further discussion in McLeod et al. 2008).
 255

256
 257

258 **Figure 4.1. Relative effect of UV-B to UV-A on trace gas emission from different plant components**
 259 **and surfaces.** The relative effect of UV-B to UV-A is here defined as emission rate in response to UV-B
 260 (312nm) relative to that in response to UV-A (375nm) when adjusted according to irradiance intensity. CH₄:
 261 Pectin, dry (McLeod et al. 2008), Pectin, aq. sol. (Bruhn et al. 2009), *Brassica oleracea* leaves, *Brassica*
 262 *napus* leaves, *Triticum* leaves, *Triticum* ears, and *citrus lemon* fruit peel (Rolsted MMM, Bruhn D, Mikkelsen
 263 TN, Ambus P unpublished); N₂O: Bruhn et al. (2014b); CO: Bruhn et al. (2013). (+) designates that natural
 264 surface wax is present, (-) designates that natural surface wax is either removed mechanically or not present
 265 at all (in the case of *Triticum* spikes). Green = CH₄ from pectin, Dark blue = CH₄ from leaf with nat. wax. Light
 266 blue = CH₄ from leaf without nat. wax, Red = N₂O, Orange=CO.
 267

268 The crude indication of an action spectrum of CH₄ emission from pectin in aqueous solution (Fig 4.1)
 269 resembles the detailed action spectrum of citrus pectin-impregnated glass fibre sheets (Fig 4.1).
 270 Importantly, though, the crude indication of action spectra of trace gas emission appear to be highly
 271 dependent on the precursor as well as the condition of that precursor (Fig 4.1). For example leaves of two
 272 *Brassica* species appear to emit relatively far more CH₄ in response to UV-B than to UV-A, as would be
 273 expected from the results of a single component such as pectin. Furthermore, in the case of the leaves of
 274 the two *Brassica* species, it appears that the removal of leaf surface wax results in a relatively higher effect
 275 of UV-B to that of UV-A in terms of CH₄ emission. This may reflect a combination of UV-A radiation
 276 penetrating deeper into the mesophyll than UV-B in all examined species (Liakoura et al. 2003) when the
 277 wax is intact. However this depends on species (Baur et al. 1998) and perhaps several plant components
 278 (incl. surface wax *per se*) are potential precursors to UV induced trace gas emission (Table 4.1).
 279

280 4.3.2 UV-response functions

281 UV induced trace gas emission is commonly reported to exhibit a near linear response function from both
 282 intact organs/tissues as well as from single plant components (McLeod et al. 2008; Bruhn et al. 2009, 2013,
 283 2014a; Derendorp et al. 2011). This, together with the fact that direct UV induced trace gas emission often
 284 occurs at constant rates over long periods (Bruhn et al. 2009), strongly indicates photochemical reactions
 285 from plentiful precursors.

286
 287 **Fig. 4.2.** Pectin CH₄ production (ng g⁻¹ h⁻¹) as a function of spectrally weighted UV-intensity (W m⁻²). The CH₄
 288 production decays one decade when the spectrum increases 80 nm, e.g. the CH₄ emission is 1 at 300 nm

289 and 0.1 at 380 nm: spectral weighting function from Fig. 1a and data from Table 1 in McLeod et al. (2008).
290 Data is from the UV313 lamp filtered with 125- μ m cellulose diacetate which filters UV wavelengths < 290
291 nm. ER = Emission Rate. Linear regressions functions shown in box. Values are means of three replicates
292 and standard errors are less than symbol size except where visible.

293

294 However, further examination (Fig. 4.2) of the only published dataset on UV-induced pectin-based CH₄
295 productions at very low UV intensities indicates release to be more responsive (by a factor of two) than
296 under higher intensities. The role of self-shading in this context remains to be fully investigated. Thus, even
297 for simple linear responses in UV-induced trace gas emission, there is reason to believe that the underlying
298 photochemical mechanisms are complex. Only Raivonen et al. (2009) have reported on an analysis of the
299 potential linearity of the direct response function of any trace gas (NO_x) emission to natural temporal
300 variation in UV intensity (UV-A).

301

302 **4.3.3 Temperature interactions**

303 In most cases trace gas emission from plant material is also observed in darkness (PAR and UV absent) and
304 with a positive response to temperature increases, although with sensitivities too low to infer underlying
305 abiotic processes (Derendorp et al. 2011a). However, reliable indications are lacking of interactions
306 between the photochemical reactions and temperature with respect to trace gas emission from plant
307 material.

308

309 **4.3.4 [O₂] or O-radicals dependency**

310 It caused quite a surprise (Kirschbaum et al. 2006, 2007) when Keppler et al. (2006) first reported an
311 aerobic plant CH₄ emission in response to solar radiation. In all examined cases of UV induced trace gas
312 emission from plant materials, there is a positive dependency on [O₂] or O-radicals (Table 4.1). Further, this
313 confirms a combination of instantaneous photochemical reactions and subsequent radical reactions in
314 most cases. UV radiation can therefore in theory act as a stimulus via an increased reactive oxygen species
315 (ROS) reaction, and consequently the actual precursor of the emitted gas does not itself need to be a good
316 absorber of UV radiation for the process to occur. However, Lee et al. (2012) provide evidence that photo-
317 oxidation may only be one of several photo degradation processes, as they observed the process occurring
318 in the absence of O₂. They speculated that the direct breakdown of chemical groups such as carboxyl,
319 carbonyl, and methoxyl groups may result in CO₂, CO, and CH₄ release.

320

321 **4.4 Precursors**

322

323 In Table 4.1 we have compiled current knowledge on the potential precursors for trace gas formation in
 324 *direct* response to UV radiation examined and/or suggested in the literature. For the C-based trace gases
 325 there are many structural components, which may act as precursors. In contrast, for the N-based trace
 326 gases the potential precursors appear to be more dependent on surface deposited molecules (Table 4.1).

327

328 **Table 4.1 Potential precursors for trace gas formation in direct response to UV radiation examined**
 329 **and/or suggested in the literature.**

Gas	Suggested source (plant or surface deposit)	Ref	Positive [O ₂] or O-radicals-dependence	Ref
CH₄	Pectin (methyl groups)	1, 2, 3, 4, 5, 6, 7	Yes	2, 7
	Wax	8	Yes	8
	(15-nonacosanone & 2-hexadecanone)			
	Cellulose	3, 4	??	
	Lignin	3, 4	??	
	Methionine	9	??	
	Ascorbic acid	10	Yes	10
CO	Cellulose	11	Yes	12, 13, 14
CO₂	Lignin	15, 16		
C2–C5 hydrocarbons	??	11, 17	Yes	11
N₂O	Wax,	18		
	Mesophyll	18		
	Surface bound N ₂ O	19		
	NO ₃ ⁻	20		
	NH ₄ NO ₃	18, 21	Yes	20, 22, 23
NO_{x,y}	needle surfaces, HNO ₃ or NO ₃ ⁻	24, 25, 26		

330

331 References: 1) Keppler et al. 2006; 2) Keppler et al. 2008; 3) Vigano et al. 2008; 4) Vigano et al. 2009; 5)
 332 McLeod et al. 2008; 6) Bruhn et al. 2009; 7) Messenger et al. 2009; 8) Bruhn et al. 2014a; 9) Bruhn et al.
 333 2012; 10) Althoff et al. 2014; 11) Schade and Crutzen 1999; 12) Tarr et al. 1995; 13) Yonemura et al. 1999;
 334 14) Derendorp et al. 2011; 15) Rozema et al. 1997; 16) Day et al. 2007; 17) Fraser et al. 2015; 18) Bruhn et
 335 al. 2014b; 19) Kim et al. 2010; 20) Rubasinghege & Grassian, 2009; 21) Rubasinghege et al. 2011; 22)
 336 Prasad, 2002; 23) Prasad and Zipf, 2008; 24) Hari et al. 2003; 25) Raivonen et al. 2006; 26) Raivonen et al.
 337 2009. ?? equals unknown.

338
339
340
341
342
343
344
345
346
347

It seems to be the consensus that the polysaccharide pectin is the most important precursor for UV induced plant CH₄ emission due to its content of methyl groups (Keppler et al. 2006, Keppler et al. 2008; Vigano et al. 2008, 2009; McLeod et al. 2008; Bruhn et al. 2009; Messenger et al. 2009; Bloom et al. 2010; Fraser et al. 2015). Whereas we agree that pectin is one of the potential precursors, we are currently not convinced that it necessarily is the most important one. Pectin is laid down in primary plant cell walls. For pectin to be reached by UV irradiation in nature, UV irradiation has to first penetrate the outer surface wax layer naturally occurring on plant organs. In Figure 4.3 we show different pairs of plant organs with natural surface wax or without surface wax, respectively.

348
349
350
351
352

Figure 4.3. Different pairs of plant material with different amounts of surface wax. A+B) Organically grown *Citrus limon* fruit with natural surface wax (A) or with wax mechanically removed (B) by gentle scrubbing with a kitchen sponge. C+D) *Brassica oleracea capitata* f. *alba* leaf with surface wax removed (C) or intact (D). E+F) *Triticum aestivum* "Tähti" (E, with almost no surface wax) and *Triticum aestivum* "Vinjett"

353 (F, with normal surface wax) ears. G+H) Detailed parts of *Triticum aestivum* "Tähti" (G) and *Triticum*
354 *aestivum* "Vinjett" (H) ears. Rolsted MMM, Bruhn D, Mikkelsen TN, Ambus P unpublished. Photos: Rolsted
355 MMM.

356

357 As described in Section 4.2, UV irradiation is to some degree screened at the surface of plant tissues and
358 organs. Thus, an approach to illustrate whether pectin is the most important precursor for CH₄ formation is
359 to measure and compare the UV induced CH₄ production from material of pairs of plant organs with more
360 or less natural UV-screening surface wax (Figure 4.3) as well as in samples of plant leaves with or without
361 the natural wax removed. We did this and found that UV-B induced CH₄ formation in both *citrus limon*
362 peels and *Cydonia oblonga* peels as well as in *Brassica oleracea capitata f. alba* leaves was halved upon
363 removal of the surface wax. This evidently contradicts the notion that in nature pectin is the most
364 important precursor for UV induced CH₄ formation, especially because both *citrus limon* peels and *Cydonia*
365 *oblonga* peels are chosen for industrial pectin extraction due to the high pectin content. Conversely, there
366 was no difference in UV-B induced CH₄ formation between *Triticum aestivum* "Tähti" (with almost no
367 surface wax) ears and *Triticum aestivum* "Vinjett" (with normal surface wax) ears, or between leaves of
368 *Triticum aestivum* when irradiated from either adaxial side (with almost no wax) or from the abaxial side
369 (with much natural surface wax). Removal of surface wax should, in theory, increase UV exposure of the
370 pectin, but in no case did wax removal result in a higher rate of CH₄ formation. Furthermore, we recently
371 demonstrated that the surface wax *per se* is resulting in CH₄ formation upon UV irradiation (Bruhn et al.
372 2014a). In conclusion, we are still far from having a clear understanding of the relative contribution of
373 different precursors in any UV induced trace gas formation and emission. Additionally, given that UV
374 radiation of different wave-lengths reaches different depths in the plant tissue (Liakoura et al. 2003) in a
375 species- dependent manner (Bauer et al. 1998), it seems most likely that each of the different precursors
376 (Table 4.1) are affected differently with respect to wave-lengths of UV irradiation. Consequently, we
377 cannot with any certainty extrapolate an action spectrum for one precursor to that of an entire tissue or
378 organ (Fig. 4.1).

379

380

381

382 5 Upscaling

383 When an unaccounted natural source is discovered there is an urgent demand for extrapolating observed
384 rates to a global scale, so the magnitude of the new source can be put into perspective. However, if

mechanisms are unknown, there is a high risk in an upscaling exercise, because driving forces and controlling factors unintentionally might be ignored, leading to the wrong outcome. On the other hand, if some factors are known, response curves can be constructed and upscaling can be conducted under defined premises, even though there are still unknown factors. Results can then be treated as a platform for understanding and as a contribution to an ongoing knowledge improvement process. Based on current knowledge, we suggest a simple global upscaling and source strength for sunlight-induced emission of the gases CH₄, CO, and N₂O at the plant surface.

5.1 Upscaling of methane, CH₄

The discovery of aerobic CH₄ emissions from plants became breaking news in 2006 (Kepler et al. 2006), mainly because their global upscaling suggested a source strength of 62-236 Tg yr⁻¹, which represented approximately 10–40% of the annual total of methane entering the modern atmosphere, and approximately 30–100% of annual methane entering the preindustrial (0 to 1700 AD) atmosphere (Ferretti et al. 2007). Four independent research groups subsequently revised the global magnitude of this potential CH₄ source by different approaches, and jointly suggested emissions in the lower end compared to the pioneering study by Kepler et al. (2006). Based on carbon stable isotope analysis (Ferretti et al. 2007), standing leaf biomass (Parsons et al. 2006), leaf-mass-based estimation and photosynthesis-based estimation (Kirschbaum et al. 2006), and extrapolation from initially reported chamber measurements (Butenhoff and Khalil 2007), aerobic CH₄ emissions from vegetation were respectively estimated at 0 – 176 Tg yr⁻¹, 42 Tg yr⁻¹, 10 – 60 Tg yr⁻¹ and 20 - 69 Tg yr⁻¹. None of the studies revealed underlying mechanisms for aerobic CH₄ emission.

At the American Geophysical Union (AGU) fall meeting in 2007, three groups presented a major driving factor, UV-radiation, for the aerobic CH₄ emission (Bruhn et al. 2007; Röckmann et al. 2007; Vigano et al. 2007), and in the following years UV generated CH₄ emission was confirmed in several publications (Vigano et al. (2008, 2009); McLeod et al. 2008; Kepler et al. 2008; Bruhn et al 2009; Messenger et al. 2009). McLeod and Kepler (2010) concluded in a review that the proposed formation of CH₄ under aerobic conditions in plants is robust, but the magnitude and significance for the global CH₄ budget remained unresolved.

After the discovery of UV as a driving factor, only one group has tried to upscale aerobic plant generated CH₄; Bloom et al. (2010) provided a putatively low global estimate of 0.2–1.0 Tg y⁻¹ plant-produced CH₄. The upscaling was only based on UV-induced CH₄ emission measured from purified pectin. Bloom et al. (2010)

416 assumed that UV-induced CH₄ emission measured in purified pectin is representative of UV-induced leaf
417 CH₄ emission when taking leaf pectin content into account. However, we believe that there is good
418 evidence in the literature to indicate that this is not the case, since Vigano et al. (2008) for example showed
419 that, at a certain UV irradiation, the CH₄ emission by commercially purified pectin was ca. 80 ng CH₄ g⁻¹ DW
420 h⁻¹, whereas that of dried perennial ryegrass leaf was almost three-fold higher at ca. 200 ng CH₄ g⁻¹ DW h⁻¹
421 – see Bruhn et al (2012) for further discussions. Therefore, the current upscaling by Bloom et al. (2010)
422 must be seen as a preliminary attempt to evaluate global significance from the basis of limited information,
423 and it is important to gain more knowledge for modelling of the UV driven CH₄ from plants at a global level.
424 We suggest that future modelling of the UV driven CH₄ from plants must include data obtained under field
425 conditions with respect to plant growth and development, and exposure to natural sunlight. Such data are
426 currently not available, and therefore methane upscaling is not included in this review.

427

428 **5.2. Upscaling of carbon monoxide, CO**

429 All natural terrestrial direct CO emissions, in the range of 50–200 Tg CO yr⁻¹, have hitherto been ascribed by
430 the IPCC (1995, 2001) to photo-induced CO emission by living plants (cf. Tarr et al. 1995). However, in
431 studies on underlying photo-induced CO emission by living plants (Seiler and Giehl 1977; Seiler et al. 1978),
432 which were incorporated into global CO budgets in the early IPCC assessment reports (IPCC 1995, 2001),
433 the UV component of (sun)light was not considered (Bruhn et al. 2013). Therefore, we still await a proper
434 global estimate of UV radiance induced CO emission by living vegetation. Bruhn et al. (2013) provides the
435 first in situ measurements of ecosystem CO emission by living plants in response to natural solar UV
436 irradiation. Importantly, Bruhn et al. (2013) find that in the studied natural grass field the photo-induced
437 CO emission due to natural solar UV radiation is more than half of the value of that due to total solar
438 spectrum at the Earth's surface. This may imply that the previous global estimate of photo-induced CO
439 emission from living plants of 50–200 Tg CO yr⁻¹ (cf. Tarr et al. 1995) should perhaps be doubled. Thus,
440 future global budgets need to include CO emission caused by natural UV irradiance.

441 Here, we use the results from Bruhn et al. (2013), and estimate the global UV driven CO emissions. The
442 upscaling is based on in situ ecosystem-atmosphere CO exchange measurements from natural vegetation
443 and under ambient UV-B conditions in September and October of 2011 at DTU Risø campus (55°41'12"N,
444 12°05'52"E) in combination with laboratory experiments with artificial UV (Figure 5.1.). For materials and
445 methods see Bruhn et al. (2013), and for the upscaling procedure, see appendices B and C. There is
446 substantial geographical variation in source strength (Fig. 5.1), which is mostly caused by the geographical

447 variation in surface UV radiation intensity, similar to the responsiveness of N₂O (Fig. 5.2). The emission of
 448 CO in response to the UV-component of natural solar radiation was also evident at the ecosystem scale.
 449 When scaled to the global level, the UV-induced emission of CO by vegetation surfaces amounts up to 22 Tg
 450 yr⁻¹, which equals 11-44% of all the natural terrestrial living plant sources hitherto accounted for, which
 451 range between 50 and 200 Tg CO yr⁻¹ (IPCC, 1995, 2001; Tarr et al. 1995).

452
 453 **Figure. 5.1** Estimated annual global CO emissions (mg CO m⁻²) from terrestrial vegetation surfaces induced
 454 by temperature and natural UV-radiation.

456 5.3 Upscaling of Nitrous oxide, N₂O

457 In order to evaluate the global significance of our new discovery of a terrestrial UV-driven N₂O source, we
 458 attempted to scale the processes of temperature- and UV-induced N₂O emission rates by vegetation to the
 459 global level (Fig. 5.2) - for materials and methods see Bruhn et al (2014) and upscaling procedure see
 460 appendices B and C. The upscaling was feasible because the magnitude of measured N₂O emission rates in
 461 response to natural sunlight, including low intensities of UV-radiation ranging from 280-400 nm, was similar
 462 to the magnitude of measured N₂O emission rates in response to high intensities of artificial UV-radiation
 463 within the 309-314 nm narrow range (Bruhn et al. 2014b). The total of these radiation-driven N₂O sources
 464 amounts to 0.65-0.78 Tg yr⁻¹, which equals 7-24% of all the natural terrestrial N₂O sources hitherto
 465 accounted for, which range between 3.3 and 9 Tg N yr⁻¹ (Solomon et al. 2007). There is substantial
 466 geographical variation in the source strength (Fig. 5.2), which is mostly caused by the geographical variation
 467 in surface UV radiation intensity – similar to the responsiveness of CO (Fig. 5.1). Importantly, the irradiance
 468 responses of N₂O emissions across all examined wave length ranges (UV-B, UV-A, PAR) is steepest at low
 469 irradiance intensities (Bruhn et al. 2014b). This intensity-dependent sensitivity is not taken into account in

470 our linear scaling of the UV-induced N₂O emission rates to the global level, and it is therefore likely that we
471 underestimate the N₂O source strength.

472

473

474

475 **Figure 5.2.** Estimated annual global N₂O emissions (mg N₂O m⁻²) from terrestrial vegetation surfaces,
476 induced by temperature and natural UV-radiation.

477 6 Perspectives and Conclusions

478 6.1 Realistic emission rates

479 Despite the many reports on directly UV induced trace gas emission (CH₄, CO, CO₂, C2-C5
480 hydrocarbons/NMVOC, N₂O and NO_x) from plant materials, there are very few studies with replicated
481 measurements of plant trace gas emission in response to natural solar radiation including UV (NO_y,
482 Raivonen et al. 2009; CO, Bruhn et al. 2013; N₂O, Bruhn et al. 2014b). In all three examples there were
483 indications that measured realistic emission rates were substantial compared to those of other
484 emission/uptake processes at the ecosystem level. At this stage it is unfortunately not possible to say
485 anything in general about realistic emission rates from a wider perspective.

486

487 6.2 Future studies

488 From the evidence listed above it becomes apparent that much research is necessary for a more
489 comprehensive understanding of mechanism, precursors and indeed *in situ* emission rates. Therefore we
490 suggest that future experiments include tests of:

- Action spectra and linearity in response function at low UV levels at intact tissues in many more species
- Responses to natural variation in UV intensities in the field
- Effect of deposition of especially N-precursors
- Direct responses to UV after the plants previously have been exposed to variable UV exposures during growth
- Investigation if stomatal conductance has any effects on the UV induced gas emission

6.3 Known gas emission stimulated by UV

It is well established that sunlight and UV in particular stimulate the production of several gases at the surface or near the surface of living plants. Currently there is documented evidence for production of the following gas species: CH₄, CO, CO₂, NMVOCs, NO_x and N₂O. The number of gases produced by UV stimulation is probably greater, but further gas screening studies are needed to assess this. Independent of gas species, the UV-induced gas emission rates documented until now are very low, and as a consequence it is very challenging with respect to equipment and experimental setup to investigate these processes. Most records concern CH₄ production, but there are still many unanswered questions for this gas with regards to dose responses and production under natural conditions. For the other mentioned gases there are even more unanswered questions, nevertheless we have enough information to provide the first attempt at a global budget of UV-induced CO and N₂O emissions based on measurements from natural vegetation under field conditions. The result indicates that UV driven CO production may contribute as much as 11-44% of all the natural terrestrial plant sources. The UV-induced N₂O source equals 7-24% of the natural terrestrial source strength. These global estimates should be regarded as a contribution to an ongoing quantification process, but this high global share emphasizes the urgent need for more work. In order to establish reliable global estimates and enable future predictions, it is apparent that much research is necessary to elucidate mechanisms, precursors, environmental relationships and establishment of relevant and realistic emission rates.

6.4. Perspectives

This newly discovered light-associated aspect of trace gas emission from living vegetation may have significant consequences for our understanding of exchange processes between the global biosphere and atmosphere. It is very likely a global phenomenon occurring on all leaf surfaces exposed to sunlight in both

522 managed and natural ecosystems. Our global estimates for CO and N₂O under the current environmental
523 conditions evidently show that radiation-driven processes are significant natural sources, and this could
524 also be true for the other gases. An important feature is that the gas production is occurring at or just
525 under the leaf surface, resulting in periodic high gas concentration within the boundary layer surrounding
526 the leaves. This could for instance reduce the gas uptake of ozone in leaves since CO accelerates the
527 reaction of O₃ with ethylene (Horie and Moortgat, 1998), a process that has so far not been considered in
528 ozone effect research. Through geological eras, the radiation-driven greenhouse gas (direct: CO₂, CH₄, N₂O
529 and indirect: CO) impact must have fluctuated with UV radiation and other processes producing or
530 consuming greenhouse gases (e.g. soil respiration, denitrification and methanogens in wetlands, and
531 methane oxidation in upland soils), and therefore the development of climate on Earth.

532 **Acknowledgements.** We are thankful to Prof Ary Hoffmann for reading the manuscript and providing
533 helpful suggestions for corrections and NordGen (The Nordic Genetic Resource Center) for providing seeds
534 for the experiments. This work was supported by the FP7 EU Eclairé project.

535

536 References

- 537 Althoff F, Jugold A, Keppler F (2010) Methane formation by oxidation of ascorbic acid using iron minerals
538 and hydrogen peroxide. *Chemosphere* 80: 286–292
- 539 Altimir N, Vesala T, Keronen P, Kulmala M, Hari, P. (2002). Methodology for direct field measurements of
540 ozone flux to foliage with shoot chambers. *Atmospheric Environment* 36(1): 19-29.
- 541 Aphalo PJ, Albert A, McLeod A, Heikkilä A, Gómez I, Figueroa FL, Robson TM, Strid Å (2012) Manipulating
542 UV radiation. In: PJ Aphalo, A Albert, LO Björn, A McLeod, TM Robson, E Rosenqvist (eds.) *Beyond the
543 visible: A handbook of best practice in plant UV photobiology*. COST Action FA0906 UV4growth. Helsinki,
544 University of Helsinki, 176 pp.
- 545 Baur P, Stulle K, Schonherr J, Uhlig B (1998) Absorption of UV-B to blue light radiation by leaf cuticles of
546 selected crop plants. *Gartenbauwissenschaft* 63: 145-152.
- 547 Beer C, Reichstein M, Tomelleri E, Ciais P, Jung M, Carvalhais N, Rödenbeck C, Arain MA, Baldocchi D, Bonan
548 GB, Bondeau A, Cescatti A, Lasslop G, Lindroth A, Lomas M, Luyssaert S, Margolis H, Oleson KW, Rouspard
549 O, Veenendaal E, Viovy N, Williams C, Woodward FI, Papale D (2010) Terrestrial gross carbon uptake: global
550 distribution and covariation with climate. *Science* 329:834–838

551 Björn LO (1996) Effects of ozone depletion and increased UV-B on terrestrial ecosystems, *Int. J. Environ.*
552 *Studies* 51: 217–243.

553 Björn LO, McLeod A, Aphalo PJ, Albert A, Lindfors AV, Heikkilä A, Kolarž P, Ylianttila L, Zipoli G, Grifoni D,
554 Huovinen P, Gómez I, Figueroa FL (2012). Quantifying UV radiation. In: PJ Aphalo, A Albert, LO Björn, A
555 McLeod, TM Robson, E Rosenqvist (eds.) *Beyond the visible: A handbook of best practice in plant UV*
556 *photobiology*. COST Action FA0906 UV4growth. Helsinki, University of Helsinki, 176 pp.

557 Bloom AA, Lee-Taylor J, Madronich S, Messinger DJ, Palmer PI, Reay DS, McLeod AR (2010) Global methane
558 emission estimates from ultraviolet irradiation of terrestrial plant foliage. *New Phytol* 187: 417–425

559 Boucher O, Randall D, Artaxo P, Bretherton C, Feingold G, Forster P, Kerminen V-M, Kondo Y, Liao H,
560 Lohmann U, Rasch P, Satheesh SK, Sherwood S, Stevens B and Zhang XY, 2013: Clouds and Aerosols. In:
561 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment*
562 *Report of the Intergovernmental Panel on Climate Change [Stocker, TF, Qin D, Plattner G-K, Tignor M, Allen*
563 *SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds.)]. Cambridge University Press, Cambridge,*
564 *United Kingdom and New York, NY, USA.*

565 Bowling DR, Miller JB, Rhodes ME, Burns SP, Monson RK, Baer D (2009). Soil, plant, and transport influences
566 on methane in a subalpine forest under high ultraviolet irradiance. *Biogeosciences* 6: 1311-1324.

567 Brown MJ, Parker GG, Posner NE (1994) A survey of ultraviolet-B radiation in forests. *Journal of Ecology* 82:
568 843-854.

569 Bruhn D, Mikkelsen TN, Ambus P (2007) Aerobic Emission of Methane by Terrestrial Plant Material in
570 Response to UV- irradiance. *Eos Trans. AGU*, 88(52), Fall Meet. Suppl., Abstract B53A-0940.

571 Bruhn D, Mikkelsen TN, Obro J, Willats WGT, Ambus P (2009). Effects of temperature, ultraviolet radiation
572 and pectin methyl esterase on aerobic methane release from plant material. *Plant Biol.* 11: 43-48.

573 Bruhn D, Mikkelsen TN, Rolsted MMM, Egsgaard H, Ambus P (2014a) Leaf surface wax is a source of plant
574 methane formation under UV radiation and in the presence of oxygen. *Plant Biol.* 16: 512-516.

575 Bruhn D, Moller IM, Mikkelsen TN, Ambus P (2012) Terrestrial plant methane production and emission.
576 *Physiol. Plant.* 144: 201-209.

577 Bruhn D, Albert KR, Mikkelsen TN, Ambus P (2014b). UV-induced N₂O emission from plants. *Atmospheric*
578 *Environment*, 99: 206-214

579 Bruhn D, Albert KR, Mikkelsen TN, Ambus P (2013) UV-induced carbon monoxide emission from living
580 vegetation. *Biogeosciences*, 10: 7877–7882

581 Caldwell MM (1981) Plant response to solar ultraviolet-B radiation. P. 170-186. In Lange et al. (ed)
582 Physiological plant ecology. Springer-Verlag, Berlin.

583 Caldwell MM, Searles PS, Flint SD, Barnes PW (1999). Terrestrial ecosystem responses to solar UV-B
584 radiation mediated by vegeta-tion, microbes and abiotic chemistry, in: M.C. Press, J.D. Scholes, M.G. Barker
585 (Eds.), Physiological Plant Ecology, Blackwell Science, Oxford, 1999, pp. 241–262.

586 Caldwell MM, Teramura AH, Tevini M, Bornman JF, Bjorn LO, Kulandaivelu G (1995) Effects of increased
587 solar ultraviolet radiation on terrestrial plants. *Ambio* 24: 166–173.

588 Cen Y P and Bornman J F (1993) The effect of exposure to enhanced UV-B radiation on the penetration of
589 monochromatic and polychromatic UV-B radiation in leaves of *Brassica-napus*, *Physiol. Plant.*, 87: 249–255.

590 Ciais P, Sabine C, Bala G, Bopp L, Brovkin V, Canadell J, Chhabra A, DeFries R, Galloway J, Heimann M, Jones
591 C, Le Quéré C, Myneni, Piao S, Thornton P (2013). Carbon and Other Biogeochemical Cycles. In: Climate
592 Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of
593 the Intergovernmental Panel on Climate Change [Stocker, TF, Qin D, Plattner G-K, Tignor M, Allen SK,
594 Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds.)]. Cambridge University Press, Cambridge, United
595 Kingdom and New York, NY, USA.

596 Day TA, Zhang ET, Ruhland CT (2007) Effects of ultraviolet radiation accelerates mass and lignin loss of
597 *Larrea tridentate* litter in the Sonoran Desert. *Plant Ecol* 193: 185-194.

598 Deckmyn GE, Cayenberghs E, Ceulemans (2001) UV-B and PAR in single and mixed canopies grown under
599 different UV-B exclusions in field. *Plant Ecol* 154: 125-133.

600 Derendorp L, Holzinger R, Röckmann T, (2011b). UV-induced emissions of C2-C5 hydrocarbons from leaf
601 litter. *Environ. Chem.* 8: 602-611.

602 Derendorp L, Quist JB, Holzinger R, Röckmann T (2011a) Emissions of H₂ and CO from leaf litter of
603 *Sequoiadendron giganteum* and their dependence on UV radiation and temperature. *Atmos Environ* 45:
604 7520-7524.

605 Dueck T A, de Visser R, Poorter H, Persijn S, Gorissen A, de Visser W, Schapendonk A, Verhagen J, Snel J,
606 Harren FJM, Ngai AKY, Verstappen F, Bouwmeester H, Voeselek LACJ, van der Werf A (2007). No evidence
607 for substantial aerobic methane emission by terrestrial plants: a ¹³C-labelling approach. *New Phytologist*
608 175: 29-35.

609 EPA (2010). Methane and Nitrous Oxide Emissions From Natural Sources. Office of Atmospheric Programs
610 (6207J), Washington, DC 20460, EPA 430-R-10-001.

611 Ferretti DF, Miller JB, White JWC, Lassey KR, Lowe DC, Etheridge DM (2007) Stable isotopes provide revised
612 global limits of aerobic methane emissions from plants. *Atmospheric Chemistry and Physics* 7: 237–241.

613 Fraser WT, Blei E, Fry SC, Newman MF, Reay DS, Smith KA, Mcleod AR, 2015. Emission of methane, carbon
614 monoxide, carbon dioxide and short-chain hydrocarbons from vegetation foliage under ultraviolet
615 irradiation. *Plant Cell Environ.* 38: 980-989.

616 Grant RH (1997) Partitioning of biologically active radiation in plant canopies. *Int J Biometeorol* 40: 26-40.

617 Hari P, Raivonen M, Vesala T, Munger JW, Pilegaard K, Kulmala M (2003) Ultraviolet light and leaf emission
618 of NO_x. *Nature* 422: 134.

619 Hartmann DL, Klein Tank AMG, Rusticucci M, Alexander LV, Brönnimann S, Charabi Y, Dentener FJ,
620 Dlugokencky EJ, Easterling DR, Kaplan A, Soden BJ, Thorne PW, Wild M, Zhai PM (2013). Observations:
621 Atmosphere and Surface. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working*
622 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, TF, Qin
623 D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds.)]. Cambridge
624 University Press, Cambridge, United Kingdom and New York, NY, USA.

625 Horie O, Moortgat GK (1998) The effect of the addition of CO on the reaction of ozone with ethene.
626 *Chemical Physics Letters*, 288, 464–472.

627 IPCC: Climate Change 2001: The Scientific Basis, Contribution of Working Group I to the Third Assessment
 628 Report of the Intergovernmental Panel on Climate Change, edited by: Houghton J T, Ding Y, Griggs D J,
 629 Noguer M, van der Linden P J, Dai X, Maskell K, Johnson C A. Cambridge University Press, Cambridge,
 630 United Kingdom and New York, NY, USA, 881 pp.

631 IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth
 632 Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, TF, Qin D, Plattner G-K,
 633 Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds.)]. Cambridge University Press,
 634 Cambridge, United Kingdom and New York, NY, USA.

635 Isaksen I, Granier C, Myhre G, Berntsen T, Dalsren S, Gauss M, Klimont Z, Benestad R, Bousquet P, Collins
 636 W, Cox T, Eyring V, Fowler D, Fuzzi S, Jckel P, Laj P, Lohmann U, Maione M, Monks P, Prevot A, Raes F,
 637 Richter A, Rognerud B, Schulz M, Shindell D, Stevenson D, Storelvmo T, Wang WC, van Weele M, Wild M,
 638 Wuebbles D (2009) Atmospheric composition change: climate chemistry interactions. *Atmos. Environ.* 43:
 639 5138-5192.

640 Jacobs JF, Koper GJM, Ursem WNJ (2007) UV protective coatings: a botanical approach, *Prog. Org. Coat.*, 58:
 641 166–171.

642 Karabourniotis G, Bornman JF (1999) Penetration of UV-A, UV-B and blue light through the leaf trichome
 643 layers of two xeromorphic plants, olive and oak, measured by optical fibre microprobes. *Physiologia*
 644 *Plantarum* 105: 655-661.

645 Keppler F, Boros M, Frankenberg C, Lelieveld J, McLeod A, Pirttila AM, Rockmann T, Schnitzler JP (2009)
 646 Methane formation in aerobic environments. *Environ. Chem.* 6: 459-465.

647 Keppler F, Hamilton JTG, Brass M, Rockmann T (2006). Methane emissions from terrestrial plants under
 648 aerobic conditions. *Nature* 439: 187-191.

649 Keppler F, Hamilton JTG, McRoberts WC, Vigano I, Brass M, Rockmann T (2008) Methoxyl groups of plant
 650 pectin as a precursor of atmospheric methane: evidence from deuterium labelling studies. *New Phytol.* 178:
 651 808-814.

652 Khalil MAK and Rasmussen RA (1990) The global cycle of carbon monoxide: Trends and mass balance,
 653 *Chemosphere*, 20: 227-242.

654 Kim KH, Watanabe K, Menzel D, Freund H-J (2010) UV photo-dissociation and photodesorption of N₂O on
 655 Ag(111). *J. Phys. Condens. Matter* 22: 084012.

656 Kirschbaum MUF, Bruhn D, Etheridge DM, Evans JR, Farquhar GD, Gifford RM, Paul KI, Winters AJ (2006) A
657 comment on the quantitative significance of aerobic methane release by plants. *Functional Plant Biology*
658 33: 521–530.

659 Kirschbaum MUF, Niinemets Ü, Bruhn D, Wintners AJ (2007) How important is aerobic methane release by
660 plants? *Funct Plant Sci Biotech* 1: 138–145

661 de Klein C, Harvey M (2015) Nitrous Oxide Chamber Methodology Guidelines. Version 1.1 ISBN 978-0-478-
662 40585-9 (online). Ministry for Primary Industries, Wellington 6140, New Zealand

663 Lee H, Rahn T, Throop HL, (2012) An accounting of C-based trace gas release during abiotic plant litter
664 degradation. *Global Change Biology* 18: 1185–1195.

665 Liakoura V, Manetas Y, Karabouniotis G (2001) Seasonal fluctuations in the concentration of UV-absorbing
666 compounds in leaves of some Mediterranean plants under field conditions. *Physiologia Plantarum* 111:
667 491-500.

668 Liakoura V, Bornmann JF, Karabouniotis G (2003) The ability of abaxial and adaxial epidermis of sun and
669 shade leaves to attenuate UV-A and UV-B radiation in relation to the UV absorbing capacity of the whole
670 leaf methanolic extracts, *Physiol. Plant.*, 117: 33–43.

671 Long LM, Patel HP, Cory WC, Stapleton AE (2003) The maize epicuticular wax layer provides UV protection.
672 *Functional Plant Biology* 30: 75-81.

673 Masson-Delmotte V, Schulz M, Abe-Ouchi A, Beer J, Ganopolski A, González Rouco JF, Jansen E, Lambeck K,
674 Luterbacher J, Naish T, Osborn T, Otto-Bliesner B, Quinn T, Ramesh R, Rojas M, Shao X, Timmermann A
675 (2013) Information from Paleoclimate Archives. In: *Climate Change 2013: The Physical Science Basis.*
676 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
677 *Change In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth*
678 *Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, TF, Qin D, Plattner G-K,*
679 *Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds.)]. Cambridge University Press,*
680 *Cambridge, United Kingdom and New York, NY, USA.*

681 McLeod AR, Fry SC, Loake GJ, Messenger DJ, Reay DS, Smith KA, Yun BW (2008). Ultraviolet radiation drives
682 methane emissions from terrestrial plant pectins. *New Phytol.* 180: 124-132.

683 McLeod A, Keppler F (2010) Vegetation. In: Reay D, Smith P, van Amstel A (eds.) *Methane and Climate*
684 *Change.* Earthscan, London and Washington, DC, pp 74–96.

685 Messenger DJ, McLeod AR, Fry SC (2009) The role of ultraviolet radiation, photosensitizers, reactive oxygen
686 species and ester groups in mechanisms of methane formation from pectin. *Plant Cell Environ* 32: 1–9

687 Mikkelsen TN, Bruhn D, Ambus P, Larsen KS, Ibrom A, Pilegaard K (2011) Is methane released from the
688 forest canopy? *Iforest-Biogeosci. For.* 4: 200-204.

689 Mikkelsen TN, Ro-Poulsen H (2002) In situ autumn ozone fumigation of mature Norway spruce - effects on
690 net photosynthesis. *Phyton* 42: 97–104.

691 Myhre G, Shindell D., Bréon F.-M., Collins W., Fuglestad J., Huang J., Koch D., Lamarque J.-F., Lee D.,
692 Mendoza B., Nakajima T., Robock A., Stephens G., Takemura T. and Zhang H., 2013: Anthropogenic and
693 Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working*
694 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, TF, Qin
695 D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds.)]. Cambridge
696 University Press, Cambridge, United Kingdom and New York, NY, USA.

697 Parsons AJ, Newton PCD, Clark H, Kelliher FM. 2006. Scaling methane emissions from vegetation. *Trends in*
698 *Ecology and Evolution* 21: 423–424.

699 Potter C S, Klooster S A, Chatfield RB (1996) Consumption and production of carbon monoxide in soils: a
700 global model analysis of spatial and seasonal variation, *Chemosphere*, 33: 1175–1193.

701 Prasad SS, Zipf EC (2008) Atmospheric production of nitrous oxide from excited ozone and its potentially
702 important implications for global change studies. *J. Geophys. Res. Atmos.* 113: D15307.

703 Raivonen M, Bonn B, Sanz MJ, Vesala T, Kulmala M, Hari P (2006) UV-induced NO_y emissions from Scots
704 pine: could they originate from photolysis of deposited HNO₃? *Atmospheric Environment* 40: 6201–6213.

705 Raivonen M, Vesala T, Pirjola L, Altimir N, Keronen P, Kulmala M, Hari P (2009) Compensation point of NO_x
706 exchange: Net result of NO_x consumption and production. *Agricultural and Forest Meteorology* 149: 1073–
707 1081.

708 Rosenqvist E, Figueroa FL, Gómez I, Aphalo PJ (2012). Plant growing conditions. In: PJ Aphalo, A Albert, LO
709 Björn, A McLeod, TM Robson, E Rosenqvist (eds.) *Beyond the visible: A handbook of best practice in plant*
710 *UV photobiology. COST Action FA0906 UV4growth. Helsinki, University of Helsinki, 176 pp.*

711 Röckmann T, Vigano I, Holzinger R, van Weelden H, Keppler F (2007) News about methane emission from
712 plant matter. *Eos Trans. AGU*, 88(52), Fall Meet. Suppl., Abstract B51F-05

713 Rozema J, van de Staaij J, Björn LO, Caldwell M (1997) UV-B as an environmental factor in plant life: stress
714 and regulation, *Trends Ecol. Evol.* 12: 22–28.

715 Rubasinghege G, Grassian VH (2009) Photochemistry of adsorbed nitrate on aluminum oxide particle
716 surfaces. *J. Phys. Chem. A* 113: 7818-7825.

717 Rubasinghege G, Spak SN, Stainer CO, Carmichael GR, Grassian VH (2011) Abiotic mechanism for the
718 formation of atmospheric nitrous oxide from ammonium nitrate. *Environ. Sci. Technol.* 45: 2691-2697.

719 Schade GW, Hofmann R-F, Crutzen PJ (1999) CO emissions from degrading plant matter, (I) Measurements,
720 *Tellus B*, 51: 889–908. 1999a.

721 Schade, GW, Crutzen PJ (1999): CO emissions from degrading plant matter (II), Estimate of a global source
722 strength, *Tellus B*, 51: 909–918.

723 Shindell DT, Faluvegi G, Koch DM, Schmidt GA, Unger N, Bauer SE (2009) Improved attribution of climate
724 forcing to emissions. *Science*, 326: 716–718.

725 Seiler W, Giehl H (1977) Influence of plants on the atmospheric carbon monoxide, *Geophys. Res. Lett.*, 8:
726 329–332.

727 Seiler W, Giehl H, Bunse G (1978) The influence of plants on atmospheric carbon monoxide and dinitrogen
728 oxide, *Pure Appl. Geophys.*, 116: 439–451.

729 Shulski MD, Walter-Shea EA, Hubbard KG, Yuen GY, Horst G (2004) Penetration of photosynthetic active
730 radiation and ultra violet radiation into Alfalfa and Tall Fescue canopies. *Agronomy Journal* 96: 1562-1571.

731 Smeets CJPP, Holzinger R, Vigano I, Goldstein AH, Rockmann T (2009) Eddy covariance methane
732 measurements at a Ponderosa pine plantation in California. *Atmos. Chem. Phys.* 9: 8365-8375.

733 Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds.) (2007)
734 Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on
735 Climate Change, 2007, Cambridge Univ. Press.

736 Stocker TF, Qin D, Plattner G-K, Alexander LV, Allen SK, Bindoff NL, Bréon F-M, Church JA, Cubasch U, Emori
737 S, Forster P, Friedlingstein P, Gillett N, Gregory JM, Hartmann DL, Jansen E, Kirtman B, Knutti R, Krishna
738 Kumar K, Lemke P, Marotzke J, Masson-Delmotte V, Meehl GA, Mokhov II, Piao S, Ramaswamy V, Randall D,
739 Rhein M, Rojas M, Sabine C, Shindell D, Talley LD, Vaughan DG Xie S-P, 2013: Technical Summary. In:
740 Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment
741 Report of the Intergovernmental Panel on Climate Change [Stocker, TF, Qin D, Plattner G-K, Tignor M, Allen
742 SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds.)]. Cambridge University Press, Cambridge,
743 United Kingdom and New York, NY, USA.

744 Sundqvist E, Crill P, Molder M, Vestin P Lindroth A (2012). "Atmospheric methane removal by boreal
745 plants." *Geophysical Research Letters* 39: L21806.

746 Tarr M A, Miller W L, Zapp RG (1995) Direct carbon monoxide photoproduction from plant matter, *J.*
747 *Geophys. Res.*, 100: 11403–11413,

748 Tiiva P, Rinnan R, Faubert P, Rasanen J, Holopainen T, Kyrö E and Holopainen JK (2007), Isoprene emission
749 from a subarctic peatland under enhanced UV-B radiation, *New Phytol.* 176: 346–355.

750 Ueta I, Mizuguchi A, Tani K, Kawakubo S Saito Y (2013) Rapid Temperature-Programmed Separation of
751 Carbon Monoxide and Carbon Dioxide on a Packed Capillary Column in Gas Chromatography: Application to
752 the Evaluation of Photocatalytic Activity of TiO₂. *Analytical Sciences* 29: 673-676.

753 Vigano I, Holzinger R & Röckman T (2007) The isotope signature of methane emitted from plant matter
754 upon irradiation with UV light. *Eos Trans. AGU*, 88(52), Fall Meet. Suppl., Abstract B53A-0938

755 Vigano I, van Weelden H, Holzinger R, Keepler F, Röckmann T (2008) Effect of UV radiation and temperature
756 on the emission of methane from plant biomass and structural components. *Biogeosciences* 5: 937–947

757 Vigano I, Röckmann T, Holzinger R, van Dijk A, Keppler F, Greule M, Brand WA, Geilmann H, van Weelden H
758 (2009). The stable isotope signature emitted from plant material under UV irradiation. *Atmos Environ* 43:
759 5637–5646.

760 Vigano I, van Weelden H, Holzinger R, Keppler F, McLeod A Röckmann T (2008). Effect of UV radiation and
761 temperature on the emission of methane from plant biomass and structural components. *Biogeosciences* 5:
762 937-947.

763 Yoshimura H, Zhu H, Wu Y, Ma R (2010) Spectral properties of plant leaves pertaining to urban landscape
764 design of broad-spectrum solar ultraviolet radiation reduction. *Int J Biometeorol* 54: 179 - 191.

765 Worden HM, Deeter MN, Frankenberg C, George M, Nichitau F, Worden J, Aben I, Bowman KW, Clerbaux C,
766 Coheur PF, de Laat ATJ, Detweiler R, Drummond JR, Edwards D P, Gille JC, Hurtmans D, Luo M. Martínez-
767 Alonso S, Massie S, Pfister G, Warner JX (2013) Decadal record of satellite carbon monoxide observations,
768 *Atmos. Chem. Phys.*, 13: 837-850.

769 Yurganov L, McMillan W, Grechko E, Dzhola A (2010) Analysis of global and regional CO burdens measured
770 from space between 2000 and 2009 and validated by ground-based solar tracking spectrometers. *Atmos.*
771 *Chem. Phys.* 10: 3479–3494.

772

773

774 **Appendix A**

775 ***Methods & techniques***

776 This appendix expands 3 methods & techniques in the text with respect to environmental factors and other
777 important issues related to measurements of UV induced gases.

778

779 **Temperature**

780 Particularly challenging is temperature control inside confinements made of transparent materials and at
781 the same time exposed to intense irradiation by lamps or natural sunlight that may lead to severe scorching
782 of plants. Under laboratory conditions, experiments may be run in well vented and temperature controlled
783 growth cabinets (Bruhn et al. 2009) or enclosures equipped with heating tape (Vigano et al. 2008) to
784 maintain stable temperature conditions. Direct temperature control of enclosures may include simple and
785 inexpensive means such as ice blocks (M. Drösler, pers. comm.) or more advanced applications in the form
786 of Peltier cooling technology (Mikkelsen and Ro-Poulsen, 2002; Bruhn et al. 2014; Sundqvist et al. 2012).
787 Temperature can be determined directly at the leaf surface with a micro-thermocouple attached to the
788 material (Keppler et al. 2006) or air temperature detected by thermocouples or conventional
789 thermometers situated in the enclosure. Exterior surface temperature of enclosures can be measured by
790 using a heat conducting steel probe connected to a high precision temperature meter (Bruhn et al. 2014).

791

792 **Chamber material**

793 Chambers should be made of materials that allow transmittance of UV-radiation without filtering.
794 Commonly used materials in transparent chamber or plant cuvettes include UV-transparent synthetic
795 quartz-glass, tradename Suprasil®, that offers optimal UV-transmissions (Vigano et al. 2008; Rosenqvist et
796 al. 2012; Bruhn et al. 2014), alternatively UV-transparent acrylic materials (Rosenqvist et al. 2012; Bruhn et
797 al. 2014) can be used. Controlled transmittance of UV in experimental setups can be achieved by the
798 application of filters to reduce or filter out specific UV-wavelengths reaching surfaces being investigated.
799 Many commercial acrylic materials, with trade names such as e.g. Plexiglas or Perspex, will attenuate UV
800 penetration and can be deployed to manipulate UV intensity (Bruhn et al. 2014). For specific and controlled
801 filtering of UV radiation, various filters can be applied either at the light source or covering the enclosure
802 windows; a comprehensive review of UV manipulation is given by Aphalo et al. (2012).

803 **Figure A1.** Plant leaves inserted in UV-B transparent vials. Note vials without leaves are used as blank
 804 controls.

805 **Surface reactions and reactive species**

806 Synthetic soft plastic and rubber materials used in growth cabinets such as hoses, tubes, pots, sealants and
 807 wire insulators, provide potential complications if exposed in experimental setups to study UV-induced gas
 808 emissions. Firstly, these materials may release phytotoxic compounds, leading to plant growth problems or
 809 plant death; for a review see Rosenqvist et al. (2012). Secondly, photochemical reactions on the surface of
 810 synthetic materials when exposed to UV-radiation can produce gases like methane (D. Bruhn, unpublished)
 811 or N₂O (Bruhn et al. 2014) that may confound experimental results. For this reason, it is strongly
 812 recommended to include empty/blank controls in the experimental protocol (Bruhn et al. 2014; Sundqvist
 813 et al. 2012; Figure A1). Presence of synthetic materials in the experimental units should be minimized, and
 814 materials shielded with (e.g.) PTFE replaced with inert materials (glass, metal) where appropriate or pre-
 815 conditioned by heating (Sundqvist et al. 2012).

816

817 **Analysis of gas mixing ratios**

818 Analysis of mixing ratios of target gases in the enclosures can be achieved principally by two different
 819 approaches. These are *i*) manual grab sampling by syringe where a subsample of enclosure headspace is
 820 transferred to the analyzer or a storage vial for subsequent analysis; incubation may also take place in vials
 821 that can be mounted directly in the analytical unit such as a GC-autosampler, avoiding the need for manual
 822 sample transfer (Bruhn et al, 2014). Alternatively, *ii*) the headspace gas concentrations can be observed in

823 real-time where the test unit is connected to an appropriate gas analyzer in a sealed gas loop for
824 continuous or cyclic analysis (Sundqvist et al. 2012).

825

826 Gas-chromatography: Conventional GC-instrumentation equipped with Flame-Ionization-Detection [FID] for
827 CH₄ (e.g. Vigano et al. 2008), methanizer-FID for CO and CO₂ (Ueta et al. 2013) and Electron-Capture-
828 Detection [ECD] for N₂O (e.g. Bruhn et al. 2014) are applicable for grab sample analysis in order to cross-
829 check the optical techniques, and where experiments with small vials prevent measurements with optical
830 systems that require additional sample volume. The reproducibility with GC-analysis is typically ±10 ppb
831 although the micro-GC system tested by Ueta et al. (2013) for combined CO and CO₂ analysis exhibited
832 detection limits of 3 to 5 ppm. Trace-gas GC analysis at ambient concentrations require sample volumes of
833 typically 0.5-1 ml.

834

835 Laser spectroscopy: During the last couple of decades laser spectrometers for sensitive, accurate and fast
836 analysis of air trace gas constituents have become available at affordable pricing. In their work with UV-
837 induced CH₄ dynamics, Vigano et al. (2008) and Sundqvist et al. (2102) used an off-axis integrated cavity
838 output spectrometer (Los Gatos Inc.) for real-time monitoring of CH₄ mixing ratios. With use of laser
839 spectroscopy, it is necessary to pay attention to cross-interference from other gas species; Vigano et al.
840 (2008) verified this for plant emission of abundant methanol (CH₃OH). For studies on UV-induced emissions
841 of carbon monoxide (Bruhn et al. 2013) and nitrous oxide (Bruhn et al. 2014), a Los Gatos laser N₂O/CO
842 spectrometer was applied following proper correction for cross-sensitivity with water and direct cross-
843 interference between CO and N₂O. The sensitivity of laser spectrometers is several fold higher compared
844 with GC analysis, and allows reproducibility in the range of ±1 ppb.

845 Stable isotope analysis: GC combined with stable isotope (SI) analysis provides a powerful tool to study
846 source partitioning and reaction pathways of trace gases emitted from surfaces. Isotope-ratio-mass-
847 spectrometry (IRMS) in combination with proper pre-concentration (e.g. cryo-trapping) and
848 chromatographic separation of analytical compounds has been used for studying carbon (¹³C), hydrogen
849 (deuterium; D) in CH₄ (Keppler et al. 2006; 2008) and nitrogen (¹⁵N) in N₂O (Bruhn et al. 2014). The
850 reproducibility of gas mixing ratios by GC-IRMS is diminished (±20-30 ppb at ambient concentrations)
851 compared with conventional GC analysis and the sample amount required for proper analysis is in the
852 range of tens of milliliters. Dueck et al. (2007) analyzed the concentration of ¹³C-methane in CH₄ emitted
853 from fully ¹³C-labeled plant material using photo-acoustic spectroscopy in combination with a continuous-

854 wave, optical parametric oscillator (OPO) and reported a detection limit of 3 ppb. Whereas the work by
855 Dueck et al. (2007) demonstrated only negligible emissions of CH₄ based on the spectrometric method,
856 later work by Vigano et al. (2008) with the same plant material showed a contrasting result with significant
857 emissions of ¹³C-CH₄, supposedly due to different analytical sensitivities (Vigano et al. 2008).

858 SI analysis encompasses recognition of the isotopic composition in gases emitted from materials with
859 isotopic abundances at natural levels (e.g. Keppler et al. 2006) as well as from isotopically enriched
860 materials (e.g. Bruhn et al. 2014). Isotopic variations arise from mass-dependent isotope fractionation in
861 biological and chemical processes, and natural abundance analysis of the trace gases, may add information
862 about the nature and origin of precursors. Application of the rare (heavy) isotope is valuable not only for
863 revealing information on specific precursor substances but also for providing a tool to study consumption
864 processes.

865 The recent development of isotopic laser spectrometers (e.g. instruments offered by Picarro, Aerodyne, Los
866 Gatos) provides new opportunities to investigate mechanisms and processes in UV-induced trace gas
867 emissions. However, so far no work taking advantage of these instruments has been reported in literature.

868 **Light sources**

869 Work on UV-induced trace gas emissions inevitably requires selection of a proper light source and
870 establishment of associated irradiation intensities, wavelengths and action spectra. The selection of a light
871 source is application driven and depends on the requirements imposed by the study. The main
872 requirements concern the intensity and spectral distribution of radiant output of the lamp. The geometry
873 of the setup, including the source-target-distance and area of exposure, sets certain limits not only on the
874 light source but also on characteristics of the monochromator if this is applied (Aphalo et al. 2012).

875 In this context, it is important to stress that the current review addresses works investigating the direct
876 photolytic effect of UV-light exposure for plant-derived trace gas emissions. In this sense, translocation
877 studies where plants are grown under natural or controlled conditions with attenuated or enhanced UV-
878 exposure and subsequently examined for historical UV effects are not considered.

879 It is beyond the limits of this manuscript to present a detailed review and recommendation on selection of
880 proper light source equipment for UV studies. Instead, the reader is referred to comprehensive reviews on
881 usage of artificial light sources in UV photobiology given by Aphalo et al. (2012), and UV quantification
882 reviewed by Björn et al. (2012). A number of potential light sources can be selected for UV work, either as

883 single light sources, or more often in combination to achieve desired optical conditions. A brief list is shown
884 below; for a detailed discussion we refer to Aphalo et al. (2012).

885 Fluorescent lamps and tubes are low pressure mercury vapor lamps that emit radiation at specific spectral
886 lines, mostly in the UV region of the spectrum

887 Xenon arc lamps are specialized light sources that produce intense visible and UV radiation. High intensity
888 water-cooled deuterium lamps (150 W) have a fairly flat radiant intensity curve in the UV-B region that is
889 appropriate for mechanistic plant UV photobiology studies.

890 Spectrographs composed of a light source and a monochromator may be used in applications requiring
891 spectrally-resolved UV radiation exposure of biological specimens. Lasers usually produce very narrow and
892 intense beams of monochromatic light. For the purpose of UV photobiology, tuneable optical parametric
893 oscillator (OPO) pump lasers (pump wavelength 355 nm) are especially useful.

894 Specific experimental setups for studying UV-effects on plant gas emissions are described in detail in
895 literature cited above, and illustrate the complexity and experimental precautions associated with such
896 studies. As an example, Vigano et al. (2008) used 6 types of lamps, one PAR lamp, four UV-A and UV-B
897 lamps, and one UV-C lamp. The UV content (UV-A and UV-B separately) was determined with a Waldmann
898 UV meter calibrated for each individual UV lamp, except for the UV-C lamp. These authors did not report on
899 the action spectrum for CH₄ release from biomass upon UV irradiation, and the UV strength was reported
900 as the non-weighted integral over the UV-A range (400–320 nm), UV-B range (320–280 nm) or total UV
901 range (400–280 nm). By choosing this approach (using unfiltered, non-weighted UV radiation) the authors
902 neglected a possible wavelength dependence of the biologically effective dose (Vigano et al. 2008). Bruhn
903 et al. (2009), in addition to PAR lamps, used four different lighting sources to obtain desired UV-B and UV-A
904 irradiance; the UV sources were placed at varying distances to yield the reported irradiances. The irradiance
905 spectra of the experimental tubes and the transmission spectra of the glass vials used were further
906 established in order to reveal wavelength dependent responses in the experiments.

907

908

909 **Appendix B**

910 **Global drivers for CO and N₂O contribution**

911 Global CO and N₂O up-scaling was based on the parameterized response equations to UV-irradiation and
912 temperature (see Bruhn et al. 2013, 2014b; Appendix C).

913 The drivers were driven by geospatial satellite information on UV-B temperature and land surface
914 classifications derived from normalized difference vegetation index (NDVI) and snow cover (SC). The
915 UV310nm data was obtained from the Giovanni OMI/Aura Online Visualization and Analysis Daily Level 3
916 Global Gridded Products (<http://giovanni.sci.gsfc.nasa.gov/giovanni/>), the temperature data from MODIS
917 (Moderate Resolution Imaging Spectroradiometer, NASA Earth Observations, <http://neo.sci.gsfc.nasa.gov/>)
918 and NDVI, LAI, and SC data from NASA Earth Observations (<http://neo.sci.gsfc.nasa.gov/>) and handled in a
919 global longitude/latitude grid (250×150). The effect of UV was scaled with the global UV Irradiance at 310
920 nm at surface level (averaged across the years 2005, 2007, 2009, Local Noon Time). Temperature
921 dependence was scaled based on daytime land surface temperatures averaged per month (over the ten
922 years 2001–2010) of available data. The temperature response parameterization did not include
923 temperatures below 0°C, while the CO/N₂O emission from grids with a temperature below 0°C were set to
924 zero in the up-scaling. Land surface area was determined from satellite land dataset information and the
925 area was calculated from longitude/latitude information. Snow cover (SC) was averaged per month (2009)
926 and we excluded areas covered by snow. The remaining land surface area was classified as being vegetation
927 covered or vegetation free based on the NDVI. The up-scaling approach was conservative in the way that
928 the surface area did not include topography. Further, most of the UV-radiation received by leaf surfaces are
929 indeed screened (absorbed or reflected) by the surface wax (Cen and Bornman, 1993; Liakoura et al. 2003;
930 Jacobs et al. 2007). Therefore, for the global estimate of the UV-effect on CO/N₂O emission by vegetation,
931 we assumed an effective Leaf Area Index (LAI) of unity. The proportions of vegetation and sand area were
932 determined by NDVI classification. Response functions for plants were applied to surface covered by
933 vegetation. The NDVI were averaged per month (2009). Correlations between LAI satellite measurement
934 and NDVI from 2009 showed that for instance NDVI around 0.3, 0.5, 0.7 and 0.75 reflects vegetation with
935 0.4, 0.9, 2.0 and 4.0 layers of leaves per ground area (LAI) respectively (data not shown). This information
936 was used to construct four groups with different distributions among vegetation and vegetation free
937 surfaces. The groups are: (1) NDVI<0.2, 100% sand; (2) 0.2<NDVI<0.4, 60% sand and 40% vegetation; 3)
938 0.4<NDVI<0.6, 20% sand and 80% vegetation; and (4) NDVI>0.6, 100% vegetation. These vegetation cover
939 values are lower than a derivation from the NDVI LAI relationship would indicate, but since vegetation
940 cover is clustered by nature with LAI values up to over 5, this must result in more vegetation free areas
941 than an average estimate would produce. For up-scaling, any LAI above 1 would give the same values. This
942 division into group categories decides the percentage area of vegetation and vegetation-free combination
943 in each of the 250×150 grid cells.

944

945 **Appendix C**

946 **Response functions for global CO and N₂O contribution**

947 For each of 250×150 grid cells (Appendix B) we estimated the CO or N₂O emission rate (*ER*) on a monthly
948 basis as

$$ER = \alpha \times e^{\beta \overline{T_{day}}} \times \frac{\overline{UV}}{50} \overline{days \times DL} \times area \times \left(1 - \frac{area_{sc}}{area}\right) \times prop_{cat}$$

949 where α is a base *ER* of ecosystem CO or N₂O emission (nmol m⁻² h⁻¹) measured at 21.4 °C and 50 mW UV-B
950 (see Bruhn et al. 2013, 2014b). We assumed a response to temperature, $(T) = \alpha \times e^{\beta T}$, as measured at
951 leaf level (Bruhn et al. 2013, 2014b) when exposed to UV-B. In the up-scaling we substituted T with a mean
952 daytime temperature for the respective grid cell, $\overline{T_{day}}$. As we have demonstrated near-linear relationships
953 between irradiance of both UV-B and UV-A and CO and N₂O emissions, respectively (Bruhn et al. 2013,
954 2014b), we scaled the base *ER* of ecosystem CO or N₂O emission with the mean UV Irradiance (mW) at 310
955 nm at surface level for the grid cell. Further, *ER* for grid cell was adjusted according to the monthly average
956 day length per month, $\overline{DL}$, number of days per month, area and category of land vegetation.

957

958