



Effect factors for terrestrial acidification in Brazil

Crespo Mendes, Natalia; Laurent, Alexis; Hauschild, Michael Zwicky

Publication date:
2016

[Link back to DTU Orbit](#)

Citation (APA):

Crespo Mendes, N., Laurent, A., & Hauschild, M. Z. (2016). Effect factors for terrestrial acidification in Brazil. Abstract from SETAC Europe 26th Annual Meeting, Nantes, France.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Effect factors for terrestrial acidification in Brazil

Natalia Crespo Mendes¹, Alexis Laurent¹ and Michael Zwicky Hauschild¹

¹Division for Quantitative Sustainability Assessment, Department of Management Engineering, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark
E-mail contact: ncme@dtu.dk

Short Abstract

To support the increased use of existing Life Cycle Impact Assessment (LCIA) methodologies across the world, new methodological elements have been developed towards spatially resolved impact assessment. Spatially resolved methods could better capture the differences of regional environmental conditions, which is an essential approach considering countries like Brazil, with high biodiversity. Previous studies have assessed the impacts of terrestrial acidification from the estimations of the potential losses of vascular plants species richness as a result of exposure to acidifying substances for 13 biomes, with 2409 species addressed for whole world. In this context this work aims to provide spatially-differentiated effect factors (EF) for terrestrial acidification in Brazil and support the development of spatially-differentiated characterization factors for Brazil. In order to maintain compatibility with existing LCIA methods the effect factors were developed using the framework adopted by LC-Impact and Impact World+ methods. Soil pH was used as an indicator of soil acidity to predict plant occurrences. From the number of plant species occurring at each 0.1 pH unit response relationships of species richness and soil pH were developed. The species richness in each ecoregion were transformed into an empirical potentially not occurring fraction, which is a zero-to-one measure used to represent the presence or absence of species. The set of data consists of 976345 records of plants occurrences in Brazil, represented by 33167 species, indicating that this is a comprehensive study. Maps of soil pH in Brazil were extracted at 1-km resolution and pH values were extracted for the depth range of 0-30cm. For each ecoregion, species richness was plotted against soil pH and the exposure-response curves for acidification described the behavior of plant species in a certain region when it is exposed to acidic conditions. From these curves it was possible to derive the effect factors for terrestrial acidification. The results of this work show that spatial differentiation is meaningful when it is possible to combine fine spatial resolutions and highly representative data and this approach can be applied for other impact categories and regions, and contribute to the development of spatial differentiated LCIA methodologies.

Keywords: Effect factors; Acidification; Life Cycle Impact Assessment methodologies, spatial differentiation