



Enzymatic polymerization of bio-based monomers for applications in hydrogels and coatings

Hoffmann, Christian; Nguyen, Hiep Dinh; Storgaard, Thomas; Khan, Anzar; Daugaard, Anders Egede

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Hoffmann, C., Nguyen, H. D., Storgaard, T., Khan, A., & Daugaard, A. E. (2015). Enzymatic polymerization of bio-based monomers for applications in hydrogels and coatings. Abstract from Nordic Polymer Days 2015, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Enzymatic polymerization of bio-based monomers for applications in hydrogels and coatings

Christian Hoffmann^a, Hiep Nguyen^a, Thomas Storgaard^a, Anzar Khan^b, Anders E. Daugaard

^aDanish Polymer Centre, Department of Chemical and Biochemical Engineering, Technical University of Denmark, Søtofts plads bygning 227, DK2800 Kgs. Lyngby, Denmark (adt@kt.dtu.dk)

^bDivision of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University (NTU), Singapore

Enzymatic polymerization has been gradually building up during the last 30 years as an alternative to classical polyesterification processes, which permits preparation of polyesters under more benign conditions with high selectivity¹. In particular, the high selectivity is an interesting property of the enzymatic catalysts that can provide control over polymer structure in functional polymers.

Lipase catalyzed polymerizations (specifically CALB) has been applied to prepare functional polyesters and to evaluate the possibilities of using less stable bio-based monomers such as itaconic acid or its derivatives. Through variation of starting materials, several different functional polyesters have been prepared with the purpose of exploiting the selectivity of the enzymes to prepare polyesters with a controlled degree of branching. Branched polyesters find application in coatings and some of the prepared systems will be discussed in relation to alkyd coatings.

By combination with longer chain water soluble polymers such as PEGs it is also possible to prepare water soluble polyesters with some control over the endgroup structure. In combination with different carboxylic acids, functional polyesters based on PEG have been prepared and functionalized through aza-michael additions as well as through thiol-ene chemistry². Thereby the enzymatically prepared polymer backbone can be considered a scaffold for functional water soluble materials. Finally, these polymers have been applied for preparation of hydrogels, aiming at the potential use of these materials in drug delivery.




Figure 1: Investigation of the possibilities of enzymatic polymerization of functional monomers in different combinations with respect to both functionality of polymers and selectivity during the polymerization.

References

- (1) Gross, R. A.; Kumar, A.; Kalra, B. *Chem. Rev.* **2001**, *101*, 2097.
- (2) Hoffmann, C.; Stuparu, M. C.; Daugaard, A.; Khan, A. *J. Polym. Sci. Part A Polym. Chem.* **2015**, *53*, 745.