



Draft Genome Sequence of *Acinetobacter johnsonii* C6, an Environmental Isolate Engaging in Interspecific Metabolic Interactions

Kaas, Rolf Sommer; Mordhorst, Hanne; Leekitcharoenphon, Pimlapas; Jensen, Jacob Dyring; Haagensen, Janus Anders Juul; Molin, Søren; Pamp, Sünje Johanna

Published in:
Genome Announcements

Link to article, DOI:
[10.1128/genomeA.00155-17](https://doi.org/10.1128/genomeA.00155-17)

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kaas, R. S., Mordhorst, H., Leekitcharoenphon, P., Jensen, J. D., Haagensen, J. A. J., Molin, S., & Pamp, S. J. (2017). Draft Genome Sequence of *Acinetobacter johnsonii* C6, an Environmental Isolate Engaging in Interspecific Metabolic Interactions. *Genome Announcements*, 5(16), [e00155-17]. DOI: 10.1128/genomeA.00155-17

General rights



Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Draft Genome Sequence of *Acinetobacter johnsonii* C6, an Environmental Isolate Engaging in Interspecific Metabolic Interactions

 Rolf Sommer Kaas,^a Hanne Mordhorst,^a Pimlapas Leekitcharoenphon,^a Jacob Dyring Jensen,^a Janus A. J. Haagenen,^b Søren Molin,^b  Sünje Johanna Pamp^a

Research Group for Genomic Epidemiology, Technical University of Denmark, Kongens Lyngby, Denmark^a;
Novo Nordisk Center for Biosustainability, Technical University of Denmark, Kongens Lyngby, Denmark^b

ABSTRACT *Acinetobacter johnsonii* C6 originates from creosote-polluted groundwater and performs ecological and evolutionary interactions with *Pseudomonas putida* in biofilms. The draft genome of *A. johnsonii* C6 is 3.7 Mbp and was shaped by mobile genetic elements. It reveals genes facilitating the biodegradation of aromatic hydrocarbons and resistance to antimicrobials and metals.

A *Acinetobacter johnsonii* C6 (formerly, *Acinetobacter* sp. strain C6) was isolated in 1994 from a microbial community of a creosote-contaminated aquifer at a gasworks in Fredensborg, Denmark (1, 2). Creosotes are mixtures of chemicals formed during natural gas production, which can contain aromatic hydrocarbons and a variety of heterocycles. Despite their toxicity, creosotes were used as medical treatment against infections, toothache, gastrointestinal, and respiratory complications.

A. johnsonii C6 forms biofilms and participates in interspecific interactions, including metabolic interactions, with *Pseudomonas putida* (3–6). The genetic determinants for these activities are largely unknown. Here, we report the draft genome sequence of *A. johnsonii* C6. It was generated using Illumina MiSeq sequencing (2 × 250 cycles), yielding 593,389 raw read pairs and a depth of coverage of ~68×. The reads were trimmed and filtered using bbduk2 (BBMap 35.82) (<http://jgi.doe.gov/data-and-tools/bbtools/>) and assembled using SPAdes 3.7.0 (7). Contigs smaller than 500 bp or with coverage below 2× were removed. The draft genome is 3,705,435 bp in 26 contigs, with a G+C content of 41.7%. It contains 3,543 genes, as predicted using Prodigal (8), 77 tRNA genes, and one rRNA operon (16S, 23S, 5S). The 16S rRNA gene sequence had >99% sequence similarity to *A. johnsonii* XBB1 (accession no. NZ_CP010350.1), *A. johnsonii* ATCC 17909^T (accession no. Z93440.1), and *A. johnsonii* DSM 6963 (accession no. X81663.1) (9–11). Putative functions for predicted proteins were assigned using PROKKA 1.1 and by comparing sequences to the public databases Pfam, KEGG, InterPro, and CARD (12–16), followed by submission-ready file conversion (https://bitbucket.org/RolfKaas/gff3_to_ena_embl).

A. johnsonii C6 encodes proteins predicted to convert aromatic hydrocarbons, such as benzyl alcohol, benzoate, fluorobenzoate, dihydroxybenzoate, methylcatechol, methylbenzyl alcohol, hydroxybenzaldehyde, hydroxymethylnaphthalene, naphthalenemethanol, benzene, toluene, chlorobenzene, and cyclohexanol. Previously, it was shown that this strain could grow on toluene, benzyl alcohol, and benzoate (4, 5).

A number of antimicrobials, as well as heavy metals (e.g., arsenate, mercury, tellurite, copper, and chromate), may be tolerated by *A. johnsonii* C6, mainly facilitated by proteins involved in their efflux, transport, reduction, and functions encoded by anti-

Received 25 February 2017 Accepted 2 March 2017 Published 20 April 2017

Citation Kaas RS, Mordhorst H, Leekitcharoenphon P, Dyring Jensen J, Haagenen JAJ, Molin S, Pamp SJ. 2017. Draft genome sequence of *Acinetobacter johnsonii* C6, an environmental isolate engaging in interspecific metabolic interactions. *Genome Announc* 5:e00155-17. <https://doi.org/10.1128/genomeA.00155-17>.

Copyright © 2017 Kaas et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sünje Johanna Pamp, sjpa@food.dtu.dk.

biotic resistance genes, such as *bla*_{OXA-334} (OXA-211 family) and *catB*. *In vitro* assays revealed that *A. johnsonii* C6 was resistant to chloramphenicol, trimethoprim, cefoxitin, and quinupristin-dalfopristin. *A. johnsonii* C6 may produce secondary metabolites, and it harbors biosynthetic gene clusters for a siderophore, aryl polyene, bacteriocin, and unknown metabolites, based on predictions by antiSMASH (17).

The *A. johnsonii* C6 draft genome encodes 19 proteins containing GGDEF and/or EAL domains involved in c-di-GMP metabolism, and proteins involved in motility (pili), and secretion (type II secretion system [T2SS], T6SS, secretory-signal recognition particle [Sec-SRP], and Tat), suggesting dynamic interactions with their environment, including with other microorganisms. The presence of features related to plasmids, phages, and insertion sequence (IS) elements suggests that mobile genetic elements have shaped the evolution and ecology of *A. johnsonii* C6.

The genome sequence of *A. johnsonii* C6 will facilitate the understanding of its physiology, evolution, and interaction with *P. putida*. Studies on *A. johnsonii* could also provide new insight into the biodegradation of aromatic hydrocarbons and resistance to antimicrobials and toxic metals, with relevance to environmental biotechnology.

Accession number(s). The draft genome sequence of *A. johnsonii* C6 is available from DDBJ/ENA/GenBank under the accession number [FUUY00000000](https://www.ncbi.nlm.nih.gov/nuccore/FUUY00000000).

ACKNOWLEDGMENTS

The support from Frank M. Aarestrup and the National Food Institute at the Technical University of Denmark are greatly appreciated.

REFERENCES

- Flyvbjerg J, Arvin E, Jensen BK, Olsen SK. 1993. Microbial degradation of phenols and aromatic hydrocarbons in creosote-contaminated groundwater under nitrate-reducing conditions. *J Contam Hydrol* 12:133–150. [https://doi.org/10.1016/0169-7722\(93\)90018-N](https://doi.org/10.1016/0169-7722(93)90018-N).
- Møller S, Pedersen AR, Poulsen LK, Arvin E, Molin S. 1996. Activity and three-dimensional distribution of toluene-degrading *Pseudomonas putida* in a multispecies biofilm assessed by quantitative *in situ* hybridization and scanning confocal laser microscopy. *Appl Environ Microbiol* 62:4632–4640.
- Hansen SK, Rainey PB, Haagensen JAJ, Molin S. 2007. Evolution of species interactions in a biofilm community. *Nature* 445:533–536. <https://doi.org/10.1038/nature05514>.
- Haagensen JAJ, Hansen SK, Christensen BB, Pamp SJ, Molin S. 2015. Development of spatial distribution patterns by biofilm cells. *Appl Environ Microbiol* 81:6120–6128. <https://doi.org/10.1128/AEM.01614-15>.
- Møller S, Sternberg C, Andersen JB, Christensen BB, Ramos JL, Givskov M, Molin S. 1998. *In situ* gene expression in mixed-culture biofilms: evidence of metabolic interactions between community members. *Appl Environ Microbiol* 64:721–732.
- Christensen BB, Haagensen JAJ, Heydorn A, Molin S. 2002. Metabolic commensalism and competition in a two-species microbial consortium. *Appl Environ Microbiol* 68:2495–2502. <https://doi.org/10.1128/AEM.68.5.2495-2502.2002>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
- Feng Y, Yang P, Wang X, Zong Z. 2016. Characterization of *Acinetobacter johnsonii* isolate XBB1 carrying nine plasmids and encoding NDM-1, OXA-58 and Per-1 by genome sequencing. *J Antimicrob Chemother* 71:71–75. <https://doi.org/10.1093/jac/dkv324>.
- Rainey FA, Lang E, Stackebrandt E. 1994. The phylogenetic structure of the genus *Acinetobacter*. *FEMS Microbiol Lett* 124:349–353. <https://doi.org/10.1111/j.1574-6968.1994.tb07307.x>.
- Ibrahim A, Gerner-Smidt P, Liesack W. 1997. Phylogenetic relationship of the twenty-one DNA groups of the genus *Acinetobacter* as revealed by 16S ribosomal DNA sequence analysis. *Int J Syst Bacteriol* 47:837–841. <https://doi.org/10.1099/00207713-47-3-837>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Finn RD, Coghill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res* 44:D279–D285. <https://doi.org/10.1093/nar/gkv1344>.
- Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. 2017. KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res* 45:D353–D361. <https://doi.org/10.1093/nar/gkw1092>.
- Finn RD, Attwood TK, Babbitt PC, Bateman A, Bork P, Bridge AJ, Chang HY, Dosztányi Z, El-Gebali S, Fraser M, Gough J, Haft D, Holliday GL, Huang H, Huang X, Letunic I, Lopez R, Lu S, Marchler-Bauer A, Mi H, Mistry J, Natale DA, Necci M, Nuka G, Orengo CA, Park Y, Pesseat S, Piovesan D, Potter SC, Rawlings ND, Redaschi N, Richardson L, Rivoire C, Sangrador-Vegas A, Sigrist C, Sillitoe I, Smithers B, Squizzato S, Sutton G, Thanki N, Thomas PD, Tosatto SCE, Wu CH, Xenarios I, Yeh L-S, Young S-Y, Mitchell AL. 2017. InterPro in 2017—beyond protein family and domain annotations. *Nucleic Acids Res* 45:D190–D199. <https://doi.org/10.1093/nar/gkw1107>.
- Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, Lago BA, Dave BM, Pereira S, Sharma AN, Doshi S, Courtot M, Lo R, Williams LE, Frye JG, Elsayegh T, Sardar D, Westman EL, Pawlowski AC, Johnson TA, Brinkman FSL, Wright GD, McArthur AG. 2017. CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res* 45:D566–D573. <https://doi.org/10.1093/nar/gkw1004>.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Bruccoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43:W237–W243. <https://doi.org/10.1093/nar/gkv437>.