



Synthesis of arabinoxylan fragments

Underlin, Emilie Nørmølle; Böhm, Maximilian F.; Madsen, Robert

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Underlin, E. N., Böhm, M. F., & Madsen, R. (2017). Synthesis of arabinoxylan fragments. Abstract from 19th European Carbohydrate Symposium, Barcelona, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



SYNTHESIS OF ARABINOXYLAN FRAGMENTS

Emilie N. Underlin¹, Maximilian F. Böhm², Robert Madsen¹

¹ Technical University of Denmark, Department of Chemistry, Kgs. Lyngby, Denmark.

² Carbosynth LLC, Oxford, England.

e-mail: emnu@kemi.dtu.dk

The cell wall of plants can be termed the skeleton of the plant. One of the parts making up the cell wall is hemicellulose. Hemicellulose is composed of a number of saccharides where one of the most abundant are the arabinoxylan-oligosaccharides. In many instances the biosynthesis and degradation of arabinoxylan oligosaccharides remain elusive. As a consequence defined arabinoxylan fragments have been chosen as synthetic targets which subsequently will be submitted to enzymatic studies. A better understanding of these processes could lead to e.g. better utilisation of the biomass for biofuel production, or production of commercial chemicals which are mainly obtained from fossil fuels today [1].

The arabinoxylan fragments have a backbone of β -1,4-linked xylans with α -L-arabinose units attached at specific positions. The synthesis utilises an efficient synthetic route, where all the xylan units can be derived from D-xylose through a common intermediate. The xylan units feature the same thiophenyl donor functionality to allow for successive coupling with the same optimised glycosylation protocol [2]. Then, the arabinose units can be attached and global deprotection yields the target fragments.




Figure 1: Retrosynthetic overview with one target as an example.

[1] H. Scheller; P. Ulvskov; Hemicelluloses. *Annu. Rev. Plant Biol.*, **2010**, *61*, 263-289

[2] D. Crich; F. Cai; F. Yang; A stable, commercially available sulfenyl chloride for the activation of thioglycosides in conjunction with silver trifluoromethanesulfonate. *Carbohydr. Res.*, **2008**, *343*, 1858-1862.