



Diagnostic herd sensitivity using environmental samples

Vigre, Håkan; Josefsen, Mathilde Hartmann; Seyfarth, Anne Mette; Agersø, Yvonne

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Vigre, H., Josefsen, M. H., Seyfarth, A. M., & Agersø, Y. (2012). Diagnostic herd sensitivity using environmental samples. Abstract from 13th Conference of the International Society for Veterinary Epidemiology and Economics, Maastricht, Netherlands.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract – oral presentation

Diagnostic herd sensitivity using environmental samples

Vigre, H., Josefsen, M., Seyfarth, A.M. and Agersø, Y., DTU Food, Denmark; hvig@food.dtu.dk

Due to logistic and economic benefits, the animal industry has an increased interest in using environmental samples to classify herds free of infections. For a valid interpretation of results obtained from environmental samples, the performance of the diagnostic method using these samples must be assessed. In this abstract we present an approach to evaluate the sensitivity of environmental samples used for herd classification. This is illustrated using results from samples for isolation of methicillin resistant *Staphylococcus aureus* (MRSA) in 48 pig herds that had previously been tested positive either at farm or slaughter. Three sample matrices were collected; dust samples (5 environmental swabs), nasal swabs (10 pools with 5 animals per pool) and air samples (1 filter). Based on the assumption that MRSA occurred in all 48 herds the overall herd sensitivity was 58% for nasal swabs, 33% for dust samples and 63% for air samples. However, we assumed that the sensitivity of environmental samples varies between herds due to variation in the amount of the analyte. This assumption can be tested by analyzing the association between animal prevalence and the result obtained using environmental samples. In our example, the prevalence of infected pigs in each herd was estimated from the pooled samples of nasal swabs. Logistic regression was used to estimate the effect of animal prevalence on the probability to detect MRSA in the dust and air samples at herd level. The results show a significant increase in the probability of isolating MRSA in the environmental samples by increasing within-herd prevalence. This relationship was strongest for dust where the sensitivity increased from 25% at 1% infected pigs, to 35% at a 25% infected pigs. The air sampling was more sensitive to detect infected herds irrespectively of the within herd prevalence, and performed almost perfectly at a prevalence of 25% infected pigs (sensitivity=99%). In general, the dependence of within herd prevalence should be considered in designing surveillance programs based on environmental samples.