



Succinic acid production from Jerusalem artichoke

Gunnarsson, Ingólfur Bragi; Karakashev, Dimitar Borisov; Angelidaki, Irini

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Gunnarsson, I. B., Karakashev, D. B., & Angelidaki, I. (2013). Succinic acid production from Jerusalem artichoke. Poster session presented at EU BC&E 21st European Biomass Conference and Exhibition, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Succinic acid production from Jerusalem artichoke tubers

Ingólfur B. Gunnarsson, Dimitar B. Karakashev and Irimi Angelidaki

Succinic acid

- Succinic acid is currently produced from petro chemicals.
- Has applications in agricultural, food, chemical and pharmaceutical industries.
- *A. succinogenes* 130Z produces succinic acid in large amounts during fermentation.
- Theoretically consumes 1 mol CO₂ per 1 mol succinic acid produced.
- Could contribute to the abatement of CO₂ emissions at a time when CO₂ levels in earth's atmosphere are at an all time high (>400 ppm).

Jerusalem artichoke tubers

- Root vegetables rich in inulin.
- Inulin is a D-fructose and D-glucose polysaccharide.
- Poor digestibility in humans.
- Investigate using Jerusalem artichoke tubers for producing succinic acid.



Results

Fermentation of pure sugars (batch bottles):

Succinic acid (SA) yield was 83% when D-glucose was the substrate compared to 72% yield SA when D-fructose was the substrate.


D-glucose utilization was 93% compared to 88.5% D-fructose utilization.

		Glucose	Fructose
0 hours	Initial reducing sugar (g/L)	48.0±0.5	46.2±0.3
48 hours	Residual reducing sugar (g/L)	3.4±0.2	5.4±0.1
	Succinic acid (g/L)	37.2±0.6	29.4±0.1
	Formic acid (g/L)	5.8±0.2	7.0±0.2
	Acetic acid (g/L)	8.2±0.1	7.2±0.5
	Sugar utilization (%)	93.0±0.5	88.5±0.2
	Succinic acid yield* (%)	83.0±1.3	72.0±0.5

Results

Fermentation of tuber hydrolysates:

SA yield was 80.2% in bottles compared to 77.9% in 3-L bioreactor. However the SA production was increased by 26.4% in 3-L bioreactor and the D-fructose utilization increased from 68.5% to 96.5%.



Conclusions

- This is the first report on the use of Jerusalem artichoke tuber hydrolysate as substrate for succinic acid production by *A. succinogenes*.
- Succinic acid was successfully produced, where the highest succinic acid concentration was 26.8 g/L and the highest succinic acid yield observed was 80.2%.
- *A. succinogenes* 130Z utilized both D-fructose and D-glucose in the tuber hydrolysates.
- The importance of pH control and its effect on D-fructose utilization during succinic acid production was clearly observed when results from batch bottles and 3-L bioreactor were compared.
- Jerusalem artichoke tubers as raw material for succinic acid production is a very attractive option compared to other types of biomass tested for succinic acid production.