



Aspergillus triggers phenazine production in Pseudomonas aeruginosa

Jensen, Britt Guillaume; Jelsbak, Lars; Søndergaard, Ib; Pedersen, Mona Højgaard; Frisvad, Jens Christian; Nielsen, Kristian Fog

Publication date:
2011

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):
Jensen, B. G., Jelsbak, L., Søndergaard, I., Pedersen, M. H., Frisvad, J. C., & Nielsen, K. F. (2011). Aspergillus triggers phenazine production in Pseudomonas aeruginosa. Abstract from European Congress on Microbial Biofilms – Basic and Clinical Aspects, Copenhagen, Denmark, .

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract for Eurobiofilms 2011, Copenhagen, Denmark, 6-8 July 2011

Aspergillus* triggers phenazine production in *Pseudomonas aeruginosa

B.G. Jensen, L. Jelsbak, I. Søndergaard, M.H. Pedersen, J.C. Frisvad, K.F. Nielsen
Department of Systems Biology, Technical University of Denmark, DK-2800 Kgs. Lyngby,
Denmark.

Objectives:

Pseudomonas aeruginosa is an opportunistic human pathogen, commonly infecting cystic fibrosis (CF) patients. Aspergilli, especially *Aspergillus fumigatus*, are also frequently isolated from CF patients. Our aim was to examine the possible interaction between *P. aeruginosa* and different *Aspergillus* species.

Methods:

A suspension of fungal spores was streaked onto WATM agar plates. After 24 hours incubation at 37 °C, a *P. aeruginosa* overnight culture was streaked out perpendicular to the fungal streak. The plates were incubated at 37 °C for five days, examined and plugs were extracted for HPLC-DAD and HPLC-DAD-MS analysis.

Results:

P. aeruginosa PAO1 suppressed growth of *A. fumigatus*, *A. niger*, *A. flavus*, *A. oryzae*, *A. terreus* and *Emericella nidulans*. HPLC and HPLC-DAD-MS results showed an increase in phenazine-1-carboxylic acid and phenazine-1-carboxamide production by *P. aeruginosa* in the contact area of *A. niger*, *A. flavus*, *A. oryzae*, but not *A. fumigatus*. In addition, other metabolites with UV chromophores similar to the phenazines were only found in the contact zone between *Aspergillus* and *Pseudomonas*. No change in secondary metabolite profiles were seen for the Aspergilli, when comparing with or without the presence of *Pseudomonas*.

Conclusion:

All Aspergilli tested, with the exception of *A. fumigatus*, triggered the upregulation of phenazine-1-carboxamide and phenazine-1-carboxylic acid production by *P. aeruginosa*. Surprisingly no changes in secondary metabolite profiles were detected in any of the Aspergilli.