



Simulation of the Initial 3-D Instability of an Impacting Drop Vortex Ring

Sigurdson, Lorenz; Wiwchar, Justin; Walther, Jens Honore

Published in:
American Physical Society. Bulletin

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Sigurdson, L., Wiwchar, J., & Walther, J. H. (2013). Simulation of the Initial 3-D Instability of an Impacting Drop Vortex Ring. American Physical Society. Bulletin, 58(8), [Abstract: R6.00005].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract Submitted
for the DFD13 Meeting of
The American Physical Society

Simulation of the Initial 3-D Instability of an Impacting Drop Vortex Ring LORENZ SIGURDSON, JUSTIN WIWCHAR, Vortex Fluid Dynamics Lab, Mechanical Engineering Department, University of Alberta, JENS WALTHER, Mechanical Engineering, Danish Technical University, VF DL TEAM — Computational vortex particle method simulations of a perturbed vortex ring are performed to recreate and understand the instability seen in impacting water drop experiments. Three fundamentally different initial vorticity distributions are used to attempt to trigger a Widnall instability, a Rayleigh centrifugal instability, or a vortex breakdown-type instability. Simulations which simply have a perturbed solitary ring result in an instability similar to that seen experimentally. Waviness of the core which would be expected from a Widnall instability is not visible. Adding an opposite-signed secondary vortex ring or an image vortex ring to the initial conditions, to trigger a Rayleigh or breakdown respectively, does not appear to significantly change the instability from what is seen with a solitary ring. This suggests that a Rayleigh or vortex breakdown-type instability are not likely at work, though tests are not conclusive. Perhaps the opposite-signed secondary vortex was not strong enough or placed appropriately. Elliptical streamlines, as expected, are visible in the core of the solitary ring at early times. Support from the Canadian Natural Sciences and Engineering Research Council grant 41747 is gratefully acknowledged.

Lorenz Sigurdson
Vortex Fluid Dynamics Lab, Mechanical Engineering Department,
University of Alberta

Date submitted: 02 Aug 2013

Electronic form version 1.4