



N2O production dynamics in nitrifying/denitrifying activated sludge under defined environmental conditions

Pellicer i Nàcher, Carles; Jensen, Marlene Mark; Petersen, Morten S.; Smets, Barth F.

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Pellicer i Nàcher, C., Jensen, M. M., Petersen, M. S., & Smets, B. F. (2013). N2O production dynamics in nitrifying/denitrifying activated sludge under defined environmental conditions. Abstract from ICON3: 3rd international conference on Nitrification, Tokoy, Japan.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ICON 3 – 3rd International conference on nitrification, 2nd-5th September 2013,
Tokyo (Japan)

N₂O production dynamics in nitrifying/denitrifying activated sludge under defined environmental conditions

Carles Pellicer-Nàcher, Marlene Mark Jensen, Morten S. Petersen, and Barth F.
Smets

Nitrous oxide (N₂O) is a gaseous pollutant emitted as an unwanted product in wastewater treatment plants during the nitrification-denitrification process. Even though the emission capacity of the process with respect to this compound is still under debate, N₂O has been identified as an important contributor to global warming and the destruction of the ozone layer. The present study makes use of unique datasets collected during controlled batch tests with activated sludge biomass to test and calibrate a pseudo-mechanistic model that predicts N₂O production by nitrifying and heterotrophic bacteria. The proposed model described successfully the observed N₂O production dynamics and confirmed that the availability of ammonia, low dissolved oxygen and nitrite accumulation were the main factors triggering N₂O production. Nitrifier-denitrification was proposed as the main pathway catalyzing the conversion of fixed nitrogen to N₂O. Heterotrophic denitrification rates were one order of magnitude lower than nitrification rates and contributed marginally to the overall N₂O production. Further data analysis allowed derivation of the overall mass transfer coefficients describing gaseous stripping and revealed that a minor portion of the N₂O produced was actually released to the gas phase. This work represents a step further in the use and calibration of process models to control and understand better N₂O production and emissions during conventional wastewater treatment.