



## PAH in tea and coffee

Duedahl-Olesen, Lene; Navarantem, Marin; Adamska, Joanna; Højgård, Arne

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Duedahl-Olesen, L., Navarantem, M., Adamska, J., & Højgård, A. (2013). PAH in tea and coffee. Abstract from 24th International Symposium on Polycyclic Aromatic Compounds (ISPAC 2013), Oregon, United States.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## PAH IN TEA AND COFFEE

\*Lene Duedahl-Olesen<sup>1</sup>, Marin Navarantem<sup>1</sup>, Joanna Adamska<sup>1</sup>, Arne Højgård<sup>2</sup>

1. National Food Institute, Technical University of Denmark, Søborg, Denmark

2. Regional Veterinary and Food Administration Center, Ministry of Food, Agriculture and Fisheries, Århus, Denmark

For food regulation in the European Union maximum limits on other foods than tea and coffee includes benzo[*a*]pyrene and the sum of PAH4 (sum of benzo[*a*]pyrene, chrysene, benzo[*a*]anthracene and benzo[*b*]fluoranthene). This study includes analysis of the above mentioned PAH in both, tea leaves, coffee beans and ready-to-drink preparations. Compared to other food matrices (e.g. fish), the analytical methods were challenged by the hot water extracts.

Preparation of tea includes roasting and drying of the tea leaves using combustion gases from burning wood, oil, or coal. These are responsible for accumulation of PAH in tea leaves. Different varieties of tea leaves were analyzed and highest concentrations were found in leaves from mate and black tea with maximum concentrations of 32 µg/kg for benzo[*a*]pyrene and 115 µg/kg for the sum of PAH4.

Also, coffee beans are roasted during processing. However, both benzo[*a*]pyrene and PAH4 concentrations were more than ten times lower for coffee beans than for tea leaves. Highest levels were found for PAH4 of solid instant coffee (5.1 µg/kg).

Data were used to calculate the exposure of benzo[*a*]pyrene (15%) and sum of PAH4 (10%) from tea and coffee for the Danish population.