

## **Bioaugmentation of an acetate-oxidising anaerobic consortium in up-flow sludge blanket reactor subjected to high ammonia loads**

**Fotidis, Ioannis; Karakashev, Dimitar Borisov; Angelidaki, Irini**

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Fotidis, I., Karakashev, D. B., & Angelidaki, I. (2013). Bioaugmentation of an acetate-oxidising anaerobic consortium in up-flow sludge blanket reactor subjected to high ammonia loads. Abstract from 13th World Congress on Anaerobic Digestion, Santiago De Compostela, Spain.

## **DTU Library** Technical Information Center of Denmark

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Bioaugmentation of an acetate-oxidising anaerobic consortium in up-flow sludge blanket reactor subjected to high ammonia loads

I. A. Fotidis, D. B. Karakashev, I. Angelidaki

Department of Environmental Engineering, Technical University of Denmark, Building 113,  
DK-2800 Kgs. Lyngby, Denmark

(E-mail: [ioanf@env.dtu.dk](mailto:ioanf@env.dtu.dk); [dbka@env.dtu.dk](mailto:dbka@env.dtu.dk); [iria@env.dtu.dk](mailto:iria@env.dtu.dk))

## Abstract

Ammonia is the major inhibitor of anaerobic digestion (AD) process leading to suboptimal utilisation of the biogas potential of the feedstocks and causing economical losses to the biogas plants. However, ammonia is mainly inhibiting the acetoclastic methanogens, while the hydrogenotrophic methanogens, in syntrophic association with acetate oxidising bacteria, are more resistant to ammonia toxicity effect. The use of syntrophic acetate oxidising methanogenic consortia could provide a new approach to tackle ammonia toxicity effect in AD. The SAO culture (i.e. *Clostridium ultunense* spp. nov. in association with *Methanoculleus* spp. strain MAB1), is an acetate oxidising methanogenic consortium that can produce methane (CH<sub>4</sub>) at high ammonia levels. In the current study the bioaugmentation of the SAO culture in a mesophilic up-flow anaerobic sludge blanket (UASB) reactor subjected to high ammonia loads was tested. The co-cultivation in fed-batch of a fast-growing hydrogenotrophic methanogen (i.e. *Methanoculleus bourgensis*) with the SAO culture was also investigated. Results obtained clearly demonstrated that bioaugmentation of SAO culture in a UASB reactor was not possible most probably due to the slow growth of the culture. The incubation period (duration of lag+exponential phase) of SAO culture was reduced more than 30% when it was co-cultivated with *Methanoculleus bourgensis*, in fed-batch reactors. Therefore, the bioaugmentation of the SAO culture along with *Methanoculleus bourgensis* in a UASB reactor is a promising approach to overcome ammonia toxicity.