



Time-resolved angular distributions of plume ions from silver at low and medium laser fluence

Christensen, Bo Toftmann; Schou, Jørgen

Publication date:
2011

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):

Christensen, B. T., & Schou, J. (2011). Time-resolved angular distributions of plume ions from silver at low and medium laser fluence. Abstract from 11th International Conference on Laser Ablation, Playa del Carmen, Mexico.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Time-resolved angular distributions of plume ions from silver at low and medium laser fluence

B. Toftmann and J. Schou

DTU Fotonik, Risø Campus, Technical University of Denmark, DK-4000 Roskilde, Denmark (josc@fotonik.dtu.dk)

Even at moderate fluence (0.6 -2.4 J/cm²) laser impact on metals in the UV regime results in a significant number of ions emitted from the surface. The ablated particles are largely neutrals at the lowest fluence, but the fraction of ions increases strongly with fluence.

We have irradiated silver in a vacuum chamber ($\sim 10^{-7}$ mbar) with a Nd:YAG laser at a wavelength of 355 nm and made detailed measurements of the time-resolved angular distribution. The ion flow in different directions has been measured with a hemispherical array of Langmuir probes, by which the time-of-flight spectra, as well as the total angular yield, in all directions can be recorded (1). The angular distribution peaks more strongly in forward direction with increasing fluence - as also reported in the literature – and can be well approximated by the model of Anisimov et al. (1,2).

Typically, the spectra of silver ions peak from 70 eV up to 145 eV in a direction normal to the target surface with increasing fluence. With increasing observation angle the time-of-flight spectra exhibit a peak at longer flight times, i.e. at a lower kinetic energy. At the highest fluence the ionized fraction of the ablated particles exceeds 0.5.

(1) Thestrup et al. *Appl. Surf. Sci.* 197-198, 175 (2002).

(2) Anisimov et al. *Phys. Rev. B* 48, 12076 (1993).