



Single-molecule conductivity of non-redox and redox molecules at pure and gold-mined Au(111)-electrode surfaces

Zhang, Jingdong; Chi, Qijin; Ulstrup, Jens; Wang, Yun; Hush, Noel S.; Reimers, Jeff R.

Publication date:
2011

[Link back to DTU Orbit](#)

Citation (APA):

Zhang, J., Chi, Q., Ulstrup, J., Wang, Y., Hush, N. S., & Reimers, J. R. (2011). Single-molecule conductivity of non-redox and redox molecules at pure and gold-mined Au(111)-electrode surfaces. Abstract from 5th Gerischer Symposium, Berlin, .

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Single-molecule conductivity of non-redox and redox molecules at pure and gold-mined Au(111)-electrode surfaces

Jingdong Zhang, Qijin Chi and Jens Ulstrup

Department of Chemistry, Technical University of Denmark, Kongens Lyngby, Denmark

Yun Wang, Noel S. Hush and Jeff R. Reimers

School of Chemistry, The University of Sydney, NSW 2006, Australia

The structure, two-dimensional organization, and function of molecules immobilized on solid surfaces can be addressed in a degree of detail that has reached the level of the single-molecule. In this context redox molecules are “smart” molecules adding sophisticated electronic function. Redox metalloproteins such as blue copper, heme, and iron-sulfur proteins as well as redox metalloenzymes are other new targets for single-molecule electrochemical and bioelectrochemical surface science. Electrochemistry combined with scanning tunnelling and atomic force microscopy in aqueous chemical or biological media supported by comprehensive theoretical frames, have emerged as core approaches in these exciting areas.

Single-molecule redox electrochemistry is rooted in two major areas. One is the preparation of well-defined (atomically planar) electrode surfaces modified by molecular monolayers (SAMs). High-resolution in situ STM combined with large-scale theoretical computations have offered detailed insight in the surface binding modes of functionalized alkanethiols on Au-surfaces in particular. Such surfaces have disclosed a variety of surface structures that involve both direct binding to a planar Au and binding to surface-mined Au-atoms. In addition the SAMs ensure protein/enzyme immobilization gentle enough that the proteins retain electron transfer or enzyme activity in a variety of local environments. The second area is the mapping and control of the immobilized redox molecules and metalloproteins themselves. Single-molecule resolution has also here been achieved, both in structural mapping and in controlled single-molecule electron transport and enzyme function on SAM surfaces such as those now characterized to (sub)-molecular resolution.

Three recent references:

1. J. Zhang, A.M. Kuznetsov, I.G. Medvedev, Q. Chi, T. Albrecht, P.S. Jensen and J. Ulstrup, Chem. Rev. 108 (2008) 2737-2791.
2. J. Zhang, A.C. Welinder, Q. Chi, and J. Ulstrup, PCCP 12 (2011) 9999-10011.
3. E. Della Pia, Q. Chi, D. Jones, J.E. MacDonald, J. Ulstrup and M. Elliot, Nano Letters 11 (2011) 176-182.