Is organic farming a mitigation option?

A study of N₂O emission from winter wheat

Carter, Mette Sustmann; Albert, Kristian Rost; Ambus, Per

Publication date:
2009

[Link back to DTU Orbit](#)

Citation (APA):

Carter, M. S., Albert, K. R., & Ambus, P. (2009). *Is organic farming a mitigation option? A study of N₂O emission from winter wheat*. Poster session presented at International Scientific Congress on Climate Change, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Is organic farming a mitigation option?

– A study on N₂O emission from winter wheat

Mette S. Carter¹, Kristian Albert² and Per Ambus¹

1) Risø National Laboratory for Sustainable Energy, Biosystems Division, Technical University of Denmark; 2) Institute of Biology, University of Copenhagen, Denmark

CROPSYS

A 4-year interdisciplinary project

The overall goal is to quantify productivity and environmental impacts of different organic and conventional cropping systems across a range of soil and climatic conditions, and to identify management measures which contribute significantly to a sustainable development of the individual cropping systems.

Background

Agricultural soils is the major source of the strong greenhouse gas nitrous oxide (N₂O). In general, organic farming is regarded as a production system with low environmental impact, but it may not always be the case. Crop production in organic farming to a large extent relies on nitrogen (N) supply via decomposition of plant residue, manure and soil organic matter, which needs to be

synchronised with the crop N demand in order to avoid N₂O losses. This is in contrast to conventional farming where mineral fertilizer is supplied when needed for plant growth. Statistical analyses have indicated that the N₂O emission factor is higher for manure (and presumably crop residues) than for mineral fertilizers (Lægveit and Aastveit, 2002).

Objective and experimental design

The objective of the study is to evaluate whether N₂O emissions from cropping systems are affected by 1) organic versus conventional farming, 2) proportion of N₂-fixing crops in the rotation and 3) use of catch crops. Nitrous oxide fluxes were measured in winter wheat field plots that belonged to four different long-term crop rotations at Flakkebjerg, Denmark:

C4-CC Conventional rotation without green manure and catch crops

O4-CC Organic rotation without green manure and catch crops

O4+CC Organic rotation without green manure, but with catch crops

O2+CC Organic rotation based on green manure with catch crops

Monitoring took place every second week from sowing in October 2007 to harvest in August 2008 using two-part static chambers (0.6×0.6 m²).

Fig 1. Conventional winter wheat (C4-CC) and organic winter wheat (O4+CC) in May 2008. The crop yields in the conventionally managed plots were twice as high as in the organic plots.

Fig. 2 Daily N₂O fluxes and mean N₂O fluxes during autumn-winter, spring and summer in four crop rotations. Bars with same letter are not significantly different (P>0.05).

Conclusions

- Organic farming could potentially be a mitigation option for atmospheric emissions of N₂O.
- During spring the N₂O emission from the conventional crop rotation (C4-CC) was higher than from the equivalent organic rotation (O4-CC; Fig. 2).
- During summer there was a tendency for a higher N₂O loss from the conventional rotation (P=0.066; Fig. 2).
- The N₂O emissions related to the production of a given amount of winter wheat was similar in the organic and conventional systems.
- The N₂O emission from the cropping systems was not affected by the presence of catch crops or by the proportion of N₂-fixing crops in the rotation (Fig. 2).

Acknowledgements

The project is linked to the International Centre for Research in Organic Food Systems and funded under the research programme: Research in Organic Food and Farming, International Research Co-operation and Organic Integrity (DARCOF III 2005-2010). Co-ordinated by Professor, J.E. Olesen, University of Aarhus.

Reference

Lægveit, M. and Aastveit, A.H., In: Petersen, S.O. and Olesen, J.E. (eds.). DIAS report, Plant Production no. 81, 2002, pp. 122-134.

Poster presented at International Scientific Congress on Climate Change, Copenhagen, 10-12 March 2009

Contact

Scientist Mette S. Carter
mthy@risoe.dtu.dk or
 Prof. Per Ambus
peam@risoe.dtu.dk