

## The effect of vocal tract impedance on the vocal folds

**Agerkvist, Finn T.; Selamtzis, Andreas**

*Published in:*  
9th Pan European Voice Conference

*Publication date:*  
2011

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Agerkvist, F. T., & Selamtzis, A. (2011). The effect of vocal tract impedance on the vocal folds. In 9th Pan European Voice Conference: Proceedings

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## The effect of vocal tract impedance on the vocal folds

Finn Agerkvist, Andreas Selamtsis

Acoustic Technology, Department of Electrical Engineering, Technical University of Denmark, Lyngby, Denmark  
{ [fa@elektro.dtu.dk](mailto:fa@elektro.dtu.dk) ; [a.selamsis@gmail.com](mailto:a.selamsis@gmail.com) }

The importance of the interaction between the acoustic impedance of the vocal tract with the flow across the vocal cords is well established. In this paper we are investigating the changes in vocal tract impedance when using the different modes of phonation according to Sadolin [1], going from the soft levels of the *Neutral* mode to the high levels of the fully ‘metallic’ *Edge* mode. The acoustic impedance of vocal tract as seen from the mouth opening is measured via a microphone placed close to the mouth when exciting the system with a volume velocity source [2]. At the same time a *Laryngograph* frontend is used to measure the electroglottograph signal which reflects the opening and closing pattern of the vocal folds. The measurements were carried out for all four modes (Neutral, Curbing, Overdrive and Edge) for the vowel [a] in three different pitches: C3(131 Hz), G3 (196 Hz) and C4 (262Hz) . The results show changes in the resonance frequencies of the vocal tract with increasing pitch, whereas the changes between the modes are less clear due to the measurement signal being weak in comparison to the louder modes, especially at high pitches. The electroglottograph shows a very different waveform for the *Neutral* mode compared to the other, so-called metallic modes. The differences in waveform between *Curbing*, *Overdrive* and *Edge* modes are minor. However, the spectrum of the *Overdrive* mode shows stronger 2<sup>nd</sup> harmonic and weaker 4<sup>th</sup> and 6<sup>th</sup> harmonic compared to *Curbing* and *Edge*.

Finally the *Overdrive* mode, which is the mode that is most limited in pitch range, was tested at its pitch limit C5 (523 Hz) under normal conditions and when the singer has inhaled Helium. When inhaling Helium the acoustic impedance of the vocal tract is reduced in magnitude and the resonances are scaled upwards in frequency due to different density and speed of sound in Helium. The electroglottograph shows a change in waveform when the singer inhales helium. The percentage of the glottal cycle when the vocal cords are open, the so-called open quotient, increases from 40 to 55%. When inhaling helium the male singer was able reach Eb5, a minor third over the normal limit for males, this seems to indicate that the vocal tract impedance is at least partially responsible for the pitch limit in ‘Overdrive’.

### References

[1] Sadolin C. Complete vocal technique. *Shout*, 2000.

[2] Smith J. Wolfe J. Epps, J. A novel instrument to measure acoustic resonances of the vocal tract during phonation. *Measurement Science and Technology*, 8, 1112., 1997.