



How about a Bayesian M/EEG imaging method correcting for incomplete spatio-temporal priors

Stahlhut, Carsten; Attias, Hagai T.; Sekihara, Kensuke; Wipf, David; Hansen, Lars Kai; Nagarajan, Srikantan S.

Published in:
Journal of Cognitive Neuroscience

Publication date:
2013

[Link back to DTU Orbit](#)

Citation (APA):
Stahlhut, C., Attias, H. T., Sekihara, K., Wipf, D., Hansen, L. K., & Nagarajan, S. S. (2013). How about a Bayesian M/EEG imaging method correcting for incomplete spatio-temporal priors. *Journal of Cognitive Neuroscience, Supplement*, 260.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

How about a Bayesian M/EEG imaging method correcting for incomplete spatio-temporal priors?

Carsten Stahlhut¹, Hagai T. Attias², Kensuke Sekihara³, David Wipf⁴
Lars K. Hansen¹, Srikantan S. Nagarajan⁵

¹Technical University of Denmark, DTU Informatics, Denmark

²Convex Imaging, San Francisco, CA, USA

³Tokyo Metropolitan University, Dept. of System Design & Engineering, Japan

⁴Visual Computing Group, Microsoft Research Asia, China

⁵UCSF, Biomagnetic Imaging Laboratory, San Francisco, CA, USA

Abstract

In this contribution we present a hierarchical Bayesian model, sAquavit, to tackle the highly ill-posed problem that follows with MEG and EEG source imaging. Our model facilitates spatio-temporal patterns through the use of both spatial and temporal basis functions. While in contrast to most previous spatio-temporal inverse M/EEG models, the proposed model benefits of consisting of two source terms, namely, a spatio-temporal pattern term limiting the source configuration to a spatio-temporal subspace and a source correcting term to pick up source activity not covered by the spatio-temporal prior belief.

We have tested the model on both artificial data and real EEG data in order to demonstrate the efficacy of the model. The model was tested at different SNRs (-10.0, -5.2, -3.0, -1.0, 0, 0.8, 3.0 dB) using white noise. At all SNRs the sAquavit performs best in AUC measure, e.g. at SNR=0dB AUC is, 0.985 (sAquavit) and 0.857 (Bolstad et al., 2009).

Our results demonstrate that the sAquavit model is capable in balancing spatio-temporal prior guidance and source correction estimation to obtain superior estimates relative to current inverse methods.