



Impact of CryoSat-2 for marine gravity field - globally and in the Arctic Ocean

Andersen, Ole Baltazar; Stenseng, Lars; Knudsen, Per; Jain, Maulik

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Andersen, O. B., Stenseng, L., Knudsen, P., & Jain, M. (2012). Impact of CryoSat-2 for marine gravity field - globally and in the Arctic Ocean. Abstract from 2012 AGU Fall Meeting, San Francisco, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

[AGU Abstract Browser](#) Beta

- [About](#)
- [Meetings](#)
- [Sections](#)
- [Index Terms](#)
- [Advanced Search](#)

Quick Search

Impact of CryoSat-2 for marine gravity field - globally and in the Arctic Ocean

Details

Meeting [2012 Fall Meeting](#)
Section [Cryosphere](#)
Session [CryoSat-2: Science and Validation I Posters](#)
Identifier C51A-0761
Authors [Andersen, O B*](#), [DTU Space, Kgs Lyngby, Denmark](#)
[Stenseng, L, DTU Space, Kgs Lyngby, Denmark](#)
[Knudsen, P, DTU Space, Kgs Lyngby, Denmark](#)
[Jain, M, DTU Space, Kgs Lyngby, Denmark](#)
[Remote sensing \[0758\]](#)
Index Terms [Gravity anomalies and Earth structure \[1219\]](#)
[Satellite geodesy: results \[1240\]](#)

Abstract

CryoSat-2 offers at least four important improvements compared with conventional altimetry which should benefit marine gravity in general but particularly at high latitude. Firstly the ESA CryoSat-2 signal to noise ratio should be a factor of two better than conventional altimetry. Secondly the 369 days repeat offered by CryoSat-2 provides denser coverage than older geodetic mission data set like ERS-1. Thirdly, the 92 degree inclination of CryoSat-2 is designed to map more of the Arctic Ocean than previous altimetric satellites. Finally, CryoSat-2 is able to operate in two new modes (SAR and SAR-in) designed to improve the sea surface /ice height mapping in the Polar Regions. We have investigated the use of two years of CryoSat-2 LRM data as well as CryoSat SAR and SAR-in data for deriving a global gravity field as well as a regional marine gravity field in the Arctic. Both conventional ESA Level 2 GDR data, NOAA LRM data, but also Level1b (LRM, SAR and SAR-in waveforms) data have been analyzed. A suite of eight different empirical retracers have been developed and investigated for their ability to predict marine gravity in the Arctic Ocean. The impact of the various improvement offered by CryoSat-2 in comparison with conventional satellite altimetry have been studied and quantified both globally but particularly for the Arctic Ocean using a large number of marine and airborne surveys providing “ground truth” marine gravity.

Cite as: Author(s) (2012), Title, Abstract C51A-0761 presented at 2012 Fall Meeting, AGU, San Francisco, Calif., 3-7 Dec.