



Updated vegetation information in high resolution WRF simulations

Nielsen, Joakim Refslund; Dellwik, Ebba; Hahmann, Andrea N.; Boegh, Eva

Published in:

Climate and Land Surface Changes in Hydrology. Proceedings

Publication date:

2013

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Nielsen, J. R., Dellwik, E., Hahmann, A. N., & Boegh, E. (2013). Updated vegetation information in high resolution WRF simulations. In *Climate and Land Surface Changes in Hydrology. Proceedings* (pp. 233-238). IAHS Press. IAHS Publications Series (Red Books), No. 359

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Updated vegetation information in high resolution WRF simulations

JOAKIM REFLUND¹, EBBA DELLWIK¹, ANDREA N HAHMANN¹ & EVA BOEGH²

*1 DTU Wind Energy, Frederiksborgvej 399, DK-4000 Roskilde, Denmark
jref@dtu.dk*

2 Roskilde University, Universitetsvej 1, 02, DK-4000 Roskilde, Denmark

Abstract Climate studies show that the frequency of heat wave events and above-average high temperatures during the summer months over Europe will increase in the coming decades. Such climatic changes and long-term meteorological conditions will impact the seasonal development of vegetation and ultimately modify the energy distribution at the land surface. In weather and climate models it is important to represent the vegetation variability accurately to obtain reliable results. The weather research and forecasting (WRF) model uses green vegetation fraction (GVF) time series to represent vegetation seasonality. The GVF of each grid cell is additionally used to scale other parameters such as LAI, roughness, emissivity and albedo within predefined intervals. However, the default GVF used by WRF does not reflect recent climatic changes or change in management practices since it was derived more than 20 years ago. In this study, a new high resolution GVF product based on MODIS images is applied in a high resolution WRF simulation over Denmark during the 2006 heat wave year. It is found that the GVF is very different in a heat wave year compared to the default GVF. The simulation is compared to a control run using the default GVF data and their performances are quantified against gridded data. The verification includes 2-m temperature and precipitation. The results show that although the simulation using the new GVF product performs well, it does not significantly improve performance compared to the default GVF, despite significant differences in vegetation fractions.

Key words green vegetation fraction; WRF; heat wave