



## EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA); Scientific Opinion on Dietary Reference Values for protein

**EFSA Publication; Tetens, Inge**

*Link to article, DOI:*  
[10.2903/j.efsa.2012.2557](https://doi.org/10.2903/j.efsa.2012.2557)

*Publication date:*  
2012

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
EFSA Publication (2012). EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA); Scientific Opinion on Dietary Reference Values for protein. Parma, Italy: European Food Safety Authority. (The EFSA Journal; No. 2557, Vol. 10(2)). DOI: 10.2903/j.efsa.2012.2557

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## SCIENTIFIC OPINION

### Scientific Opinion on Dietary Reference Values for protein<sup>1</sup>

#### EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA)<sup>2,3</sup>

European Food Safety Authority (EFSA), Parma, Italy

#### ABSTRACT

This opinion of the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) deals with the setting of Dietary Reference Values (DRVs) for protein. The Panel concludes that a Population Reference Intake (PRI) can be derived from nitrogen balance studies. Several health outcomes possibly associated with protein intake were also considered but data were found to be insufficient to establish DRVs. For healthy adults of both sexes, the average requirement (AR) is 0.66 g protein/kg body weight per day based on nitrogen balance data. Considering the 97.5<sup>th</sup> percentile of the distribution of the requirement and assuming an efficiency of utilisation of dietary protein for maintenance of 47 %, the PRI for adults of all ages was estimated to be 0.83 g protein/kg body weight per day and is applicable both to high quality protein and to protein in mixed diets. For children from six months onwards, age-dependent requirements for growth estimated from average daily rates of protein deposition and adjusted by a protein efficiency for growth of 58 % were added to the requirement for maintenance of 0.66 g/kg body weight per day. The PRI was estimated based on the average requirement plus 1.96 SD using a combined SD for growth and maintenance. For pregnancy, an intake of 1, 9 and 28 g/d in the first, second and third trimesters, respectively, is proposed in addition to the PRI for non-pregnant women. For lactation, a protein intake of 19 g/d during the first six months, and of 13 g/d after six months, is proposed in addition to the PRI for non-lactating women. Data are insufficient to establish a Tolerable Upper Intake Level (UL) for protein. Intakes up to twice the PRI are regularly consumed from mixed diets by some physically active and healthy adults in Europe and are considered safe.

© European Food Safety Authority, 2012

#### KEY WORDS

Protein, amino acids, nitrogen balance, factorial method, efficiency of utilisation, digestibility, health outcomes.

<sup>1</sup> On request from the European Commission, Question No EFSA-Q-2008-468, adopted on 19 January 2012.

<sup>2</sup> Panel members: Carlo Agostoni, Jean-Louis Bresson, Susan Fairweather-Tait, Albert Flynn, Ines Golly, Hannu Korhonen, Pagona Lagiou, Martinus Løvik, Rosangela Marchelli, Ambroise Martin, Bevan Moseley, Monika Neuhäuser-Berthold, Hildegard Przyrembel, Seppo Salminen, Yolanda Sanz, Sean (J.J.) Strain, Stephan Strobel, Inge Tetens, Daniel Tomé, Hendrik van Loveren and Hans Verhagen. Correspondence: [nda@efsa.europa.eu](mailto:nda@efsa.europa.eu)

<sup>3</sup> Acknowledgement: The Panel wishes to thank the members of the WG on Population Reference Intakes: Carlo Agostoni, Jean-Louis Bresson, Susan Fairweather-Tait, Albert Flynn, Ambroise Martin, Monika Neuhäuser-Berthold, Hildegard Przyrembel, Sean (J.J.) Strain, Inge Tetens, Daniel Tomé and EFSA's staff member Anja Brønstrup for the preparatory work on this scientific opinion.

## SUMMARY

Following a request from the European Commission, the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) was asked to deliver a scientific opinion on Population Reference Intakes for the European population, including protein.

Dietary proteins are the source of nitrogen and indispensable amino acids which the body requires for tissue growth and maintenance. The main pathway of amino acid metabolism is protein synthesis. In this opinion, “protein” is total nitrogen x 6.25 and protein requirements are based on nitrogen content. Protein digestion takes place in the stomach and in the small intestine. In healthy humans, the absorption and transport of amino acids is usually not limited by the availability of digestive enzymes or transport mechanisms, but some protein escapes digestion in the small intestine and is degraded in the colon through bacterial proteolysis and amino acid catabolism. By the time digesta are excreted as faeces, they consist largely of microbial protein. Therefore, when assessing protein digestibility, it is important to distinguish between faecal and ileal digestibility, as well as apparent and true nitrogen and amino acid digestibility.

The concept of protein requirement includes both total nitrogen and indispensable amino acid requirements. The quantity and utilisation of indispensable amino acids is considered to be an indicator of the dietary protein quality, which is usually assessed using the Protein Digestibility-Corrected Amino Acid Score (PD-CAAS). It is important to determine to what extent the nitrogen from dietary protein is retained in the body. Different values for the efficiency of protein utilisation have been observed for maintenance and for tissue deposition/growth; at maintenance, the efficiency of nitrogen utilisation for retention is about 47 % in healthy adults in nitrogen balance on mixed diets.

Foods of animal origin with a high protein content are meat, fish, eggs, milk and dairy products. Bread and other grain-based products, leguminous vegetables, and nuts are plant foods high in protein. Most of the animal sources are considered high quality protein having an optimal indispensable amino acid composition for human needs and a high digestibility, whereas the indispensable amino acid content of plant proteins and/or their digestibility is usually lower. In European countries the main contributors to dietary protein intake are meat and meat products, grains and grain-based products, and milk and dairy products.

Data from dietary surveys show that the average protein intakes in European countries vary between 67 to 114 g/d in adult men and 59 to 102 g/d in women, or about 12 to 20 % of total energy intake (E %) for both sexes. Few data are available for the mean protein intakes on a body weight basis, which vary from 0.8 to 1.25 g/kg body weight per day for adults.

In order to derive Dietary Reference Values (DRVs) for protein the Panel decided to use the nitrogen balance approach to determine protein requirements. Nitrogen balance is the difference between nitrogen intake and the amount lost in urine, faeces, via the skin and other routes. In healthy adults who are in energy balance the protein requirement (maintenance requirement) is defined as that amount of dietary protein sufficient to achieve zero nitrogen balance. The requirement for dietary protein is considered to be the amount needed to replace obligatory nitrogen losses, after adjustment for the efficiency of dietary protein utilisation and the quality of the dietary protein. The factorial method is used to calculate protein requirements for physiological conditions such as growth, pregnancy or lactation in which nitrogen is not only needed for maintenance but also for the deposition of protein in newly formed tissue or secretions (milk).

According to a meta-analysis of available nitrogen balance data as a function of nitrogen intake in healthy adults, the best estimate of average requirement for healthy adults was 105 mg N/kg body weight per day (0.66 g high quality protein/kg per day). The 97.5<sup>th</sup> percentile was estimated as 133 mg N/kg body weight per day (0.83 g high quality protein/kg per day) from the distribution of the logarithm of the requirement, with a coefficient of variation (CV) of about 12 %. The Panel considers that the value of 0.66 g/kg body weight per day can be accepted as the Average Requirement (AR) and the value of 0.83 g/kg body weight per day as the Population Reference Intake (PRI) derived for

proteins with a PD-CAAS value of 1.0. This value can be applied to usual mixed diets in Europe which are unlikely to be limiting in their content of indispensable amino acids. For older adults, the protein requirement is considered to be equal to that for adults. The lower energy requirement of sedentary elderly people means that the protein to energy ratio of their requirement may be higher than for younger age groups.

For infants, children and adolescents, the Panel accepted the approach of WHO/FAO/UNU (2007) in which estimates of the protein requirements from six months to adulthood were derived factorially as the sum of requirements for maintenance and growth corrected for efficiency of protein utilisation. An average maintenance value of 0.66 g protein/kg body weight per day was applied. Average daily needs for dietary protein for growth were estimated from average daily rates of protein deposition, calculated from studies on whole-body potassium deposition, and from an efficiency of utilisation of dietary protein for growth of 58 %. The PRI was estimated based on the average requirement plus 1.96 SD using a combined SD for growth and maintenance.

For pregnant women, the Panel accepted the factorial method for deriving protein requirements during pregnancy which was based on the newly deposited protein in the foetus and maternal tissue, and on the maintenance requirement associated with the increased body weight. Because of the paucity of data in pregnant women and because it is unlikely that the efficiency of protein utilisation decreases during pregnancy, the efficiency of protein utilisation was taken to be 47 % as in non-pregnant women. Thus, for pregnant women a PRI for protein of 1, 9 and 28 g/d in the first, second and third trimesters, respectively, is proposed in addition to the PRI for non-pregnant women.

For lactation, the Panel accepted the factorial method which requires assessing milk volumes produced and the content of both protein nitrogen and non-protein nitrogen, as well as calculating the amount of dietary protein needed for milk protein production. As the efficiency of protein utilisation for milk protein production is unknown, the same efficiency as in the non-lactating adult (47 %) was assumed. The PRI was estimated by adding 1.96 SD to give an additional 19 g protein/d during the first six months of lactation (exclusive breastfeeding), and 13 g protein/d after six months (partial breastfeeding).

The Panel also considered several health outcomes that may be associated with protein intake. The available data on the effects of an additional dietary protein intake beyond the PRI on muscle mass and function, on body weight control and obesity (risk) in children and adults, and on insulin sensitivity and glucose homeostasis do not provide evidence that can be considered as a criterion for determining DRVs for protein. Likewise, the available evidence does not permit the conclusion that an additional protein intake might affect bone mineral density and could be used as a criterion for the setting of DRVs for protein.

Data from food consumption surveys show that actual mean protein intakes of adults in Europe are at, or more often above, the PRI of 0.83 g/kg body weight per day. In Europe, adult protein intakes at the upper end (90-97.5<sup>th</sup> percentile) of the intake distributions have been reported to be between 17 and 27 E%. The available data are not sufficient to establish a Tolerable Upper Intake Level (UL) for protein. In adults an intake of twice the PRI is considered safe.

DRVs have not been derived for indispensable amino acids since amino acids are not provided as individual nutrients but in the form of protein. In addition, the Panel notes that more data are needed to obtain sufficiently precise values for indispensable amino acid requirement.

**TABLE OF CONTENTS**

Abstract .....	1
Summary .....	2
Table of contents .....	4
Background as provided by the European Commission.....	6
Terms of reference as provided by European Commission.....	6
Assessment.....	8
1. Introduction .....	8
2. Definition / category .....	8
2.1. Definition .....	8
2.2. Protein digestion and metabolism.....	9
2.2.1. Intestinal protein digestion and amino acid absorption .....	9
2.2.2. Protein turnover, amino acid metabolism and amino acid losses .....	10
2.3. Protein quality from digestibility and indispensable amino acid composition .....	10
2.3.1. Measurement of protein digestibility.....	10
2.3.2. The indispensable amino acid scoring method.....	11
2.4. Nitrogen retention and efficiency of dietary protein utilisation.....	11
3. Dietary protein sources and intake data.....	12
3.1. Nitrogen and protein content in foodstuffs – the nitrogen conversion factor .....	12
3.2. Dietary sources.....	13
3.3. Dietary intake.....	14
4. Overview of Dietary Reference Values and recommendations.....	15
4.1. Dietary Reference Values and recommendations for protein for adults .....	15
4.1.1. Older adults.....	16
4.2. Dietary Reference Values and recommendations for protein for infants and children .....	16
4.3. Dietary Reference Values and recommendations for protein during pregnancy .....	19
4.4. Dietary Reference Values and recommendations for protein during lactation .....	19
4.5. Requirements for indispensable amino acids.....	20
5. Criteria (endpoints) on which to base Dietary Reference Values (DRVs).....	21
5.1. Protein intake and protein and nitrogen homeostasis.....	21
5.1.1. Methods for the determination of protein requirement.....	21
5.1.1.1. Nitrogen balance .....	21
5.1.1.2. Indicator amino acid oxidation method.....	21
5.1.1.3. The factorial method .....	21
5.1.1.4. Protein quality and reference pattern for indispensable amino acids.....	22
5.1.2. Protein requirement of adults.....	23
5.1.2.1. Older adults.....	23
5.1.3. Protein requirement of infants and children .....	24
5.1.4. Protein requirement during pregnancy .....	25
5.1.5. Protein requirement during lactation .....	26
5.2. Protein intake and health consequences.....	26
5.2.1. Muscle mass.....	26
5.2.2. Body weight control and obesity .....	27
5.2.2.1. Infants .....	27
5.2.2.2. Adults.....	28
5.2.3. Insulin sensitivity and glucose control.....	28
5.2.4. Bone health .....	28
5.2.5. Kidney function .....	29
5.2.6. Capacity of the urea cycle.....	29
5.2.7. Tolerance of protein.....	30
6. Data on which to base Dietary Reference Values (DRVs).....	30
6.1. Protein requirement of adults .....	30
6.2. Protein requirement of infants and children.....	30
6.3. Protein requirement during pregnancy.....	31

6.4. Protein requirement during lactation.....	31
6.5. Safety of protein intakes above the PRI.....	31
Conclusions .....	32
References .....	33
Appendix 1: Main food contributors to dietary protein intake (%) of adults (18-64 years) in European countries as estimated with the EFSA Comprehensive European Food Consumption Database.....	48
Appendix 2a: Population, methods and period of dietary assessment in children and adolescents in European countries .....	49
Appendix 2b: Protein intake of children aged ~1-3 years in European countries .....	52
Appendix 2c: Protein intake of children aged ~4-6 years in European countries .....	53
Appendix 2d: Protein intake of children aged ~7-9 years in European countries .....	54
Appendix 2e: Protein intake of children aged ~10-14 years and over in European countries.....	55
Appendix 2f: Protein intake of adolescents aged ~15-18 years and over in European countries.....	56
Appendix 3a: Population, methods and period of dietary assessment in adults in European countries .....	57
Appendix 3b: Protein intake of adults aged ~19-65 years in European countries.....	60
Appendix 3c: Protein intake of adults aged ~19-34 years in European countries .....	61
Appendix 3d: Protein intake of adults aged ~35-64 years in European countries.....	62
Appendix 3e: Protein intake of adults aged ~65 years and over in European countries .....	63
Appendix 4: Calculation of PRI for infants, children and adolescents.....	64
Glossary / Abbreviations .....	65

## BACKGROUND AS PROVIDED BY THE EUROPEAN COMMISSION

The scientific advice on nutrient intakes is important as the basis of Community action in the field of nutrition, for example such advice has in the past been used as the basis of nutrition labelling. The Scientific Committee for Food (SCF) report on nutrient and energy intakes for the European Community dates from 1993. There is a need to review and if necessary to update these earlier recommendations to ensure that the Community action in the area of nutrition is underpinned by the latest scientific advice.

In 1993, the SCF adopted an opinion on nutrient and energy intakes for the European Community<sup>4</sup>. The report provided Reference Intakes for energy, certain macronutrients and micronutrients, but it did not include certain substances of physiological importance, for example dietary fibre.

Since then new scientific data have become available for some of the nutrients, and scientific advisory bodies in many European Union Member States and in the United States have reported on recommended dietary intakes. For a number of nutrients these newly established (national) recommendations differ from the reference intakes in the SCF (1993) report. Although there is considerable consensus between these newly derived (national) recommendations, differing opinions remain on some of the recommendations. Therefore, there is a need to review the existing EU Reference Intakes in the light of new scientific evidence, and taking into account the more recently reported national recommendations. There is also a need to include dietary components that were not covered in the SCF opinion of 1993, such as dietary fibre, and to consider whether it might be appropriate to establish reference intakes for other (essential) substances with a physiological effect.

In this context the EFSA is requested to consider the existing Population Reference Intakes for energy, micro- and macronutrients and certain other dietary components, to review and complete the SCF recommendations, in the light of new evidence, and in addition advise on a Population Reference Intake for dietary fibre.

For communication of nutrition and healthy eating messages to the public it is generally more appropriate to express recommendations for the intake of individual nutrients or substances in food-based terms. In this context the EFSA is asked to provide assistance on the translation of nutrient based recommendations for a healthy diet into food based recommendations intended for the population as a whole.

## TERMS OF REFERENCE AS PROVIDED BY EUROPEAN COMMISSION

In accordance with Article 29 (1)(a) and Article 31 of Regulation (EC) No. 178/2002, the Commission requests EFSA to review the existing advice of the Scientific Committee for Food on Population Reference Intakes for energy, nutrients and other substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

In the first instance the EFSA is asked to provide advice on energy, macronutrients and dietary fibre. Specifically advice is requested on the following dietary components:

- Carbohydrates, including sugars;
- Fats, including saturated fatty acids, poly-unsaturated fatty acids and mono-unsaturated fatty acids, *trans* fatty acids;
- Protein;
- Dietary fibre.

---

<sup>4</sup> Scientific Committee for Food, Nutrient and energy intakes for the European Community, Reports of the Scientific Committee for Food 31st series, Office for Official Publication of the European Communities, Luxembourg, 1993.

Following on from the first part of the task, the EFSA is asked to advise on Population Reference Intakes for micronutrients in the diet and, if considered appropriate, other essential substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

Finally, the EFSA is asked to provide guidance on the translation of nutrient based dietary advice into guidance, intended for the European population as a whole, on the contribution of different foods or categories of foods to an overall diet that would help to maintain good health through optimal nutrition (food-based dietary guidelines).



## ASSESSMENT

### 1. Introduction

Dietary proteins are an essential component of the diet by supplying the body with nitrogen (N) and amino acids which are used to synthesise and maintain the around 25,000 proteins encoded within the human genome as well as other non-protein metabolically active nitrogenous substances like peptide hormones, neurotransmitters, nucleic acids, glutathione or creatine. In addition, amino acids are also subjected to deamination and their carbon skeleton is used in different metabolic pathways or as energy substrate.

### 2. Definition / category

#### 2.1. Definition

Proteins are built from amino acids joined together by peptide bonds between the carboxyl and the amino (or imino in the case of proline) group of the next amino acid in line. These polypeptide chains are folded into a three dimensional structure to form the protein. The primary structure or sequence of amino acids in proteins is pre-determined in the genetic code. Twenty of the naturally occurring amino acids are so-called proteinogenic amino acids which build proteins in living organisms. With few exceptions, only L-isomers are incorporated into proteins.

Dietary proteins are the source of nitrogen and indispensable amino acids for the body. Both in the diet and in the body, 95 % of the nitrogen is found in the form of proteins and 5 % is found in the form of other nitrogenous compounds, i.e. free amino acids, urea or nucleotides. A conversion factor of 6.25 is usually used for the conversion of nitrogen to protein for labelling purposes, assessment of protein intake, and for protein reference values. Total N x 6.25 is called crude protein and [total minus non-protein-N] x 6.25 is called true protein. For other purposes, protein specific nitrogen conversion factors can be used (see Section 3.1.). In this opinion, unless specifically mentioned, "protein" is total N x 6.25 and protein requirements are calculated from nitrogen content.

The 20 proteinogenic amino acids are classified as indispensable or dispensable amino acids. Nine amino acids are classified as indispensable in humans (histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine) as they cannot be synthesised in the human body from naturally occurring precursors at a rate to meet the metabolic requirement. The remaining dietary amino acids are dispensable (alanine, arginine, cysteine, glutamine, glycine, proline, tyrosine, aspartic acid, asparagine, glutamic acid, and serine). Among the nine indispensable amino acids, lysine and threonine are strictly indispensable since they are not transaminated and their deamination is irreversible. In contrast, the seven other indispensable amino acids can participate in transamination reactions. In addition, some of the dispensable amino acids which can under normal physiological conditions be synthesised in the body, can become limiting under special physiological or pathological conditions, such as in premature neonates when the metabolic requirement cannot be met unless these amino acids are supplied in adequate amounts with the diet; they are then called conditionally indispensable amino acids (arginine, cysteine, glutamine, glycine, proline, tyrosine) (IoM, 2005; NNR, 2004).

Besides being a building block for protein synthesis, each amino acid has its own non-proteogenic metabolic pathways. Some amino acids are used as precursors for nitrogenous compounds such as glutathione, various neurotransmitters, nitrogen monoxide, creatine, carnitine, taurine or niacin. Glutamine, aspartate and glycine are used for the synthesis of ribo- and deoxyribonucleotides, precursors for the synthesis of the nucleic acids RNA and DNA. Arginine and glutamine are precursors of non-proteinogenic amino acids including ornithine and citrulline that play a role in inter-organ exchange of nitrogen. Glutamine and glutamate are precursors of Krebs cycle components and are also important energy substrates for various cells. Amino acids are used after deamination as energy substrates and in gluconeogenesis and ketogenesis. Some of the amino acids can also directly or indirectly act as intracellular signal molecules. Glutamate is a well known neurotransmitter, tryptophan is the precursor of serotonin, tyrosine is the precursor of catecholamines and dopamine, as well as of thyroid hormones, and histidine is the precursor of histamine. Arginine is an activator of the first step of  $\text{NH}_4^+/\text{NH}_3$  elimination in the hepatic urea cycle, acts as a secretagogue for  $\beta$ -cells of pancreatic Langerhans

islets, and is - via nitric oxide synthase activity - the precursor of nitrogen monoxide that regulates blood pressure. Lastly, leucine has been subjected to numerous studies for its role as a signal for protein synthesis via the mTOR (mammalian target of rapamycin) signalling pathway. These non-proteogenic metabolic pathways and signalling activities are included in the concept of protein requirement when nitrogen balance is achieved and indispensable amino acid requirements are met. As a consequence, they are not used as additional markers for the determination of protein requirement.

## 2.2. Protein digestion and metabolism

Protein metabolism comprises the processes that regulate protein digestion, amino acid metabolism and body protein turnover. These processes include the absorption and supply of both dispensable and indispensable dietary amino acids and the *de novo* synthesis of dispensable amino acids, protein hydrolysis, protein synthesis, and amino acid utilisation in catabolic pathways or as precursors for nitrogenous compounds.

### 2.2.1. Intestinal protein digestion and amino acid absorption

The fluxes of nitrogen, amino acids and protein in the gut exhibit a rather complex pattern. In humans, ingested dietary proteins (about 40–110 g/d), endogenous protein secreted into the gut (20–50 g/d) and molecules containing non-protein nitrogen (urea and other molecules) secreted into the gut are mixed in the lumen of the stomach and the small intestine and are subjected to transit, digestion and absorption (Gaudichon et al., 2002). The main part is transferred into the body by absorption across the intestinal mucosa whereas a smaller part remains in the lumen and reaches the terminal ileum. This, along with other undigested luminal components, passes from the terminal ileum into the large intestine, and the whole is subjected to fermentation by the microflora.

Protein digestion starts in the stomach and is continued in the small intestine. In healthy humans, digestive enzymes and the transport across the brush border membrane through a variety of transporters are not limiting factors for amino acid absorption (Johnson et al., 2006). The metabolic activity of the small intestine is high and the small intestinal mucosa metabolises a significant proportion of both dispensable and indispensable amino acids in the course of absorption. In the absorptive state, dietary rather than systemic amino acids are the major precursors for mucosal protein synthesis. Glutamine and glutamate, which are the most important fuels for intestinal tissue, are mostly used by the intestine, and their appearance in the portal circulation is usually very low. Fifty to sixty percent of dietary threonine is used by the intestine mainly for mucin synthesis by goblet cells. Of the amino acids lysine, leucine or phenylalanine, 15–30 % is used by the intestine whereas the other fraction appears in the portal circulation. Catabolism dominates the intestinal utilisation of dietary amino acids, since only 12 % of the amino acids extracted by the intestine are used for mucosal protein synthesis.

Approximately 15 g protein/d remains in the intestinal lumen and enters the colon. There it is degraded into peptides and amino acids through bacterial proteolysis, and amino acids are further deaminated and decarboxylated. This process is considered to be a major pathway for amino acid losses at maintenance intake of dietary protein (Gaudichon et al., 2002). The microflora possesses ureolytic activity so that nitrogen of urea secreted into the intestine can be recycled both by microbial amino acid synthesis and by the uptake of ammonia from the gut. The ammonia is captured especially into alanine, aspartate/asparagine and glutamate/glutamine, from which it may be incorporated into most amino acids by transamination. This mechanism of urea recycling might be of value in conserving nitrogen (Fouillet et al., 2008; Jackson, 1995).

As a consequence of the activities of the intestinal microbiota, by the time digesta are excreted as faeces their protein content is largely of microbial origin. Therefore, faecal or ileal digestibility measurements, as well as apparent and true nitrogen and amino acid digestibility measurements (see Section 2.3.1.), have very different significance and can be used for different objectives. Measurements at the ileal level are critical for determining amino acid losses of both dietary and endogenous origin, whereas measurements at the faecal level are critical in assessing whole-body nitrogen losses (Fuller and Tome, 2005). The impact of the recycling of intestinal nitrogen, and of amino acids synthesised by bacteria, on whole-body requirement of nitrogen, amino acids and protein is not clear. Other bacteria-derived amino acid metabolites include short

chain fatty acids, sulphides, ammonia, phenols or indoles. The health consequences of changes in the luminal concentration of these products have not been extensively studied.

### 2.2.2. Protein turnover, amino acid metabolism and amino acid losses

The main pathway of amino acid metabolism is protein synthesis. In a 70 kg adult man, the body protein pool represents 10-12 kg, of which 42 % is in skeletal muscle, 15 % each in skin and blood, and 10 % in visceral organs. Four proteins (collagen, myosin, actin and haemoglobin) account for half of the body protein pool, and 25 % of the proteins of the body are present as collagen. The 10-12 kg body protein pool is in continuous turnover and exchanges with the free amino acid pool, which is approximately 100 g, via the proteosynthesis and proteolysis pathways at a rate of 250-300 g/d in the 70 kg adult man (Waterlow, 1995, 1996). This protein turnover is 2-3 times higher than the usual dietary protein intake (NNR, 2004). Moreover, the synthesis and turnover rates vary between the different body proteins. Visceral tissues have a fast protein turnover whereas peripheral tissues have a lower rate.

Amino acids are irreversibly lost in the faeces (25-30 % of total amino acid losses), by metabolic oxidation (70-75 % of total amino acid losses) and as miscellaneous losses in urine (about 0.6 g amino acids or 40 mg nitrogen in male adults), hair, skin, bronchial and other secretions, and in lactating women as milk (SCF, 1993). These amino acid losses need to be balanced by the supply of dietary protein-derived amino acids (50-100 g/d). When protein intake is increased the metabolic oxidative losses are also increased in order to achieve amino acid and nitrogen balance (Forslund et al., 1998; Morens et al., 2003; Pacy et al., 1994; Price et al., 1994).

### 2.3. Protein quality from digestibility and indispensable amino acid composition

The nutritional value of dietary proteins is related to their ability to satisfy nitrogen and amino acid requirements for tissue growth and maintenance. According to current knowledge this ability mainly depends on the digestibility of protein and amino acids, and the dispensable and indispensable amino acid composition of the proteins.

#### 2.3.1. Measurement of protein digestibility

The aim of measuring protein digestibility is to predict the quantity of absorbed nitrogen or amino acids following protein consumption. Though several *in vitro* methods requiring enzymatic hydrolysis have been proposed, the classical approach uses *in vivo* digestibility in an animal model or in humans. The classical *in vivo* procedure is based on faecal collection and determination of the nitrogen output for several days. *Apparent digestibility* of protein is measured from the difference between nitrogen ingested and nitrogen excreted in the faeces. It does not take into account the presence of endogenous nitrogen secretion and colonic metabolism. Apparent digestibility is one component in the assessment of whole-body nitrogen losses. For the determination of *true* (or *real*) *digestibility*, discrimination between exogenous nitrogen (food) and endogenous nitrogen losses (secretions, desquamations etc.) is needed. Individual amino acid digestibility is usually related to whole protein nitrogen digestibility. Alternatively, individual amino acid digestibility can be determined.

Both direct and indirect methods have been proposed to distinguish and quantify the endogenous and dietary components of nitrogen and amino acids in ileal chyme or faeces. These approaches include the administration of a protein-free diet, the enzyme-hydrolysed protein method, different levels of protein intake, or multiple regression methods, in which it is assumed that the quantity and amino acid composition of endogenous losses is constant and independent of diet (Baglieri et al., 1995; Fuller and Reeds, 1998; Fuller and Tome, 2005). Substantial advances in the ability to discriminate between exogenous (dietary) and endogenous nitrogen have been achieved using stable isotopes (Fouillet et al., 2002). By giving diets that are isotopically labelled (usually carbon or nitrogen of amino acids), the endogenous flow is estimated from the dilution of the isotopic enrichment in the digesta (Fouillet et al., 2002; Gaudichon et al., 1999; Tome and Bos, 2000). Regarding the dietary amino acid fraction, it is also questionable whether protein (overall nitrogen) digestibility is a good proxy for individual ileal amino acid digestibility because some studies have reported modest ranges of variation of individual amino acid digestibility around the value for nitrogen

digestibility (Fuller and Tome, 2005). It appears that in some cases there are substantial differences in true digestibility among amino acids (Fouillet et al., 2002; Gaudichon et al., 2002; Tome and Bos, 2000).

The unabsorbed amino acids are mostly metabolised by colonic bacteria. Therefore, the apparent digestibility measured in ileal effluent should be considered as a critical biological parameter for dietary amino acid digestibility (Fuller and Tome, 2005). Digestibility values obtained by the faecal analysis method usually overestimate those obtained by the ileal analysis method. In humans, intestinal effluents for the estimation of apparent digestibility are obtained either from ileostomy patients or, preferably, in healthy volunteers by using naso-intestinal tubes. These approaches are not, however, straightforward, and are too demanding for the routine evaluation of food, but can be used as reference methods (Fouillet et al., 2002; Fuller et al., 1994). An alternative is the use of animal models, most commonly the rat and the pig. The rat is used for the determination of protein quality in human diets (FAO/WHO, 1991). However, differences in protein digestibility have been observed between rats, pigs and humans (Fuller and Tome, 2005).

The usefulness of the values obtained by digestibility measurements depends on the objective. *In vitro* digestibility measurements can only be used to compare products with one another, and can never serve as independent reference values. Measurement of apparent and real digestibility is critical for determining amino acid losses of both dietary and endogenous origin. Data in humans are preferred whenever possible. The determination of individual amino acid digestibility is also preferred whenever possible. An unresolved aspect of digestibility assessments is how to take into account the recycling of intestinal nitrogen and bacterial amino acids to the body.

### 2.3.2. The indispensable amino acid scoring method

The concept of protein requirement includes both total nitrogen and indispensable amino acids requirements. Therefore, the content and utilisation of indispensable amino acids can be considered as valuable criteria for the evaluation of dietary protein quality (WHO/FAO/UNU, 2007). This idea leads to the use of the amino acid scoring approach in which the indispensable amino acid composition of the dietary protein is compared to a reference pattern of indispensable amino acids which is assumed to meet requirements for indispensable amino acids at a protein supply which corresponds to the average protein requirement. The reference pattern of indispensable amino acids is derived from measurements of indispensable amino acid requirements (WHO/FAO/UNU, 2007) (see Section 4.5.). Originally, the chemical score was based on the complete analysis of the food amino acid content and its comparison to the amino acid pattern of a chosen reference protein (e.g. egg or milk protein).

In the traditional scoring method, the ratio between the content in a protein and the content in the reference pattern is determined for each indispensable amino acid. The lowest value is used as the score. The Protein Digestibility-Corrected Amino Acid Score (PD-CAAS) corrects the amino acid score by the digestibility of the protein (FAO/WHO, 1991) or of each individual amino acid. The accuracy of the scoring approach depends on the precision of amino acid analysis and on the measurement of protein digestibility. A more precise approach is to use the specific ileal digestibility of individual amino acids. The PD-CAAS can be used as a criterion for the protein quality of both foods and diets. A PD-CAAS  $<1$  indicates that at least one amino acid is limiting, whereas a score  $\geq 1$  indicates that there is no limiting amino acid in the food or diet.

### 2.4. Nitrogen retention and efficiency of dietary protein utilisation

A traditional approach for evaluating the efficiency of protein utilisation has been to consider the interaction with a physiological process such as growth. The Protein Efficiency Ratio (PER) that relates the average animal (rat) weight gain to the amount of ingested protein over 28 days (Association of Official Analytical Chemists, 1984; Satterlee et al., 1979) is simple but presents several shortcomings and inaccuracies. The main difficulty lies in the significance of extrapolation to humans.

Determination of the nutritional efficiency of protein in the diet is in most cases based on estimating the extent to which dietary protein nitrogen is absorbed and retained by the organism and is able to balance daily nitrogen losses. It is determined by measuring faecal, urinary and miscellaneous nitrogen losses. Net Protein Utilisation (NPU) is the percentage of ingested nitrogen that is retained in the body, and the Biological Value (BV) gives the percentage of absorbed nitrogen that is retained. BV is the product of NPU and digestibility.

As with digestibility, NPU values are true or apparent depending on whether the loss of endogenous nitrogen is taken into account or not, and this is critical to precisely determining the efficiency of dietary protein utilisation and the quality of the different dietary protein sources. The true NPU can be calculated as follows:

$$\text{True NPU} = \frac{\text{total } N_{\text{ingested}} - [(\text{total } N_{\text{faeces}} - \text{endogenous } N_{\text{faeces}}) + (\text{total } N_{\text{urine}} - \text{endogenous } N_{\text{urine}})]}{\text{total } N_{\text{ingested}}}$$

Endogenous intestinal (faecal) and metabolic (urinary) nitrogen losses can be obtained with a protein-free diet, be derived from the y-intercept of the regression line relating nitrogen intake to retention at different levels of protein intake, or be directly determined from experiments using isotopically labelled dietary proteins.

As the post-prandial phase is critical for dietary protein utilisation, the measurement of the immediate retention of dietary nitrogen following meal ingestion represents a reliable approach for the assessment of protein nutritional efficiency. In the net post-prandial protein utilisation (NPPU) approach true dietary protein nitrogen retention is directly measured in the post-prandial phase from experiments using <sup>15</sup>N-labelled dietary proteins (Fouillet et al., 2002). Dietary proteins are considered to have a mean NPPU value of 70 % (Bos et al., 2005). This NPPU approach represents the maximal potential NPU efficiency of the dietary protein sources when determined in optimised controlled conditions in healthy adults, and it can be modified by different factors including food matrix, diet and physiological conditions.

From nitrogen balance studies, an NPU value of 47 % (median value, 95 % CI 44–50 %) was derived from the slope of the regression line relating nitrogen intake to retention for healthy adults at maintenance, and no differences were found between the results when the data were grouped by sex, diet or climate (Rand et al., 2003; WHO/FAO/UNU, 2007). The results suggested a possible age difference in nitrogen utilisation with a lower efficiency in individuals aged above 55 years (31 % compared with 48 % for adults up to 55 years,  $p=0.003$ ), but because of the apparent interaction between age and sex in the data, the extreme variability in the younger men, and the fact that the lower values for the older adults came from a single study, these results were not accepted as conclusive (Rand et al., 2003). Different values are used for efficiency of protein utilisation for maintenance (47 %) and for tissue deposition/growth in different populations and age groups including infants, and pregnant or lactating women (IoM, 2005; King et al., 1973; WHO/FAO/UNU, 2007).

The Panel considers that methods related to growth in the rat (protein efficiency ratio, PER) are not reliable for humans. Methods related to nitrogen retention (NPPU, NPU, BV) are preferable as they reflect more accurately the protein nutritional value, and can be used as reference methods. From available data in healthy adults at maintenance the mean optimal NPU value determined as NPPU is 70 %, and the usual NPU value as determined from nitrogen balance studies is approximately 47 %.

### 3. Dietary protein sources and intake data

#### 3.1. Nitrogen and protein content in foodstuffs – the nitrogen conversion factor

Assuming an average nitrogen content of 160 mg/g protein, a conversion factor of 6.25 is used for the calculation of the (crude) protein content of a food from the total nitrogen content. Specific conversion factors for different proteins have been proposed (Jones, 1941; Leung et al., 1968; Pellett and Young, 1980), including, for instance, milk and milk products (6.38), other animal products (6.25), wheat (5.83) or soy protein (5.71). Besides variations in the nitrogen content of different proteins, the presence or absence of a non-protein fraction of the total nitrogen content of a food will influence the calculated crude protein content (SCF, 2003).

Conversion factors based on the amino acid composition of a protein have been proposed to define more accurately the true protein content of different foodstuffs (AFSSA, 2007; SCF, 2003). The choice of one or several conversion factors depends on the objective, and if the aim is to indicate a product's capacity to supply nitrogen a single coefficient is enough. However, if the objective is to indicate a product's potential to supply amino acids, the use of specific coefficients based on amino acid-derived nitrogen content is more relevant. Such protein amino acid composition-derived conversion factors have been determined for different

protein sources: milk and milk products (5.85), meat, fish and eggs (5.6), wheat and legumes (5.4), and a default conversion factor (5.6) (AFSSA, 2007).

### 3.2. Dietary sources

Dietary proteins are found in variable proportions in different foods resulting in variability of dietary protein intake within and between populations. Proteins differ in their amino acid composition and indispensable amino acid content. Foods of animal origin with a high protein content are meat, fish, eggs, milk and dairy products. Most of these animal dietary protein sources are high in indispensable amino acids. Plant-derived foods with a high protein content are bread and other grain-based products, leguminous vegetables, and nuts. The protein content differs from one plant source to another accounting for 20-30 % (w/w) for uncooked legume seeds or around 10 % for cereal seeds. The indispensable amino acid content of plant proteins is usually lower than in animal proteins. In addition, technological treatments applied to proteins during extraction processes and during the production of foodstuffs may modify the characteristics, properties and nutritional quality of food proteins.

Examples of the range of protein content of some animal- and plant-derived foods are provided in Table 1. The water and energy contents of these foods can greatly differ.

**Table 1:** Protein content (N x 6.25, g/100 g of edible food) of some animal- and plant-derived foods

<b>Animal-derived foods</b>	<b>Protein content (N x 6.25, g/100 g)</b>	<b>Plant-derived foods</b>	<b>Protein content (N x 6.25, g/100 g)</b>
Red meat (raw and cooked)	20-33	Vegetables	1-5
Poultry (raw and cooked)	22-37	Legumes	4-14
Fish	15-25	Fruits	0.3-2
Eggs	11-13	Nuts and seeds	8-29
Cheese, hard	27-34	Pasta and rice (cooked)	2-6
Cheese, soft	12-28	Breads and rolls	6-13
Milk products	2-6	Breakfast cereals	5-13

Data adapted from the ANSES/CIQUAL French food composition table version 2008 (ANSES/CIQUAL, 2008)

In most European countries, the main contributor to the dietary protein intake of adults is meat and meat products, followed by grains and grain-based products, and milk and dairy products. These three food groups contribute to about 75 % of the protein intake (see Appendix 1).

Several methods exist for assessing protein quality, for example the content of indispensable amino acids. One of the food composition tables providing the most detailed amino acid profiles of various foodstuffs is the table of the United States Department of Agriculture (2009). High quality protein has an optimal indispensable amino acid composition for human needs and a high digestibility. Most dietary protein of animal origin (meat, fish, milk and egg) can be considered as such high quality protein. In contrast, some dietary proteins of plant origin can be regarded as being of lower nutritional quality due to their low content in one or several indispensable amino acids and/or their lower digestibility. It is well established that lysine is limiting in cereal protein and that sulphur-containing amino acids (cysteine and methionine) are limiting in legumes. Most of the Western diets have a PD-CAAS equal to or higher than 1 because high quality proteins dominate over low quality proteins. Although proteins limited in one amino acid can complement proteins in the diet which are limited in another amino acid, a high level of cereal in the diet in some countries can lead to a PD-CAAS lower than 1 mainly because of a low content in lysine. For example, as reported in Table 2, most protein from animal sources has a higher PD-CAAS than protein from vegetable sources, but differences also exist within proteins from vegetable sources. For adults, the PD-CAAS value of animal

proteins is usually higher than 1 (but truncated to 1). For plant proteins the PD-CAAS value is close to 1 for soy protein, somewhat lower for other legumes and around 0.5-0.65 for cereal protein.

**Table 2:** Example of values for Protein Digestibility-Corrected Amino Acid Score (PD-CAAS) values of different foods for adults (adapted from (AFSSA, 2007; Michaelsen et al., 2009; WHO/FAO/UNU, 2007))

	PD-CAAS (%)	Limiting amino acid(s)
<b>Animal sources</b>		
Egg	>1.0	-
Milk, cheese	>1.0	-
Meat, fish	>1.0	-
<b>Vegetable sources</b>		
Soy	~0.95	Met+Cys
Beans	~0.7-0.75	Met+Cys
Rice	~0.65	Lys
Wheat	~0.5	Lys
Maize	~0.5	Lys

Due to the high content in indispensable amino acids in animal proteins, a diet rich in animal protein usually has a content of each indispensable amino acid above the requirement. It is widely accepted that a balance between dispensable and indispensable amino acids is a more favourable metabolic situation than a predominance of indispensable amino acids since indispensable amino acids consumed in excess of requirement are either converted to dispensable amino acids or directly oxidised.

### 3.3. Dietary intake

Typical intakes of (crude) protein of children and adolescents from 20 countries (Appendix 2) and of adults from 24 countries in Europe (Appendix 3) are presented. The data refer to individual-based food consumption surveys, conducted from 1989 onwards. Most studies comprise nationally representative population samples.

As demonstrated in the appendices, there is a large diversity in the methodology used to assess the individual intakes of children, adolescents and adults. Because the different methods apply to different time frames, this inevitably results in variability in both the quality and quantity of available data, which makes direct comparison difficult. Moreover, age classifications are in general not uniform. Comparability is also hampered by differences in the food composition tables used for the conversion of food consumption data to estimated nutrient intakes (Deharveng et al., 1999).

Although these differences may have an impact on the accuracy of between country comparisons, the presented data give a rough overview of the protein intake in a number of European countries. Most studies reported mean intakes and standard deviations (SD), or mean intakes and intake distributions. In most studies the contribution of protein to energy intake is based on total energy intake (including energy from alcohol).

In adults, average protein intakes in absolute amounts range from approximately 67 to 114 g/d in men and from 59 to 102 g/d in women. Available data suggest an average intake of 0.8 to 1.25 g/kg body weight per day for adults. Average protein intake varies in infants and young children from about 29 to 63 g/d. Average daily intakes increase with age to about 61 to 116 g/d in adolescents. In general, males have higher intakes than females. Only a few countries present data per kg body weight. However, the estimated mean intakes vary from  $\geq 3$  g/kg body weight per day in the youngest age groups to approximately 1.2 to 2.0 g/kg body weight per day in children and adolescents aged 10-18 years.

When expressed as % of energy intake (E%), average total protein intakes range from about 12 to 20 E% in adults, with within population ranges varying from about 10-15.5 E% at the lower (2.5–10<sup>th</sup> percentile) to about 17-27 E% at the upper (90-97.5<sup>th</sup> percentile) end of the intake distributions. Average intakes of 17 E%

and higher are observed, for example in France, Ireland, Finland, Romania, Portugal and Spain. Available data show that average protein intakes in children and adolescents in European countries vary from about 11 to 18.5 E%. Within population ranges vary from about 6-13 E% (2.5-10<sup>th</sup> percentile) to 14-22 E% (90-97.5<sup>th</sup> percentile).

#### 4. Overview of Dietary Reference Values and recommendations

A number of national and international organisations and authorities have set Dietary Reference Values (DRVs) or recommendations for protein and other energy-providing nutrients, as well as for dietary fibre. Generally, the reference intakes for protein are expressed as g/kg body weight per day and g/d (adjusting for reference body weights), and as percentage of total energy intake (E%), and refer to high quality protein (e.g. milk and egg protein).

##### 4.1. Dietary Reference Values and recommendations for protein for adults

Table 3 lists reference intakes for adult humans set by various organisations.

In its report, FAO/WHO/UNU (1985) used nitrogen balance to derive a population average requirement of 0.6 g/kg body weight per day and, adding two SD (2 x 12.5 %) to allow for individual variability, a “safe level of intake” of 0.75 g/kg body weight per day. UK COMA (DoH, 1991) and SCF (1993) accepted the values adopted by FAO/WHO/UNU (1985). The Netherlands (Health Council of the Netherlands, 2001) also used the approach of FAO/WHO/UNU (1985), but applied a coefficient of variation (CV) of 15 % to allow for individual variability, and derived a recommended intake of 0.8 g/kg body weight per day. The Nordic Nutrition recommendations (NNR, 2004), taking account of the fact that diets in industrialised countries have high protein contents, set a desirable protein intake of 15 E% for food planning purposes with a range of 10-20 E% for adults. This translates into protein intakes of well above 0.8 g/kg body weight per day. The US Institute of Medicine (IoM, 2005) recommended 0.8 g/kg body weight per day of good quality protein for adults. The criterion of adequacy used for the estimated average requirement (EAR) of protein is based on the lowest continuous intake of dietary protein that is sufficient to achieve body nitrogen equilibrium (zero balance).

WHO/FAO/UNU (2007) re-evaluated their recommendations from 1985. Based on a meta-analysis of nitrogen balance studies in humans by Rand et al. (2003), which involved studies stratified for a number of subpopulations, settings in different climates, sex, age and protein source, a population average requirement of 0.66 g/kg body weight per day resulted as the best estimate. The “safe level of intake” was identified as the 97.5<sup>th</sup> percentile of the population distribution of requirement, which was equivalent to 0.83 g/kg body weight per day of high quality protein (WHO/FAO/UNU, 2007). The French recommendations (AFSSA, 2007) established a PRI of 0.83 g/kg body weight per day for adults based on the WHO/FAO/UNU (2007) report. The German speaking countries (D-A-CH, 2008) used the average requirement for high quality protein of 0.6 g/kg body weight per day (estimated by FAO/WHO (1985)), included an allowance for individual variability (value increased to 0.75 g/kg body weight per day), and took account of frequently reduced protein digestibility in mixed diets to establish a recommended intake of 0.8 g/kg body weight per day for adults.



**Table 3:** Overview of Dietary Reference Values and recommendations for protein for adults

	FAO/ WHO/UNU (1985)	DoH (1991)	SCF (1993)	Health Council of the Netherlands (2001)	NNR (2004)	IoM (2005)	WHO/ FAO/UNU (2007)	AFSSA (2007)	D-A-CH (2008)
<b>AR - Adults</b> (g/kg bw x d <sup>-1</sup> )	0.60	0.60	0.60	0.60	-	0.66	0.66	0.66	0.60
<b>PRI - Adults</b> (g/kg bw x d <sup>-1</sup> )	0.75 <sup>1</sup>	0.75	0.75	0.80	-	0.80 <sup>2</sup>	0.83 <sup>1</sup>	0.83	0.80
<b>PRI - Adult Males</b> (g/d)	-	56	56	59	-	56	-	-	59
<b>PRI - Adult Females</b> (g/d)	-	45	47	50	-	46	-	-	47
<b>Recommended intake range – Adults</b> (E%)	-	-	-	-	10-20	10-35 <sup>3</sup>	-	-	-

<sup>1</sup>Safe level of intake; <sup>2</sup> RDA; <sup>3</sup>Acceptable Macronutrient Distribution Range

#### 4.1.1. Older adults

In 1985, FAO/WHO/UNU recommended an intake of 0.75 g/kg body weight per day of good quality protein for adults, and the same recommendation was made for adults over the age of 60 years because, although efficiency of protein utilisation is assumed to be lower in older adults, the smaller amount of lean body mass per kg body weight will result in a higher figure per unit lean body mass than in younger adults (FAO/WHO/UNU, 1985).

The recommended intake for adults in the Netherlands (Health Council of the Netherlands, 2001) is 0.8 g/kg body weight per day and no additional allowance was considered necessary for adults aged >70 years. The US Institute of Medicine (IoM, 2005) recommended 0.8 g/kg body weight per day of good quality protein for adults. For adults aged 51-70 years and >70 years, no additional protein allowance beyond that of younger adults was considered necessary since no significant effect of age on protein requirement expressed per kg body weight was observed in the analysis by Rand et al. (2003), recognising that lean body mass as % body weight and protein content of the body both decrease with age.

The WHO/FAO/UNU (2007) report also concluded that the available data did not provide convincing evidence that the protein requirement of elderly people (per kg body weight, no age range given) differs from the protein requirement of younger adults. The conclusion is partly supported by data on nitrogen balance (Campbell et al., 2008) which showed that the mean protein requirement was not different between younger (21–46 years) and older (63–81 years) healthy adults: 0.61 (SD 0.14) compared with 0.58 (SD 0.12) g protein/kg body weight per day. However, the low energy requirement of sedentary elderly people means that the protein to energy ratio of their requirement is higher than for younger age groups. Thus, unless the elderly people are physically active they may need a more protein-dense diet.

In France, an intake of 1.0 g/kg body weight per day has been recommended for people ≥75 years based on considerations about protein metabolism regulation in the elderly (AFSSA, 2007). The German speaking countries (D-A-CH, 2008) recommended an intake of 0.8 g protein/kg body weight per day for adults and the same recommendation was made for adults aged 65 years and older since it was considered that the available evidence was insufficient to prove a higher requirement for the elderly.

#### 4.2. Dietary Reference Values and recommendations for protein for infants and children

Table 4 lists reference intakes set by various organisations for infants and children.

In its report, FAO/WHO/UNU (1985) calculated the protein requirements of children from six months onwards by a modified factorial method. Maintenance requirements were interpolated between the values from nitrogen balance studies for children aged one year and those for young adults aged 20 years. A CV of 12.5 % was used to allow for individual variability. The growth component of the protein requirement was set at 50 % above that based on the theoretical daily amount of nitrogen laid down, corrected for an

efficiency of dietary protein utilisation of 70 %. The average requirement was then estimated as the sum of the maintenance and growth requirements. The “safe level of intake” was estimated based on the average requirement plus two SD corresponding to a CV of 12-16 %.

In its re-evaluation, WHO/FAO/UNU (2007) calculated a maintenance value of 0.66 g protein/kg body weight per day for children and infants from 6 months to 18 years. The maintenance level was derived from a regression analysis of nitrogen balance studies on children from 6 months to 12 years. Protein deposition needs were calculated from combined data of two studies and assuming an efficiency of utilisation for growth of 58 %. The average requirement was then estimated as the sum of the maintenance and growth requirements. The “safe level of intake” was estimated based on the average level plus 1.96 SD. Requirements fall rapidly in the first two years of life (safe level at six months of age: 1.31 g/kg body weight per day; at two years of age: 0.97 g/kg body weight per day). Thereafter, the decrease towards the adult level is very slow (WHO/FAO/UNU, 2007).

Dewey et al. (1996) reviewed the approach by FAO/WHO/UNU (1985) and suggested revised estimates for protein requirements for infants and children. The German speaking countries (D-A-CH, 2008) followed the proposal of Dewey et al. (1996). For infants aged from 6 to under 12 months the maintenance requirement was estimated at 0.56 g/kg body weight per day from nitrogen balance studies. Age-dependent additions of between 35 and 31 % for the increase in body protein were made to take into account inter-individual variability of maintenance and growth requirements (Dewey et al., 1996). A recommended intake of 1.1 g/kg body weight per day (10 g/d) of high quality protein was established from 6 to under 12 months. Recommended intakes were established for children aged 1 to under 4 years (1.0 g/kg body weight per day) and 4 to under 15 years, and for boys aged 15 to under 19 years (0.9 g/kg body weight per day) and girls aged 15 to under 19 years (0.8 g/kg body weight per day). The maintenance requirement was estimated at 0.63 g/kg body weight per day (Dewey et al., 1996) and total requirement, allowing for the decreasing requirement for growth with age, was estimated to range from 0.63-0.7 g/kg body weight per day. An additional 30 % allowance was made to account for inter-individual variability in protein utilisation and digestibility.

The Nordic Nutrition recommendations (NNR, 2004) also followed the approach of Dewey et al. (1996) to establish recommended intakes of 1.1 and 1.0 g/kg body weight per day for infants aged 6-11 months and children aged 1-1.9 years, respectively. For children aged 2-17 years a recommended intake of 0.9 g/kg body weight per day was established, in agreement with the values in other recommendations (D-A-CH, 2008; Health Council of the Netherlands, 2001; IoM, 2005). The French recommendations (AFSSA, 2007) also followed the approach of Dewey et al. (1996).

The Health Council of the Netherlands (2001) used a factorial method derived from nitrogen balance experiments to estimate the protein requirements of infants over 6 months, children and adolescents. For infants aged 6-11 months a recommended intake of 1.2 g/kg body weight per day (10 g/d) of high quality protein was established. This was based on an average requirement for maintenance and growth of 0.9 g/kg body weight per day, with a CV of 15 % to allow for individual variability, and assuming an efficiency of dietary protein utilisation of 70 %. Recommended intakes were established for children aged 1 to 13 years (0.9 g/kg body weight per day) and 14 to 18 years (0.8 g/kg body weight per day) on the same basis but using an average requirement for maintenance and growth of 0.8 g/kg body weight per day for children aged 1 to 3 years and 0.7 g/kg body weight per day for children aged 4 to 18 years (Health Council of the Netherlands, 2001).

**Table 4:** Overview of Dietary Reference Values for protein for children

	FAO/ WHO/ UNU (1985) <sup>1</sup>	SCF (1993) <sup>1</sup>	Health Council of the Netherlands (2001)	NNR (2004)	IoM (2005) <sup>2</sup>	WHO/ FAO/ UNU (2007) <sup>1</sup>	AFSSA (2007)	D-A-CH (2008)
<b>Age</b>	6-9 months	7-9 months	6-11 months	6-11 months	7-12 months	6 months	6-12 months	6-<12 months
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	1.65 (m + f)	1.65 (m + f)	1.2 (m + f)	1.1 (m + f)	1.2 (m + f)	1.31 (m + f)	1.1 (m + f)	1.1 (m + f)
<b>Age</b>	9-12 months	10-12 months	1-13 y	1-1.9 y	1-3 y	1 y	12-24 months	1- <4 y
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	1.50 (m + f)	1.48 (m + f)	0.9 (m + f)	1.0 (m + f)	1.05 (m + f)	1.14 (m + f)	1.0 (m+f)	1.0 (m + f)
<b>Age</b>	1-2 y	1-1.5 y				1.5 y		
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	1.20 (m + f)	1.26 (m + f)				1.03 (m + f)		
<b>Age</b>	2-3 y	2-3 y		2-17 y		2 y	24-36 months	
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	1.15 (m + f)	1.13 (m + f)		0.9 (m + f)		0.97 (m + f)	0.9 (m+f)	
<b>Age</b>	3-5 y	4-5 y				3 y	3-10 y	
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	1.10 (m + f)	1.06 (m + f)				0.90 (m + f)	0.9 (m+f)	
<b>Age</b>	5-12 y	6-9 y			4-13 y	4-6 y	10-12 y (m), 10-11 y (f)	4-<15 y
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	1.0 (m + f)	1.01 (m + f)			0.95 (m + f)	0.87 (m + f)	0.85 (m), 0.9 (f)	0.9 (m + f)
<b>Age</b>	12-14 y	12 y				7-10 y	12-13 y (m), 11-14 y (f)	
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	1.0 (m) 0.95 (f)	1.0 (m) 0.96 (f)				0.92 (m + f)	0.9 (m), 0.85 (f)	
<b>Age</b>	14-16 y	14 y	14-18 y		14-18 y	11-14 y	13-17 y (m), 14-16 y (f)	
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	0.95 (m) 0.9 (f)	0.96 (m) 0.90 (f)	0.8 (m + f)		0.85 (m + f)	0.90 (m) 0.89 (f)	0.85 (m), 0.8 (f)	
<b>Age</b>	16-18 y	16 y				15-18 y	17-18 y (m), 16-18 y (f)	15-<19 y
<b>PRI</b> (g/kg bw x d <sup>-1</sup> )	0.9 (m) 0.8 (f)	0.90 (m) 0.83 (f)				0.87 (m) 0.84 (f)	0.8 (m + f)	0.9 (m) 0.8 (f)

<sup>1</sup> Safe level of intake; <sup>2</sup> RDA

The US Institute of Medicine (IoM, 2005) recommended intakes ranging from 1.2 g/kg body weight per day of high quality protein for infants aged 6-12 months to 0.85 g/kg body weight per day for 14 to 18 year-old boys and girls based on estimates of requirements for maintenance, with additions for growth. Maintenance requirements were estimated from short-term nitrogen balance studies as 110 mg N/kg body weight per day for older infants and children aged 7 months through 13 years, and as 105 mg N/kg body weight per day (estimated from short-term nitrogen balance studies in adults and based on a meta-analysis by Rand et al. (2003)) for ages 14 through 18 years. Growth requirements were estimated in infants and children from estimated rates of nitrogen accretion calculated from rates of weight gain and from estimates of the nitrogen content of tissues. The efficiency of dietary protein utilisation was assumed to be 58 % for ages 7 months through 13 years and 47 % for ages 14 through 18 years, estimated from the slopes of the nitrogen balance data. The EAR was thus estimated as 1.0 g/kg body weight per day for infants aged 7-12 months, 0.87 and 0.76 for boys and girls aged 1-3 and 4-13 years, respectively, and 0.73 and 0.71 g/kg body weight per day for boys and girls aged 14-18 years, respectively. A CV of 12 % for maintenance and of 43 % for growth was used in the calculation of the RDA to allow for individual variability (IoM, 2005).

#### 4.3. Dietary Reference Values and recommendations for protein during pregnancy

FAO/WHO/UNU (1985) recommended an average additional intake of 6 g/d throughout pregnancy based on derived additional levels of protein intake of 1.2 g/d, 6.1 g/d and 10.7 g/d for the first, second and third trimester, respectively. This was based on the calculated average increment of 925 g protein during pregnancy, plus 30 % (2 SD of birth weight), adjusted for the efficiency with which dietary protein is converted to foetal, placental and maternal tissues (estimated as 70 %) (FAO/WHO/UNU, 1985). WHO/FAO/UNU (2007) revised this value and recommended 1, 9 and 31 g of additional protein/d in the first, second and third trimester, respectively, as “safe intake levels”. Based on a theoretical model (Hyttén and Chamberlain, 1990), the total deposition of protein in the foetus and maternal tissue has been estimated as 925 g (assuming a 12.5 kg gestational weight gain), of which 42 % is deposited in the foetus, 17 % in the uterus, 14 % in the blood, 10 % in the placenta and 8 % in the breasts. Protein deposition has also been estimated indirectly from measurements of total body potassium accretion, measured by whole-body counting in a number of studies with pregnant women (Butte et al., 2003; Forsum et al., 1988; King et al., 1973; Pipe et al., 1979). From these studies, mean protein deposition during pregnancy was estimated as 686 g (WHO/FAO/UNU, 2007). Based on the study by Butte et al. (2003), protein deposition per trimester was then calculated for well-nourished women achieving a gestational weight gain of 13.8 kg (the mid-point of the recommended weight gain range for women with normal pre-pregnancy weight) (IoM, 1990). The efficiency of dietary protein utilisation was taken to be 42 % in pregnant women (in comparison to 47 % in non-pregnant adults) (WHO/FAO/UNU, 2007).

In Europe, UK COMA (DoH, 1991) accepted the value proposed by FAO/WHO/UNU (1985). SCF (1993) used the approach of FAO/WHO/UNU (1985) but recommended an additional intake of 10 g/d throughout pregnancy because of uncertainty about changes in protein metabolism associated with pregnancy (SCF, 1993). The Dutch (Health Council of the Netherlands, 2001) recommended an additional intake of 0.1 g/kg body weight per day throughout pregnancy. AFSSA (2007) followed the approach of FAO/WHO/UNU (1985) and recommended an intake between about 0.82 and 1 g/kg body weight per day for a woman of 60 kg (calculated from 50, 55 and 60 g/d for each trimester of pregnancy). The German speaking countries (D-A-CH, 2008) recommended an additional intake of 10 g/d (for the second and third trimesters).

The US Institute of Medicine (IoM, 2005) set the EAR at 21 g/d above the average protein requirement of non-pregnant women, averaging the overall protein needs over the last two trimesters of pregnancy. It recommended an additional intake of 25 g/d (RDA for the second and third trimesters), assuming a CV of 12 % and rounding to the nearest 5 g/d. The EAR for additional protein needs was based upon an estimated average protein deposition of 12.6 g/d over the second and third trimesters (calculated from potassium retention studies for accretion of 5.4 g protein/d, and assuming an efficiency of dietary protein utilisation of 43 %), plus an additional 8.4 g/d for maintenance of the increased body tissue.

#### 4.4. Dietary Reference Values and recommendations for protein during lactation

FAO/WHO/UNU (1985) recommended an additional intake of 16 g/d of high quality protein during the first six months of lactation, 12 g/d during the second six months, and 11 g/d thereafter. This is based on the average protein content of human milk, an efficiency factor of 70 % to adjust for the conversion of dietary protein to milk protein, and a CV of milk volume of 12.5 % (FAO/WHO/UNU, 1985). WHO/FAO/UNU (2007) revised this value and recommended an additional protein intake of 19 g/d in the first six months of lactation and 12.5 g/d after six months. This is based on the increased nitrogen needs of lactating women in order to synthesise milk proteins, with the assumption that the efficiency of milk protein production is the same as the efficiency of protein synthesis in non-lactating adults, i.e. 47 %. Therefore, the additional “safe intake” of dietary protein was calculated using an amount of dietary protein equal to milk protein, divided by an efficiency of 47 % and adding 1.96 SD corresponding to a CV of 12 % (WHO/FAO/UNU, 2007).

In Europe, UK COMA (DoH, 1991) recommended an additional intake of 11 g/d for the first six months and an additional intake of 8 g/d thereafter. The approach used was similar to that of FAO/WHO/UNU (1985) except that the values for human milk protein content used were lower because of correction for the amount (up to 25 %) of non-protein nitrogen present. SCF (1993) accepted the values proposed by FAO/WHO/UNU (1985), i.e. an additional intake of 16 g/d of high quality protein during the first six months of lactation and

12 g/d during the second six months. The Netherlands (Health Council of the Netherlands, 2001) recommended an additional intake of 0.2 g/kg body weight per day during lactation to allow for the additional protein loss of about 7 g/d in human milk. AFSSA (2007) considered the quantity of protein and non-protein nitrogen excreted in milk and its change during lactation, and recommended an additional intake of 16 g/d for the first six months, resulting in a recommended intake of about 1.1 g/kg body weight per day for a woman of 60 kg. The German speaking countries (D-A-CH, 2008) recommended an additional intake of 15 g/d during lactation based on a mean protein loss of 7-9 g/d in human milk, assuming an efficiency of utilisation of 70 % and adding 2 SD to account for inter-individual variability.

The US Institute of Medicine (IoM, 2005) calculated the EAR of additional protein during lactation (21 g/d) from the average protein equivalent of milk nitrogen output and an assumed efficiency of utilisation of 47 %. Adding 2 SD (24 %) to account for inter-individual variability yielded an RDA of +25 g/d, or a recommended protein intake of 1.3 g/kg body weight per day during lactation.

#### 4.5. Requirements for indispensable amino acids

Different approaches have been used to determine indispensable amino acid requirements. These requirements were first determined in adults using a nitrogen balance approach (Rose, 1957). The values obtained by this approach are usually considered to underestimate the requirements (Rand and Young, 1999; WHO/FAO/UNU, 2007; Young and Marchini, 1990). More recent data in adults have been obtained using amino acids labelled with stable isotopes, and are based on the measurement of amino acid oxidation as a function of intake (Bos et al., 2002). This includes the indicator amino acid balance method (Young and Borgonha, 2000), the indicator amino acid oxidation method (Elango et al., 2008a, 2008b; Pencharz and Ball, 2003), the 24 h-indicator amino acid oxidation method (Kurpad et al., 2001) and the protein post-prandial retention method (Bos et al., 2005; Millward et al., 2000).

The rationale for deriving DRVs for each indispensable amino acid remains questionable since as a rule amino acids are not provided as individual nutrients in the diet but in the form of protein. Moreover, the values obtained for indispensable amino acid requirement are not yet sufficiently precise and require further investigation (AFSSA, 2007; WHO/FAO/UNU, 2007). Only the US introduced specific RDAs for indispensable amino acids, derived from the average values of requirements deduced from amino acid oxidation methods and adding 2 CV (of 12 %) (IoM, 2005).

Average indispensable amino acid requirements are used to calculate the indispensable amino acid reference pattern, which is used in the assessment of protein quality according to the chemical score approach and the PD-CAAS. The mean values for indispensable amino acid requirements were provided in the WHO/FAO/UNU (2007) report (Table 5).

**Table 5:** Mean requirements for indispensable amino acids in adults (WHO/FAO/UNU, 2007)

	mg/kg bw x d <sup>-1</sup>		mg/kg bw x d <sup>-1</sup>
Histidine	10	Phenylalanine+tyrosine	25
Isoleucine	20	Threonine	15
Leucine	39	Tryptophan	4
Lysine	30	Valine	26
Methionine+cysteine	15 <sup>1</sup>	Total	184
<i>methionine</i>	10.4		
<i>cysteine</i>	4.1		

<sup>1</sup> resulting from rounding

The amino acid requirements of infants and children have been derived using a factorial method, based on the estimated protein requirements for maintenance and growth (Dewey et al., 1996; WHO/FAO/UNU, 2007) (Table 6). It is assumed that the required amino acid pattern for maintenance is the same as that for adults, and that the amino acid pattern required for growth is given by the amino acid composition of whole-body tissue protein (Davis et al., 1993; Dewey et al., 1996; Widdowson et al., 1979).

**Table 6:** Mean requirements for indispensable amino acids in infants, children and adolescents (WHO/FAO/UNU, 2007)

	Mean amino acid requirement at different ages (mg/kg bw x d <sup>-1</sup> )				
	0.5 years	1-2 years	3-10 years	11-14 years	15-18 years
Histidine	22	15	12	12	11
Isoleucine	36	27	23	22	21
Leucine	73	54	44	44	42
Lysine	64	45	35	35	33
Methionine+cysteine	31	22	18	17	16
Phenylalanine+tyrosine	59	40	30	30	28
Threonine	34	23	18	18	17
Tryptophan	9.5	6.4	4.8	4.8	4.5
Valine	49	36	29	29	28

## 5. Criteria (endpoints) on which to base Dietary Reference Values (DRVs)

Current DRVs for protein are based on protein homeostasis measured as nitrogen balance. DRVs also take into account protein quality, which is related to the capacity of a protein source to meet both the requirement for nitrogen and the requirement for indispensable amino acids as limiting precursors for body protein synthesis. Other criteria taking into account the functional and health consequences of protein intake may also be considered to derive DRVs for protein.

### 5.1. Protein intake and protein and nitrogen homeostasis

#### 5.1.1. Methods for the determination of protein requirement

##### 5.1.1.1. Nitrogen balance

Nitrogen balance is the classical approach for the determination of protein requirement and in initial studies of indispensable amino acid requirements (FAO/WHO/UNU, 1985). Nitrogen balance is the difference between nitrogen intake and the amount lost in urine, faeces, via the skin and via other miscellaneous ways such as nasal secretions, menstrual losses, or seminal fluid (IoM, 2005). In healthy adults at energy balance the protein requirement (maintenance requirement) is defined as that amount of dietary protein which is sufficient to achieve zero nitrogen balance. It is assumed that nitrogen balance will be negative when protein intakes are inadequate. In infants and children, nitrogen balance has to be positive to allow for growth. While there are substantial practical limitations of the method mainly related to the accuracy of the measurements and the interpretation of the results (WHO/FAO/UNU, 2007), nitrogen balance remains the method of choice for determining protein requirement in adults (Rand et al., 2003).

##### 5.1.1.2. Indicator amino acid oxidation method

As an alternative method the indicator amino acid oxidation method has been discussed (Elango et al., 2008a), but very few data are available using this indirect method for the determination of protein requirements. The values provided for the protein requirement of seven school-age children (Elango et al., 2011), of eight healthy men (Humayun et al., 2007) and of 20 young women (Tian et al., 2011) are (considerably) higher than the requirements derived from nitrogen balance measurements and there is no explanation for the origin of these differences.

##### 5.1.1.3. The factorial method

The factorial method is based on the assessment of the extent to which dietary protein nitrogen is absorbed and retained by the organism, and is able to balance daily nitrogen losses and allow additional protein deposition in newly formed tissue for growth, and in specific physiological conditions such as pregnancy or lactation. Obligatory nitrogen losses are estimated from subjects fed a diet that meets energy needs but is essentially protein-free, or more reliably is derived from the y-intercept of the slope of the regression line

relating nitrogen intake to nitrogen retention. The requirement for dietary protein is considered to be the amount needed to replace nitrogen losses and to allow additional protein deposition, after adjustment for the efficiency of dietary protein utilisation (see Section 2.4.) and the quality of the dietary protein. The factorial method is used to calculate protein requirements in physiological conditions such as growth, pregnancy or lactation. A critical factor is the value used for efficiency of dietary protein utilisation (Table 7).

**Table 7:** Previously used values for efficiency of dietary protein utilisation in different population groups and values used by EFSA in this Scientific Opinion

Population group	Previously used values (%)	Values used by EFSA (%)
Adults	70 <sup>(1)</sup> , 47 <sup>(2, 3)</sup>	47
Infants and children (for growth)	70 <sup>(1)</sup> , 58 <sup>(2)</sup> , 58/47 <sup>(3)</sup>	58
Pregnant women (for protein deposition)	70 <sup>(1)</sup> , 42 <sup>(2)</sup> , 43 <sup>(3)</sup>	47
Lactating women	70 <sup>(1)</sup> , 47 <sup>(2, 3)</sup>	47

<sup>1</sup>FAO/WHO/UNU (1985); <sup>2</sup>WHO/FAO/UNU (2007); <sup>3</sup>IoM (2005)

In healthy adults, the mean post-prandial protein efficiency in controlled optimal conditions is considered to be 70 %, and this value was first used as a reference for the different population groups including infants and women during pregnancy and lactation (FAO/WHO/UNU, 1985). However, the NPU value can be modified by various factors including the food matrix, the diet and certain physiological conditions. More recently, a value of 47 % was derived from nitrogen balance studies in healthy adults under maintenance conditions (Rand et al., 2003). For children, WHO/FAO/UNU (2007) estimated the NPU for protein deposition with growth to be 58 % from 6 months to 18 years, whereas IoM (2005) estimated it to be 58 % from 7 months to 13 years and 47 % from 14 to 18 years. During lactation the NPU was estimated to be 47 % and not to be different from that in non-lactating healthy adults (WHO/FAO/UNU, 2007). For ten pregnant adolescents, King et al. (1973) derived a relatively low value of nitrogen retention of 30 %. From different nitrogen balance studies, Calloway (1974) calculated a nitrogen retention of 25-30 %. However, in healthy pregnant women, nitrogen efficiency was found to be increased in comparison with non-pregnant women receiving the same nitrogen intake above the requirement (Mojtahedi et al., 2002). From the study by King et al. (1973), IoM (2005) recalculated an NPU value of 43 % based on those six adolescents who demonstrated a positive efficiency at multiple levels of protein intake (IoM, 2005) and WHO/FAO/UNU (2007) recalculated the efficiency of utilisation of dietary protein to be 42 % after omitting the two subjects who gave negative gradients. Eight Indian pregnant women utilised 47 % of the dietary nitrogen when 60-118 g/d of mixed protein was consumed. The nitrogen intake of the Indian women was unrelated to nitrogen retention unless intakes above 0.45 g N/kg body weight per day were omitted (Jayalakshmi et al., 1959). A similar range of values has been observed in pregnant sows (Dunn and Speer, 1991; Jones and Maxwell, 1982; King and Brown, 1993; Renteria-Flores et al., 2008; Theil et al., 2002).

The Panel considers that for healthy adults a protein efficiency value of 47 % is reasonable since it is the value derived from the nitrogen balance studies used to define nitrogen requirement in adults. There is no convincing scientific evidence that protein efficiency for maintenance of body protein and for protein deposition is lower during pregnancy or lactation. As a consequence, the same value can be considered as that determined for healthy adults (47 %). For infants and children, a value of 58 % for growth is justified because of an increased efficiency of dietary protein utilisation for growth.

#### 5.1.1.4. Protein quality and reference pattern for indispensable amino acids

The protein requirement is dependent on the dietary protein quality, which is mainly determined by the pattern of indispensable amino acids in the protein. The reference pattern of amino acids for infants <0.5 years is the amino acid pattern of human milk. The reference pattern of amino acids (mg/g protein) for the assessment of protein quality for adults is derived from proposed data on the requirement for individual indispensable amino acids (WHO/FAO/UNU, 2007) by dividing the requirement (mg amino acid/kg body weight per day) by the average requirement for protein (g/kg body weight per day). Age-specific scoring patterns for dietary proteins can be derived by dividing the requirement of each indispensable amino acid by the protein requirement of the selected age group (WHO/FAO/UNU, 2007) (Table 8).

In practice, three reference patterns are used: the amino acid pattern of human milk for infants <0.5 years, the 3-10 years reference pattern for infants and children, and the adult reference pattern.

**Table 8:** Scoring pattern (indispensable amino acid reference profiles) for infants, children, adolescents and adults (WHO/FAO/UNU, 2007)

	Infants, children, adolescents (mg/g protein)					Adults (mg/g protein)
	0.5 years	1-2 years	3-10 years	11-14 years	15-18 years	
Histidine	20	18	16	16	16	15
Isoleucine	32	31	31	30	30	30
Leucine	66	63	61	60	60	59
Lysine	57	52	48	48	47	45
Methionine+cysteine	28	26	24	23	23	22
Phenylalanine+tyrosine	52	46	41	41	40	30
Threonine	31	27	25	25	24	23
Tryptophan	8.5	7.4	6.6	6.5	6.3	6
Valine	43	42	40	40	40	39

### 5.1.2. Protein requirement of adults

In a meta-analysis by Rand et al. (2003), available nitrogen balance data as a function of nitrogen intake among healthy persons were analysed. Data obtained from 235 individuals, each studied at  $\geq 3$  test protein intakes, were gathered from 19 primary and secondary studies and used for estimating the average requirement. Subjects were classified by sex and age ( $\leq 55$  (n=221) and  $>55$  years of age (n=14)), diets by the main source of protein (animal ( $>90\%$  of total protein intake from animal sources), vegetable ( $>90\%$  of total protein intake from vegetable sources) or mixed), and climate was classified as temperate or tropical. As the distribution of individual requirements was significantly skewed and kurtotic, the mean was not a robust estimate of the centre of the population and the median was taken as the average requirement.

The Panel notes that the study by Rand et al. (2003) concluded that the best estimate of average requirement for 235 healthy adults from 19 studies was 105 mg N/kg body weight per day (0.66 g high quality protein/kg body weight per day). The 97.5<sup>th</sup> percentile of the population distribution of the requirement was estimated from the log median plus 1.96 times the SD of 0.12 and found to be 133 mg N/kg body weight per day (0.83 g high quality protein/kg body weight per day). Thus, 0.83 g protein/kg body weight per day can be expected to meet the requirements of most (97.5 %) of the healthy adult population. This value can be considered to fulfil the function of a PRI although derived differently. The data did not provide sufficient statistical power to establish different requirements for different adult groups based on age, sex, or dietary protein source (animal or vegetable proteins) (Rand et al., 2003). The Panel notes that by considering only the primary studies based on 32 data points the requirement would be 101.5 mg/kg body weight per day, but that the statistical power is greatly reduced and that this value is not significantly different to the value of 105 mg N/kg body weight per day.

The Panel considers that the value of 0.66 g/kg body weight per day can be accepted as the AR and the value of 0.83 g/kg body weight per day as the PRI derived for proteins with a PD-CAAS value of 1.0. This value can be applied to usual mixed diets in Europe which are unlikely to be limiting in their content of indispensable amino acids (WHO/FAO/UNU, 2007).

#### 5.1.2.1. Older adults

Few and contradictory data are available on the protein requirement of older adults compared to young and middle-aged adults. The hypothesis that the PRI for older adults may be greater than that for younger adults (0.83 g/kg body weight per day) (Gaffney-Stomberg et al., 2009; Thalacker-Mercer et al., 2010; Wolfe et al., 2008) was particularly discussed on the basis of an assumed, although not significantly lower efficiency of protein utilisation in the elderly (AFSSA, 2007; Rand et al., 2003).

Several studies concluded that the PRI for protein (0.83 g protein/kg body weight per day) is also adequate for older adults to reach nitrogen balance (Campbell et al., 2008; Pannemans et al., 1995a; Pannemans et al.,



1995b; Rand et al., 2003). This conclusion was previously adopted by WHO/FAO/UNU (2007), mainly based on the comprehensive meta-analysis of Rand et al. (2003).

Following the meta-analysis by Rand et al. (2003), one more recent study did not show differences between younger (21-46 years) and older (63-81 years) subjects after short-term assessment of nitrogen balance (Campbell et al., 2008). In contrast, some studies observed that older adults consuming protein at the level of the PRI (0.8 g protein/kg body weight per day) were in negative nitrogen balance and that a protein intake around 0.9-1.0 g/kg body weight per day seems to be more adequate (Bunker et al., 1987; Campbell et al., 2001; Kortebein et al., 2007; Pannemans et al., 1998; Pannemans et al., 1997). A negative nitrogen balance was observed in six elderly females (69 ± 5 years) consuming a diet providing 0.8 g protein/kg body weight per day for two weeks (Pannemans et al., 1997), and the same level of intake was associated with a decrease in the mid-thigh muscle area in ten men and women (aged 55-77 years) during 14 weeks, although whole-body leucine metabolism and body composition were not affected (Campbell et al., 2001). In the observational Health, Aging, and Body Composition (Health ABC) Study (Houston et al., 2008), protein intake was negatively associated with 3-year changes in lean mass (p=0.004) and appendicular lean mass<sup>5</sup> (p=0.001); however, losses in lean mass in subjects ingesting 0.8 g protein/kg body weight per day did not significantly differ from those in the highest quintile of protein intake (1.1 g/kg body weight per day) and no differences in lean mass losses between the five quintiles of protein intake were observed in weight stable subjects.

The difficulty in assessing protein requirements in older subjects arises from the different factors that can influence protein efficiency, including protein quality (animal versus plant protein), distribution of protein intake over the day, and physical activity. Some have suggested that protein quality and the protein composition of the diet may modify the level of protein required to reach nitrogen balance in older adults (Pannemans et al., 1998). In addition, for the same level of protein consumed daily the mode of ingestion (even distribution between meals vs. bolus) may influence the efficiency of the anabolic protein response (Cuthbertson et al., 2005; Katsanos et al., 2005; Paddon-Jones et al., 2004; Paddon-Jones and Rasmussen, 2009; Symons et al., 2007). Lastly, exercise has been demonstrated to improve nitrogen balance and to decrease losses of lean mass in older adults, and the studies concluded that an increase in dietary protein alone above the usual recommended intake does not change body composition or improve lean body mass unless accompanied by physical training programmes (Campbell and Leidy, 2007; Iglay et al., 2009; Paddon-Jones and Rasmussen, 2009).

The Panel concludes that the available data are insufficient to specifically determine the protein requirement in older adults and that at least the same level of protein intake as for young adults is required for older adults. As sedentary older adults have a lower energy requirement the protein to energy ratio of this subgroup is higher than for younger adults.

### 5.1.3. Protein requirement of infants and children

The protein requirement of infants and children includes two components, i.e. maintenance requirement and growth requirement. This protein requirement can be defined as the minimum intake that will allow a positive nitrogen equilibrium to allow for growth in normally growing subjects who have an appropriate body composition, are in energy balance and moderately physically active (WHO/FAO/UNU, 2007).

In the report by WHO/FAO/UNU (2007), estimates of the protein requirement from 6 months to 18 years were derived factorially as the sum of requirements for maintenance and growth corrected for efficiency of dietary protein utilisation. An average maintenance requirement of 0.66 g protein/kg body weight per day was applied to infants and children from 6 months to 18 years (Tables 9 and 10). Regression analysis of nitrogen balance studies on children from 6 months to 12 years resulted in a maintenance level of 110 mg N/kg body weight per day. Because this value was close to the adult maintenance value of 105 mg N/kg body weight per day and it could not be determined with certainty that maintenance values for infants and children differ from those for adults, the latter value was selected as the maintenance value for ages from 6 months onwards. Average daily needs for dietary protein for growth were estimated from average daily

<sup>5</sup> Lean mass of the arms and legs

rates of protein deposition, calculated from studies on whole-body potassium deposition, and adjusted by an efficiency of utilisation of dietary protein of 58 %. The average requirement for protein was adjusted according to the expected variability of maintenance and growth to give a value equivalent to the 97.5<sup>th</sup> percentile of the distribution as a measure of the PRI, based on the average requirement plus 1.96 SD.

The Panel agrees with this analysis of the data.

**Table 9:** Average protein requirement of infants from 6 months onwards and children up to 10 years of age derived by WHO/FAO/UNU (2007)

Age (years)	0.5	1	1.5	2	3	4	5	6	7	8	9	10
Maintenance requirement (g/kg bw x d <sup>-1</sup> )	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66
Growth requirement (g/kg bw x d <sup>-1</sup> )	0.46	0.29	0.19	0.13	0.07	0.03	0.03	0.06	0.08	0.09	0.09	0.09
Average requirement (g/kg bw x d <sup>-1</sup> )	1.12	0.95	0.85	0.79	0.73	0.69	0.69	0.72	0.74	0.75	0.75	0.75

**Table 10:** Average protein requirement of adolescents derived by WHO/FAO/UNU (2007)

Age (years)	11	12	13	14	15	16	17	18
Maintenance requirement (g/kg bw x d <sup>-1</sup> )	0.66	0.66	0.66	0.66	0.66	0.66	0.66	0.66
Growth requirement (g/kg bw x d <sup>-1</sup> )	0.09 (m) 0.07 (f)	0.08 (m) 0.06 (f)	0.07 (m) 0.05 (f)	0.06 (m) 0.04 (f)	0.06 (m) 0.03 (f)	0.05 (m) 0.02 (f)	0.04 (m) 0.01 (f)	0.03 (m) 0.00 (f)
Average requirement (g/kg bw x d <sup>-1</sup> )	0.75 (m) 0.73 (f)	0.74 (m) 0.72 (f)	0.73 (m) 0.71 (f)	0.72 (m) 0.70 (f)	0.72 (m) 0.69 (f)	0.71 (m) 0.68 (f)	0.70 (m) 0.67 (f)	0.69 (m) 0.66 (f)

#### 5.1.4. Protein requirement during pregnancy

The protein requirement during pregnancy has to take into account the requirements for deposition of new protein and for the maintenance of the weight gained, in addition to the requirement in the non-pregnant state. It can be determined by using either the nitrogen balance approach or the factorial method.

In the nitrogen balance approach, nitrogen requirement is derived from nitrogen balance studies. This requires balance measurements in women at different levels of protein intake in order to determine the maximal nitrogen deposition potential and to derive nitrogen requirement from this maximal nitrogen deposition (Calloway, 1974). However, it appears from the available studies that there is a linear increase in apparent nitrogen deposition with increasing protein intake in pregnant women. The linear relationship between nitrogen intake and deposition towards the end of pregnancy is statistically significant<sup>6</sup> (Calloway, 1974; Jayalakshmi et al., 1959; Johnstone et al., 1981; King et al., 1973).

According to the slope of these equations, the average nitrogen efficiency is very low, i.e. between 21 and 47 %. The linear nature of the relation between nitrogen intake and retention does not permit the determination of a maximal nitrogen deposition potential, or to derive a nitrogen requirement related to this maximal nitrogen deposition. The cause for this linear relationship remains unclear. This linear relation and the low level of nitrogen efficiency derived from the slopes indicate uncertainties and errors in the measurement of nitrogen balance, and implicate important limitations for the use of this approach to determine the nitrogen requirement in pregnant women.

<sup>6</sup> In the study by Jayalakshmi et al. (1959), a linear relationship was only obtained after exclusion of four values indicating nitrogen retention for intakes >0.45 g N/kg body weight per day.

The alternative approach is the factorial method used by IoM (2005) and WHO/FAO/UNU (2007). The maintenance costs were based upon the mid-trimester increase in maternal body weight, and the maintenance value of 0.66 g/kg body weight per day was derived from the average requirement in healthy adults, assuming a CV of 12 %. Protein deposition in the foetus and the maternal tissue has been estimated indirectly from measurements of total body potassium accretion. However, studies show that protein is not deposited equally throughout pregnancy. For well-nourished women with a gestational weight gain of 13.8 kg and a total protein deposition during pregnancy of 686 g, daily protein deposition was estimated as 1.9 g in the second trimester and 7.4 g in the third trimester (Butte et al., 2003; WHO/FAO/UNU, 2007). For protein deposition towards the end of pregnancy, IoM (2005) derived a mean value of 7.2 g/d based on six studies estimating the increase in whole-body potassium during pregnancy in 120 women. They then assumed that nitrogen accretion during the second trimester is only about half of that observed in the third trimester, leading to an estimated value for protein deposition of 3.6 g/d for the second trimester.

Based on an efficiency of dietary protein utilisation of 42 %, WHO/FAO/UNU (2007) estimated that an additional 1, 9 and 31 g protein/d in the first, second and third trimesters, respectively, are required to support a gestational weight gain of 13.8 kg.

The Panel notes that a value of 42 % for the efficiency of dietary protein utilisation is low, and cannot see a plausible reason to depart from the value of 47 % derived for adults for maintenance of body protein (see also Section 5.1.1.3.).

#### **5.1.5. Protein requirement during lactation**

The additional protein requirement for milk production can be estimated factorially from milk protein output and the efficiency of dietary protein utilisation for milk protein production. The efficiency of protein utilisation for milk protein production is unknown and was taken to be the same as for protein deposition in the non-lactating adult (47 %). In the report of WHO/FAO/UNU (2007) mean rates of milk production by well-nourished women exclusively breastfeeding their infants during the first six months postpartum, and partially breastfeeding in the second six months postpartum, together with the mean concentrations of protein and non-protein nitrogen in human milk, were used to calculate mean milk protein output. The factor of 6.25 was used to convert milk nitrogen to protein. Thus, the additional dietary protein requirement during lactation will be an amount of dietary protein equal to milk protein output, divided by an efficiency of protein utilisation of 47 %. Assuming a CV of 12 %, the additional protein intakes during the first six months of lactation were estimated as 19 g protein/d, falling to 13 g protein/d after six months.

The Panel accepts the approach of WHO/FAO/UNU (2007).

## **5.2. Protein intake and health consequences**

Protein requirement and PRI are derived from nitrogen balance but several health outcomes associated with protein intake could also be considered as criteria for setting DRVs for protein. It is conceivable that in the event of sufficient evidence for a positive effect on health, a PRI for protein above the PRI derived from nitrogen balance and factorial estimates would result. In addition, potentially adverse effects on health should be taken into account when assessing a protein intake above the PRI derived from nitrogen balance.

### **5.2.1. Muscle mass**

The major anabolic influences on muscle are contractile activity and feeding. Ingestion of sufficient dietary energy and protein is a prerequisite for muscle protein synthesis and maintenance of muscle mass and function.

As a result of feeding, anabolism occurs chiefly by an increase in protein synthesis. Insulin has a permissive role in increasing synthesis, and the availability of amino acids is crucial for net anabolism. *In vivo*, amino acids display an anabolic effect (Giordano et al., 1996; Volpi et al., 1996) and were shown to stimulate muscle protein synthesis (Bohe et al., 2003; Liu et al., 2002; Nair and Short, 2005; Nygren and Nair, 2003). There was no effect of a dietary protein level above the PRI on muscle mass and protein content, and a high protein diet of around 2 g/kg body weight per day has not been demonstrated to modulate skeletal protein

synthesis in both exercising and non-exercising human subjects (Bolster et al., 2005; IoM, 2005; Juillet et al., 2008) or animals (Almurshed and Grunewald, 2000; Chevalier et al., 2009; Masanés et al., 1999; Morens et al., 2001; Taillandier et al., 2003). However, increasing protein intake above the individual requirement increases amino acid oxidation and modifies protein turnover. When protein intake is increased from around 1 g/kg body weight per day to 2 g/kg body weight per day, the increase of amino acid oxidation is associated with the stimulation of protein breakdown rates in the fasted state, and a strong inhibition in the fed state, whereas whole-body protein synthesis rates are little affected (Forsslund et al., 1998; Fouillet et al., 2008; Harber et al., 2005; Morens et al., 2003; Pacy et al., 1994; Price et al., 1994).

The branched-chain amino acids (BCAA) (leucine, valine, isoleucine), particularly leucine, have been demonstrated to act as a signal for muscle protein synthesis *in vitro* (Buse and Reid, 1975; Busquets et al., 2002; Dardevet et al., 2000; Fulks et al., 1975; Hong and Layman, 1984; Kimball et al., 1998; Kimball et al., 1999; Li and Jefferson, 1978; Mitch and Clark, 1984; Mordier et al., 2000; Tischler et al., 1982). *In vivo* experiments in animal models have been less consistent but confirm *in vitro* results that leucine acts as a signal that up-regulates muscle protein synthesis and/or down-regulates muscle protein degradation (Anthony et al., 2000; Dardevet et al., 2002; Funabiki et al., 1992; Juillet et al., 2004; Layman and Grogan, 1986; McNurlan et al., 1982; Nagasawa et al., 2002; Rieu et al., 2003). In contrast, there is limited information available on the influence of leucine alone on muscle protein synthesis in humans (Koopman et al., 2005; Nair et al., 1992; Schwenk and Haymond, 1987; Sherwin, 1978; Tessari et al., 1985). At present, there is no convincing evidence that chronic leucine supplementation above the requirement of 39 mg/kg body weight per day is efficient in promoting an increase in muscle mass (Balage and Dardevet, 2010; Leenders et al., 2011). Thus, when the intake of protein is at the PRI based on nitrogen balance, and when amino acid requirements are met, an additional intake of leucine has no further effect on muscle mass.

The Panel considers that in healthy adults the available data on the effects of dietary protein intake on muscle mass and function do not provide evidence that it can be considered as a criterion for setting a PRI for protein. There are no data showing that an additional intake of protein would increase muscle mass in different age groups who are in nitrogen balance, including subjects undertaking endurance or resistance exercise. There are also no data showing that an additional protein intake would increase muscle growth in children.

## 5.2.2. Body weight control and obesity

### 5.2.2.1. Infants

It has been proposed that the well-known difference in growth observed between formula-fed and breast-fed infants may be related to differences in protein intake estimated to be 55-80 % higher in formula-fed infants compared to breast-fed infants (Alexy et al., 1999). In addition, it has been suggested that a higher protein intake may contribute to an enhanced insulin secretion and release of insulin-like growth factor (IGF)-1 and IGF-binding protein (IGFBP)-1, which was observed in prospective feeding studies with infant formulae of different protein content (13, 15 or 18 g protein/L) and a breast-fed control group (Axelsson, 2006).

In a double-blind, randomised controlled manner the European Childhood Obesity Project explored whether two types of infant formulae (standard infant formula and follow-on formula) with either lower or higher protein content (1.8 vs. 2.9 g/100 kcal for infant formula and 2.2 vs. 4.4 g/100 kcal for follow-on formula, all complying with European regulatory standards) fed during the first year of life resulted in different growth in the first two years of life (Grote et al., 2010; Koletzko et al., 2009). A reference group of breast-fed infants was also studied. The mean weight attained at 24 months was 12.4 kg and 12.6 kg for the lower- and higher-protein group, respectively; the adjusted z-score for weight-for-length was 0.20 (95% CI 0.06–0.34; p=0.005) higher in the higher-protein formula group than in the lower-protein formula group. Children fed lower-protein formula did not differ from breast-fed children with respect to weight-for-length and BMI, but weight and length were higher. Whether this statistically significant but small difference in growth observed in infants fed higher-protein formula persists and is related to obesity risk in later life is the subject of ongoing investigations. Currently, these preliminary results do not allow conclusions to be made on the effects of protein intake with regard to obesity development.

The Panel considers that the results from these studies are not suitable for the derivation of a PRI or a UL for protein for infants and children.

#### 5.2.2.2. Adults

Controlled studies in humans have investigated whether an increase in protein intake (as E%) *ad libitum* induces a decrease in body weight and adiposity. However, these studies are difficult to interpret with respect to whether the effects observed are due to an increase in dietary protein intake or to the concomitant modification of carbohydrate and/or fat intakes, and whether any observed effect of an increase in dietary protein intake would be sustainable (Brehm et al., 2003; Foster et al., 2003; Larsen et al., 2010; Samaha et al., 2003; Skov et al., 1999b; Weigle et al., 2005; Westterterp-Plantenga et al., 2004; Yancy et al., 2004). A recent review of the literature concluded that there is strong and consistent evidence that when energy intake is controlled, the macronutrient proportion of the diet is not directly related to weight loss (USDA/HHS, 2010).

The Panel considers that these data cannot be used to derive a PRI for protein for adults.

#### 5.2.3. Insulin sensitivity and glucose control

Contradictory results have been obtained for the effects of an increase in protein intake above the PRI on insulin sensitivity and glucose tolerance. Some human studies showed no effects of a high protein intake on insulin sensitivity and glucose tolerance (Kitagawa et al., 1998; Tsunehara et al., 1990), but a high protein intake was found to be accompanied by an increased insulin secretion and demand (Linn et al., 2000). In other studies, a high protein intake was shown to improve insulin sensitivity and glucose tolerance in humans (Baba et al., 1999; Gannon et al., 2003; Layman et al., 2003; Piatti et al., 1994; Sharman et al., 2002; Volek et al., 2002) and animals (Karabatas et al., 1992; Lacroix et al., 2004; Wang et al., 1998). A beneficial effect of a high-protein diet on insulin resistance and glucose homeostasis has also been reported with a reduced calorie diet regardless of weight-loss (Farnsworth et al., 2003).

Studies conducted *in vitro* or in animal models have suggested that exposure to high levels of BCAA could have a deleterious effect on insulin signalling leading to impaired insulin sensitivity and glucose tolerance (Nair and Short, 2005; Patti et al., 1998; Tremblay and Marette, 2001). Some studies using metabolomic profiling in humans also suggested a BCAA-related metabolite signature that could be correlated with insulin resistance in obese subjects (Newgard et al., 2009) or with future diabetes (Wang et al., 2011). In contrast, prolonged leucine supplementation (7.5 g/d) in elderly type 2 diabetics habitually consuming an adequate amount of dietary protein did not modulate their glycaemic control (Leenders et al., 2011) and BCAA levels were even associated with a decrease in insulin resistance with weight loss (Shah et al., 2012). Other studies, including animal experiments, also did not confirm that increasing dietary leucine was always associated with a deleterious effect on insulin signalling (Leenders and van Loon, 2011; Macotela et al., 2011; Noatsch et al., 2010). The Panel considers that there are insufficient data to conclude on the possible effect of BCAA on insulin sensitivity and glucose tolerance.

The Panel considers that these data cannot be used to derive a PRI or a UL for protein for healthy subjects.

#### 5.2.4. Bone health

Protein and calcium are main components of bone structure and it is widely accepted that protein deficiency increases the risk of bone fragility and fracture (Dawson-Hughes, 2003; Hannan et al., 2000; Kerstetter et al., 2000; Munger et al., 1999; Promislow et al., 2002; Skov et al., 2002; Zernicke et al., 1995). In several epidemiological studies, bone mineral density was positively related to protein intake (Chiu et al., 1997; Cooper et al., 1996; Devine et al., 2005; Geinoz et al., 1993; Hannan et al., 2000; Lau et al., 1998; Promislow et al., 2002; Teegarden et al., 1998; Tucker et al., 2001).

Although protein is essential for bone health, it has been observed that an increase in protein intake could also be associated with an increase in urinary calcium excretion. It was first hypothesised that this could originate from an activation of bone resorption in order to provide calcium for the neutralisation of the acid load produced by the oxidation of sulphur amino acids (Barzel and Massey, 1998). However, an increase in

protein intake is often associated with an increase in calcium intake (Heaney, 1998), and also induces an increase in calcium absorption (Kerstetter et al., 1998, 2003) that can be related to the increased urinary calcium. In addition, the regulation of body acid load is a complex process in which urinary acidity is not directly related to blood acidity; moreover, the theory that considers bone mineral mobilisation as the main physiological system involved in the regulation of extracellular hydrogen ion concentration is questionable since it does not take into account the major role of both the respiratory and the renal tubular systems in this regulation (Fenton et al., 2009). Some studies have shown a positive relationship between protein intake and the risk of bone fracture (Abelow et al., 1992; Frassetto et al., 2000; Hegsted, 1986), whereas others have found no clear association (Meyer et al., 1997; Mussolino et al., 1998) or have shown an inverse association (Munger et al., 1999). Intervention studies did not show clear effects of a protein intake above the PRI on markers of bone formation or resorption (Cao et al., 2011; Darling et al., 2009; Fenton et al., 2009).

The Panel considers that the available evidence is insufficient to be taken into consideration when deriving a PRI or a UL for protein.

### 5.2.5. Kidney function

Protein intake is a modulator of renal function and increases the glomerular filtration rate (GFR) (Brändle et al., 1996). An increase in amino acid catabolism induced by an increase in protein intake increases the production of amino acid-derived metabolites such as bicarbonate, ammonia and urea which require elimination from the body, e.g. via the kidneys.

High protein diets have been found to be associated with increases in blood urea levels and urinary urea excretion, to promote plasma vasopressin, to increase creatinine clearance, and to result in a transient increase in kidney size in humans (Brändle et al., 1996; Diamond, 1990; Gin et al., 2000; Jenkins et al., 2001; Lentine and Wrone, 2004; Zeller, 1991) and animals (Dunger et al., 1997; Hammond and Janes, 1998; Lacroix et al., 2004; Schoknecht and Pond, 1993). High intakes of protein by patients with renal disease contribute to the deterioration of kidney function, and a reduction of protein intake is usually beneficial to subjects with renal insufficiency (Klahr et al., 1994; Knight et al., 2003; Maroni and Mitch, 1997) and possibly also to subjects with microalbuminuria (Friedman, 2004). In contrast, protein intake at the PRI based on nitrogen balance is not a risk factor for renal insufficiency in healthy subjects (Locatelli et al., 1991; Skov et al., 1999a; Wiegmann et al., 1990). According to the available evidence (WHO/FAO/UNU, 2007), the decline of GFR that occurs with advancing age in healthy subjects cannot be attenuated by reducing dietary protein intake below the PRI based on nitrogen balance.

As reported in the opinion on DRVs for water (EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2010) urine osmolarity is physiologically limited between about 50 and 1,400 mOsm/L, and dehydration of more than 10 % at high ambient temperatures is a serious risk for a life-threatening heat stroke, with elevated body temperature, inadequate cardiac output leading to reduced perfusion of tissues and eventually to rhabdomyolysis (i.e. rapid breakdown of skeletal muscle), and organ failure (Bouchama and Knochel, 2002). This risk is particularly high in infants with gastro-enteritis and receiving a formula with a high potential renal solute load (Fomon, 1993). Water required for the excretion of solutes is determined by the composition of the diet and by the concentrating capacity of the kidneys. Because the protein content of the diet is, as a rule, the main determinant of the potential renal solute load, which needs water for excretion, a very high protein intake (around 20 E%, e.g. through exclusive consumption of cow's milk), with a consecutive increased production of urea, can severely impair the water balance of infants, particularly when no other liquids are consumed and/or extrarenal water losses, e.g. through diarrhoea, are increased.

The Panel considers that the available evidence is insufficient to be taken into consideration when deriving a UL for protein.

### 5.2.6. Capacity of the urea cycle

It is established that there is adequate capacity in the human metabolism to adapt to a large range of protein intakes above the PRI based on nitrogen balance. This is mainly due to the adaptation of amino acid catabolic pathways and it is established that amino acid oxidation varies at a rate dependent on the habitual protein intake. The level of protein intake has been evaluated in relation to the capacity of the urea cycle to

control the transfer of ammonia released from amino acid deamination to urea (AFSSA, 2007). It was concluded that for a healthy human male adult, protein intakes between 0.83 and 2.2 g/kg body weight per day (around 10 to 27 E%) are considered as safe, whilst IoM (2005) concludes that the maximum rate of urea production of a 70 kg male not habitually consuming a high-protein diet corresponds to a protein intake of 250 g/d or about 40 E%.

The Panel considers that the available evidence is insufficient to be taken into consideration when deriving a UL for protein.

### 5.2.7. Tolerance of protein

IoM (2005) quotes some reports of very high protein intakes up to 35 E% without adverse effects, whereas acute adverse effects were reported for intakes  $\geq 45$  E% and lethal outcomes occurred when such a diet was consumed by adults for several weeks. In Europe, adult protein intakes at the upper end of the intake distributions (90-97.5<sup>th</sup> percentile) have been reported to be between 17 and 27 E% (Appendix 3B).

The available data on the tolerance of dietary protein are not sufficient to derive a UL for protein.

## 6. Data on which to base Dietary Reference Values (DRVs)

### 6.1. Protein requirement of adults

The criterion of adequacy for the protein intake is the lowest intake that is sufficient to achieve body nitrogen equilibrium (zero balance), during energy balance. The analysis of available nitrogen balance data performed by Rand et al. (2003) concluded that the best estimate of average requirement for healthy adults was the median requirement of 105 mg N/kg body weight per day or 0.66 g protein/kg body weight per day ( $N \times 6.25$ ). The 97.5<sup>th</sup> percentile of the distribution of requirements within a population was estimated as 133 mg N/kg body weight per day, or 0.83 g protein/kg body weight per day. This quantity should meet the requirement of most (97.5 %) of the healthy adult population, and is therefore proposed as the PRI for protein for adults. For older adults, the protein requirement is considered to be equal to that of adults, as data are insufficient to establish that the requirement for healthy older adults is different from that of healthy younger adults. Thus, the PRI of 0.83 g/kg body weight per day is proposed for all adults, including older adults. The protein requirement per kg body weight is considered to be the same for both sexes and for all body weights. The PRI of 0.83 g/kg body weight per day is applicable both to high quality protein and to protein in mixed diets.

### 6.2. Protein requirement of infants and children

The protein requirement of infants and children can be defined as the minimum intake that will allow nitrogen equilibrium at an appropriate body composition during energy balance at moderate physical activity, plus the needs associated with the deposition of tissues consistent with growth and good health (WHO/FAO/UNU, 2007).

The Panel accepted the approach of WHO/FAO/UNU (2007) in which estimates of the protein requirement from six months to adulthood were derived from a factorial model. In selecting values for maintenance and growth efficiency for ages greater than six months, the likelihood that mixed diets consumed after weaning are utilised less efficiently is taken into account.

An average maintenance value of 0.66 g protein/kg body weight per day was applied to children and infants from 6 months to 18 years. Average daily needs for dietary protein for growth were estimated from average daily rates of protein deposition, and an efficiency of utilisation of dietary protein for growth of 58 % was assumed. The average requirement was then estimated as the sum of the maintenance and growth requirements.

The PRI was estimated based on the average requirement plus 1.96 SD; for this, a combined SD was calculated from the SD for growth for the respective age (see Appendix 4), which was adjusted for efficiency of dietary protein utilisation (58 %), and from the SD for maintenance (based on a CV of 12 % for all ages).

### 6.3. Protein requirement during pregnancy

The Panel follows the approach (WHO/FAO/UNU, 2007) in which the additional protein intake needed during pregnancy was derived from the newly deposited protein, taking into account efficiency of protein utilisation and the maintenance costs associated with increased body weight. Mean total protein deposition and daily protein deposition in each trimester was estimated indirectly from measurements of total body potassium accretion, and calculated for an average weight gain of 13.8 kg (the mid-point of the recommended weight gain range for women with normal pre-pregnancy weight) (IoM and NRC, 2009; WHO/FAO/UNU, 2007). Efficiency of protein utilisation was taken by the Panel to be 47 %. The additional maintenance costs were based upon the mid-trimester gain in maternal body weight, and on the adult maintenance value of 0.66 g/kg body weight per day. The PRI was estimated by adding 1.96 SD, with the SD calculated on the basis of a CV of 12 % to give an additional 1, 9 and 28 g protein/d in the first, second and third trimesters, respectively (Table 11).

**Table 11:** Derivation of Dietary Reference Values for protein during pregnancy

Trimester	Mid-trimester weight gain (kg)	Additional protein for maintenance (g/d) <sup>1</sup>	Protein deposition (g/d)	Protein deposition, adjusted for efficiency <sup>2</sup> (g/d)	Additional protein requirement (g/d)	PRI, additional intake <sup>3</sup> (g/d)
1	0.8	0.5	0	0	0.5	1
2	4.8	3.2	1.9	4.0	7.2	9
3	11	7.3	7.4	15.7	23	28

<sup>1</sup>Mid-trimester increase in weight x average requirement (AR) for maintenance of protein for adults of 0.66 g/kg body weight per day

<sup>2</sup>Protein deposition adjusted for the efficiency of protein utilisation during pregnancy: 47 %

<sup>3</sup>Calculated as the average requirement plus allowance for estimated coefficient of variation of 12 %

### 6.4. Protein requirement during lactation

The Panel accepted the factorial method via milk protein output assessment (from milk volumes and from the content of both protein nitrogen and NPN) and via calculation of the amount of dietary protein needed for milk protein production with an efficiency of utilisation of 47 %. The factor 6.25 was used to convert nitrogen to protein. The PRI was estimated by adding 1.96 SD, with the SD calculated on the basis of a CV of 12 % to give an additional 19 g protein/d during the first six months of lactation, and 13 g protein/d after six months.

### 6.5. Safety of protein intakes above the PRI

A UL cannot be derived. Concerns about the potential detrimental effects of very high protein intake remain controversial. Acute adverse effects have been reported for protein intakes  $\geq 45$  E%, but very high protein intakes up to 35 E% have not been associated with adverse effects in some reports. It can be concluded that in adults an intake of twice the PRI is safe. Such intakes from mixed diets are regularly consumed by some physically active and healthy individuals in Europe. Intakes of 3–4 times the PRI have been observed without apparent adverse effects or benefits.

Data from food consumption surveys show that actual mean protein intakes of adults in Europe are at, or more often above, the PRI of 0.83 g/kg body weight per day. Protein intakes as high as 1.7 g/kg body weight per day (95<sup>th</sup> percentile of protein intake of Dutch men aged  $\geq 65$  years) or 27 E% have been observed (see Appendix 3B).

In infants, a very high protein intake (around 20 E%) can severely impair the water balance, particularly when no other liquids are consumed and/or extrarenal water losses are increased. Consequently, such high protein intakes should be avoided in the first year of life.



## CONCLUSIONS

The Panel concludes that an Average Requirement (AR) and a Population Reference Intake (PRI) for protein can be derived for adults, infants and children, and pregnant and lactating women based on nitrogen balance studies and on factorial estimates of the nitrogen needed for deposition of newly formed tissue and for milk output. The Panel also considered several health outcomes that may be associated with protein intake; however, the available data were considered insufficient to help in setting DRVs.

The Panel concludes that the available data are not sufficient to establish a Tolerable Upper Intake Level (UL) for protein.

**Table 12:** Summary of Dietary Reference Values for protein

Age (years)	AR (g/kg bw x d <sup>-1</sup> )	PRI (g/kg bw x d <sup>-1</sup> )	Reference weight (kg) <sup>1</sup>		PRI (g/d)	
			males (m)	females (f)	m	f
			0.5	1.12	1.31	7.7
1	0.95	1.14	10.2	9.5	12	11
1.5	0.85	1.03	11.6	10.9	12	11
2	0.79	0.97	12.7	12.1	12	12
3	0.73	0.90	14.7	14.2	13	13
4	0.69	0.86	17.0	16.4	15	14
5	0.69	0.85	19.2	18.7	16	16
6	0.72	0.89	21.5	21.1	19	19
7	0.74	0.91	24.3	23.8	22	22
8	0.75	0.92	27.4	26.8	25	25
9	0.75	0.92	30.6	30.0	28	28
10	0.75	0.91	33.8	33.7	31	31
11	0.75 (m), 0.73 (f)	0.91 (m), 0.90 (f)	37.3	37.9	34	34
12	0.74 (m), 0.72 (f)	0.90 (m), 0.89 (f)	41.5	42.6	37	38
13	0.73 (m), 0.71 (f)	0.90 (m), 0.88 (f)	46.7	47.5	42	42
14	0.72 (m), 0.70 (f)	0.89 (m), 0.87 (f)	52.7	51.6	47	45
15	0.72 (m), 0.69 (f)	0.88 (m), 0.85 (f)	59.0	54.6	52	46
16	0.71 (m), 0.68 (f)	0.87 (m), 0.84 (f)	64.1	56.4	56	47
17	0.70 (m), 0.67 (f)	0.86 (m), 0.83 (f)	67.5	57.4	58	48
18-59	0.66	0.83	74.6	62.1	62	52
≥ 60	0.66	0.83	73.5	66.1	61	55
Pregnant women <sup>2</sup>						
1 <sup>st</sup> trimester						+1
2 <sup>nd</sup> trimester						+9
3 <sup>rd</sup> trimester						+28
Lactating women <sup>2</sup>						
0-6 months <i>post-partum</i>						+19
>6 months <i>post-partum</i>						+13

<sup>1</sup> For infants and children, based upon the 50<sup>th</sup> percentile of the reference body weights (kg) of European children (van Buuren et al., 2012). For adults, based upon weighted median body weights (kg) of European men and women (SCF, 1993)

<sup>2</sup> In addition to the PRI for non-pregnant women

## REFERENCES

- Abelow BJ, Holford TR and Insogna KL, 1992. Cross-cultural association between dietary animal protein and hip fracture: a hypothesis. *Calcified Tissue International*, 50, 14-18.
- Abrasheva M, Angelova K, Baikova D, Duleva V, Ivanova L, Jordanov B, Ovcharova D, Petrova S, Vatrlova K and Vukov M, 1998. Dietary and nutritional status survey of the population in Bulgaria. National Center of Hygiene, Medical Ecology and Nutrition, 230 pp.
- AFSSA (Agence Française de Sécurité Sanitaire des Aliments), 2007. Apport en protéines: consommation, qualité, besoins et recommandations [Protein intake: dietary intake, quality, requirements and recommendations]. 461 pp.
- Alexy U, Kersting M, Sichert-Hellert W, Manz F and Schoch G, 1999. Macronutrient intake of 3- to 36-month-old German infants and children: results of the DONALD Study. Dortmund Nutritional and Anthropometric Longitudinally Designed Study. *Annals of Nutrition and Metabolism*, 43, 14-22.
- Almurshed KS and Grunewald KK, 2000. Dietary protein does not affect overloaded skeletal muscle in rats. *Journal of Nutrition*, 130, 1743-1748.
- Andersen N, Fagt S, Groth M, Hartkopp H, Møller A, Ovesen L and Warming D, 1996. Danskernes kostvaner 1995. Hovedresultater [The Danish diet 1995. Main results]. Levnedsmiddelstyrelsen, Søborg, Denmark, 298 pp.
- Anonymous, 2008. Ergebnisbericht, Teil 2. Nationale Verzehrsstudie II [Report, Part 2. National Nutrition Survey II]. Max Rubner Institut, Bundesforschungsinstitut für Ernährung und Lebensmittel, 193 pp.
- ANSES/CIQUAL (Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement, et du travail/Centre d'information sur la qualité des aliments), 2008. French food composition table version 2008. Available from: <http://www.afssa.fr/TableCIQUAL/index.htm>, accessed 19 January 2012.
- Anthony JC, Yoshizawa F, Anthony TG, Vary TC, Jefferson LS and Kimball SR, 2000. Leucine stimulates translation initiation in skeletal muscle of postabsorptive rats via a rapamycin-sensitive pathway. *Journal of Nutrition*, 130, 2413-2419.
- Association of Official Analytical Chemists, 1984. Bioassay methods, "Protein Efficiency Ratio, Rat Bioassay, Final action", sections 43.253-43.257 and "Protein Efficiency Ratio, Calculation Method, First Action", sections 43.258 to 43.267. In: AOAC Official Methods of Analysis. Ed Association of Official Analytical Chemists. Association of Official Analytical Chemists, Gaithersburg, USA, 877-880.
- Axelsson I, 2006. Effects of high protein intakes. Nestlé Nutrition Workshop Series. Paediatric Programme, 58, 121-129.
- Baba NH, Sawaya S, Torbay N, Habbal Z, Azar S and Hashim SA, 1999. High protein vs high carbohydrate hypoenergetic diet for the treatment of obese hyperinsulinemic subjects. *International Journal of Obesity and Related Metabolic Disorders*, 23, 1202-1206.
- Baglieri A, Mahe S, Benamouzig R, Savoie L and Tome D, 1995. Digestion patterns of endogenous and different exogenous proteins affect the composition of intestinal effluents in humans. *Journal of Nutrition*, 125, 1894-1903.
- Balage M and Dardevet D, 2010. Long-term effects of leucine supplementation on body composition. *Current Opinion in Clinical Nutrition and Metabolic Care*, 13, 265-270.
- Barzel US and Massey LK, 1998. Excess dietary protein can adversely affect bone. *Journal of Nutrition*, 128, 1051-1053.
- Bates E, Lennox A, Bates C and Swan G, 2011. National Diet and Nutrition Survey. Headline results from Years 1 and 2 (combined) of the Rolling Programme (2008/2009 - 2009/2010). A Survey carried out on the behalf of the Food Standards Agency and the Department of Health, 68 pp.

- Becker W and Pearson M, 2002. Riksmaten 1997-1998. Befolkningens kostvanor och näringsintag. Metod- och resultatanalys [Riksmaten 1997-1998. Dietary habits and nutrient intake in Sweden. Benchmarking analysis]. Livsmedelsverket, 201 pp.
- Biro L, Regoly-Merei A, Nagy K, Peter S, Arato G, Szabo C, Martos E and Antal M, 2007. Dietary habits of school children: representative survey in metropolitan elementary schools. Part two. *Annals of Nutrition and Metabolism*, 51, 454-460.
- Bohe J, Low A, Wolfe RR and Rennie MJ, 2003. Human muscle protein synthesis is modulated by extracellular, not intramuscular amino acid availability: a dose-response study. *Journal of Physiology*, 552, 315-324.
- Bolster DR, Pikosky MA, Gaine PC, Martin W, Wolfe RR, Tipton KD, Maclean D, Maresh CM and Rodriguez NR, 2005. Dietary protein intake impacts human skeletal muscle protein fractional synthetic rates after endurance exercise. *American Journal of Physiology, Endocrinology and Metabolism*, 289, E678-683.
- Bos C, Gaudichon C and Tome D, 2002. Isotopic studies of protein and amino acid requirements. *Current Opinion in Clinical Nutrition and Metabolic Care*, 5, 55-61.
- Bos C, Juillet B, Fouillet H, Turlan L, Dare S, Luengo C, N'Tounda R, Benamouzig R, Gausseres N, Tome D and Gaudichon C, 2005. Postprandial metabolic utilization of wheat protein in humans. *American Journal of Clinical Nutrition*, 81, 87-94.
- Bouchama A and Knochel JP, 2002. Heat stroke. *New England Journal of Medicine*, 346, 1978-1988.
- Brändle E, Sieberth HG and Hautmann RE, 1996. Effect of chronic dietary protein intake on the renal function in healthy subjects. *European Journal of Clinical Nutrition*, 50, 734-740.
- Brehm BJ, Seeley RJ, Daniels SR and D'Alessio DA, 2003. A randomized trial comparing a very low carbohydrate diet and a calorie-restricted low fat diet on body weight and cardiovascular risk factors in healthy women. *Journal of Clinical Endocrinology and Metabolism*, 88, 1617-1623.
- Bunker VW, Lawson MS, Stansfield MF and Clayton BE, 1987. Nitrogen balance studies in apparently healthy elderly people and those who are housebound. *British Journal of Nutrition*, 57, 211-221.
- Buse MG and Reid SS, 1975. Leucine. A possible regulator of protein turnover in muscle. *Journal of Clinical Investigation*, 56, 1250-1261.
- Busquets S, Alvarez B, Lopez-Soriano FJ and Argiles JM, 2002. Branched-chain amino acids: a role in skeletal muscle proteolysis in catabolic states? *Journal of Cellular Physiology*, 191, 283-289.
- Butte NF, Ellis KJ, Wong WW, Hopkinson JM and Smith EO, 2003. Composition of gestational weight gain impacts maternal fat retention and infant birth weight. *American Journal of Obstetrics and Gynecology*, 189, 1423-1432.
- Calloway DH, 1974. Nitrogen balance during pregnancy. *Current Concepts in Nutrition*, 2, 79-94.
- Campbell WW, Trappe TA, Wolfe RR and Evans WJ, 2001. The recommended dietary allowance for protein may not be adequate for older people to maintain skeletal muscle. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 56, M373-380.
- Campbell WW and Leidy HJ, 2007. Dietary protein and resistance training effects on muscle and body composition in older persons. *Journal of the American College of Nutrition*, 26, 696S-703S.
- Campbell WW, Johnson CA, McCabe GP and Carnell NS, 2008. Dietary protein requirements of younger and older adults. *American Journal of Clinical Nutrition*, 88, 1322-1329.
- Cao JJ, Johnson LK and Hunt JR, 2011. A diet high in meat protein and potential renal acid load increases fractional calcium absorption and urinary calcium excretion without affecting markers of bone resorption or formation in postmenopausal women. *Journal of Nutrition*, 141, 391-397.

- Chevalier L, Bos C, Gryson C, Luengo C, Walrand S, Tome D, Boirie Y and Gaudichon C, 2009. High-protein diets differentially modulate protein content and protein synthesis in visceral and peripheral tissues in rats. *Nutrition*, 25, 932-939.
- Chiu JF, Lan SJ, Yang CY, Wang PW, Yao WJ, Su LH and Hsieh CC, 1997. Long-term vegetarian diet and bone mineral density in postmenopausal Taiwanese women. *Calcified Tissue International*, 60, 245-249.
- Cifkova R and Skodova Z, 2004. Dlouhodobé trendy hlavních rizikových faktorů kardiovaskulárních onemocnění v české populaci [Longitudinal trends in major cardiovascular disease risk factors in the Czech population]. *Casopis Lekarů Ceských [Czech medical journal]*, 143, 219-226.
- Cooper C, Atkinson EJ, Hensrud DD, Wahner HW, O'Fallon WM, Riggs BL and Melton LJ, 3rd, 1996. Dietary protein intake and bone mass in women. *Calcified Tissue International*, 58, 320-325.
- Cuthbertson D, Smith K, Babraj J, Leese G, Waddell T, Atherton P, Wackerhage H, Taylor PM and Rennie MJ, 2005. Anabolic signaling deficits underlie amino acid resistance of wasting, aging muscle. *FASEB Journal*, 19, 422-424.
- D-A-CH (Deutsche Gesellschaft für Ernährung - Österreichische Gesellschaft für Ernährung - Schweizerische Gesellschaft für Ernährungsforschung - Schweizerische Vereinigung für Ernährung), 2008. Referenzwerte für die Nährstoffzufuhr [Reference values for nutrient intakes]. Umschau Braus Verlag, Frankfurt am Main, Germany, 240 pp.
- Dardevet D, Sornet C, Balage M and Grizard J, 2000. Stimulation of in vitro rat muscle protein synthesis by leucine decreases with age. *Journal of Nutrition*, 130, 2630-2635.
- Dardevet D, Sornet C, Bayle G, Prugnaud J, Pouyet C and Grizard J, 2002. Postprandial stimulation of muscle protein synthesis in old rats can be restored by a leucine-supplemented meal. *Journal of Nutrition*, 132, 95-100.
- Darling AL, Millward DJ, Torgerson DJ, Hewitt CE and Lanham-New SA, 2009. Dietary protein and bone health: a systematic review and meta-analysis. *American Journal of Clinical Nutrition*, 90, 1674-1692.
- Davis TA, Fiorotto ML and Reeds PJ, 1993. Amino acid compositions of body and milk protein change during the suckling period in rats. *Journal of Nutrition*, 123, 947-956.
- Dawson-Hughes B, 2003. Interaction of dietary calcium and protein in bone health in humans. *Journal of Nutrition*, 133, 852S-854S.
- de Boer EJ, Hulshof KFAM and Doest DT, 2006. Voedselconsumptie van jonge peuters [Food consumption of young children]. TNO report V6269, 37 pp.
- De Vriese S, Huybrechts I, Moreau M and van Oyen H, 2006. De Belgische Voedselconsumptiepeiling 1 – 2004 [The Belgian food consumption survey 1 - 2004]. WIV/EPI REPORTS B 2006 –016. Wetenschappelijk Instituut Volksgezondheid. Available from: <https://www.wiv-isp.be/epidemio/epinl/foodnl/table04.htm>, accessed 19 January 2012.
- Deharveng G, Charrondiere UR, Slimani N, Southgate DA and Riboli E, 1999. Comparison of nutrients in the food composition tables available in the nine European countries participating in EPIC. *European Prospective Investigation into Cancer and Nutrition. European Journal of Clinical Nutrition*, 53, 60-79.
- Devine A, Dick IM, Islam AF, Dhaliwal SS and Prince RL, 2005. Protein consumption is an important predictor of lower limb bone mass in elderly women. *American Journal of Clinical Nutrition*, 81, 1423-1428.
- Dewey KG, Beaton G, Fjeld C, Lonnerdal B and Reeds P, 1996. Protein requirements of infants and children. *European Journal of Clinical Nutrition*, 50 Suppl 1, S119-147.
- Diamond JR, 1990. Effects of dietary interventions on glomerular pathophysiology. *American Journal of Physiology*, 258, F1-8.
- DoH (Department of Health), 1991. Dietary Reference Values for food energy and nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. HMSO, London, UK, 212 pp.

- Dunger A, Berg S, Kloting I and Schmidt S, 1997. Functional alterations in the rat kidney induced either by diabetes or high protein diet. *Experimental and Clinical Endocrinology and Diabetes*, 105 Suppl 2, 48-50.
- Dunn JM and Speer VC, 1991. Nitrogen requirement of pregnant gilts. *Journal of Animal Science*, 69, 2020-2025.
- EFSA (European Food Safety Authority), 2011a. Evaluation of the FoodEx, the food classification system applied to the development of the EFSA Comprehensive European Food Consumption Database. *EFSA Journal* 2011; 9(3):1970, 27 pp.
- EFSA (European Food Safety Authority), 2011b. Use of the EFSA Comprehensive European Food Consumption Database in exposure assessment. *EFSA Journal* 2011;9(3):2097, 34 pp.
- EFSA Panel on Dietetic Products Nutrition and Allergies (NDA), 2010. Scientific Opinion on Dietary Reference Values for water. *EFSA Journal*, 8(3):1459, 48 pp.
- Elango R, Ball RO and Pencharz PB, 2008a. Indicator amino acid oxidation: concept and application. *Journal of Nutrition*, 138, 243-246.
- Elango R, Ball RO and Pencharz PB, 2008b. Individual amino acid requirements in humans: an update. *Current Opinion in Clinical Nutrition and Metabolic Care*, 11, 34-39.
- Elango R, Humayun MA, Ball RO and Pencharz PB, 2011. Protein requirement of healthy school-age children determined by the indicator amino acid oxidation method. *American Journal of Clinical Nutrition*, 94, 1545-1552.
- Elmadfa I, Meyer A, Nowak V, Hasenegger V, Putz P, Verstraeten R, Remaut-DeWinter AM, Kolsteren P, Dostalova J, Dlouhy P, Trolle E, Fagt S, Biltoft-Jensen A, Mathiessen J, Velsing Groth M, Kambek L, Gluskova N, Voutilainen N, Erkkila A, Vernay M, Krems C, Strassburg A, Vasquez-Caicedo AL, Urban C, Naska A, Efstathopoulou E, Oikonomou E, Tsiotas K, Bountziouka V, Benetou V, Trichopoulou A, Zajkas G, Kovacs V, Martos E, Heavey P, Kelleher C, Kennedy J, Turrini A, Selga G, Sauka M, Petkeviciene J, Klumbiene J, Holm Totland T, Andersen LF, Halicka E, Rejman K, Kowrygo B, Rodrigues S, Pinhao S, Ferreira LS, Lopes C, Ramos E, Vaz Almeida MD, Vlad M, Simcic M, Podgrajsek K, Serra Majem L, Roman Vinas B, Ngo J, Ribas Barba L, Becker V, Fransen H, Van Rossum C, Ocke M and Margetts B, 2009a. European Nutrition and Health Report 2009. *Forum of Nutrition*, 62, 1-405.
- Elmadfa I, Freisling H, Nowak V, Hofstädter D, Hasenegger V, Ferge M, Fröhler M, Fritz K, Meyer AL, Putz P, Rust P, Grossgut R, Mischek D, Kiefer I, Schätzer M, Spanblöchel J, Sturtzel B, Wagner K-H, Zilberszac A, Vojir F and Plsek K, 2009b. Österreichischer Ernährungsbericht 2008 [Austrian Nutrition Report 2008]. Institut für Ernährungswissenschaften der Universität Wien, Bundesministerium für Gesundheit, 454 pp.
- Enghardt-Barbieri H, Pearson M and Becker W, 2006. Riksmaten – Barn 2003. Livsmedels – och näringsintag bland barn i Sverige [Riksmaten - Children 2003. The food and nutritional intake among children in Sweden]. *Livsmedelsverket*, 216 pp.
- FAO/WHO (Food and Agriculture Organization of the United Nations/World Health Organization), 1991. Protein quality evaluation. Report of the joint FAO/WHO expert consultation, Bethesda, MD, USA, 4-8 December 1989. *FAO Food and Nutrition Paper No. 51*, 66 pp.
- FAO/WHO/UNU (Food and Agriculture Organization of the United Nations/World Health Organization/United Nations University), 1985. Energy and protein requirements. Report of a Joint WHO/FAO/UNU Expert Consultation, Rome, 5-17 October 1981. *World Health Organization Technical Report Series No 724*, 206 pp.
- Farnsworth E, Luscombe ND, Noakes M, Wittert G, Argyiou E and Clifton PM, 2003. Effect of a high-protein, energy-restricted diet on body composition, glycemic control, and lipid concentrations in overweight and obese hyperinsulinemic men and women. *American Journal of Clinical Nutrition*, 78, 31-39.

- Fenton TR, Lyon AW, Eliasziw M, Tough SC and Hanley DA, 2009. Meta-analysis of the effect of the acid-ash hypothesis of osteoporosis on calcium balance. *Journal of Bone and Mineral Research*, 24, 1835-1840.
- Fomon SJ, 1993. *Nutrition of normal infants*. Mosby, St. Louis, USA, 475 pp.
- Forslund AH, Hambræus L, Olsson RM, El-Khoury AE, Yu YM and Young VR, 1998. The 24-h whole body leucine and urea kinetics at normal and high protein intakes with exercise in healthy adults. *American Journal of Physiology*, 275, E310-320.
- Forsum E, Sadurskis A and Wager J, 1988. Resting metabolic rate and body composition of healthy Swedish women during pregnancy. *American Journal of Clinical Nutrition*, 47, 942-947.
- Foster GD, Wyatt HR, Hill JO, McGuckin BG, Brill C, Mohammed BS, Szapary PO, Rader DJ, Edman JS and Klein S, 2003. A randomized trial of a low-carbohydrate diet for obesity. *New England Journal of Medicine*, 348, 2082-2090.
- Fouillet H, Bos C, Gaudichon C and Tome D, 2002. Approaches to quantifying protein metabolism in response to nutrient ingestion. *Journal of Nutrition*, 132, 3208S-3218S.
- Fouillet H, Juillet B, Bos C, Mariotti F, Gaudichon C, Benamouzig R and Tome D, 2008. Urea-nitrogen production and salvage are modulated by protein intake in fed humans: results of an oral stable-isotope-tracer protocol and compartmental modeling. *American Journal of Clinical Nutrition*, 87, 1702-1714.
- Frassetto LA, Todd KM, Morris RC, Jr. and Sebastian A, 2000. Worldwide incidence of hip fracture in elderly women: relation to consumption of animal and vegetable foods. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 55, M585-592.
- Friedman AN, 2004. High-protein diets: potential effects on the kidney in renal health and disease. *American Journal of Kidney Diseases*, 44, 950-962.
- Fulks RM, Li JB and Goldberg AL, 1975. Effects of insulin, glucose, and amino acids on protein turnover in rat diaphragm. *Journal of Biological Chemistry*, 250, 290-298.
- Fuller MF, Milne A, Harris CI, Reid TM and Keenan R, 1994. Amino acid losses in ileostomy fluid on a protein-free diet. *American Journal of Clinical Nutrition*, 59, 70-73.
- Fuller MF and Reeds PJ, 1998. Nitrogen cycling in the gut. *Annual Review of Nutrition*, 18, 385-411.
- Fuller MF and Tome D, 2005. In vivo determination of amino acid bioavailability in humans and model animals. *Journal of AOAC International*, 88, 923-934.
- Funabiki R, Yagasaki K, Hara H, Nyumura N, Yoshizawa F and Saito K, 1992. In vivo effect of L-leucine administration on protein synthesis in mice. *The Journal of Nutritional Biochemistry*, 3, 401-407.
- Gabrijelčič Blenkuš M, Gregorič M, Tivadar B, Koch V, Kostanjevec S, Turk Fajdiga V, Žalar A, Lavtar D, Kuhar D and Rozman U, 2009. Prehrambene navade odraslih prebivalcev Slovenije z vidika varovanja zdravja [Food habits of the adult population of Slovenia in terms of health]. Inštitut za varovanje zdravja Republike Slovenije, Pedagoška fakulteta, Univerza v Ljubljani, 186 pp.
- Gaffney-Stomberg E, Insogna KL, Rodriguez NR and Kerstetter JE, 2009. Increasing dietary protein requirements in elderly people for optimal muscle and bone health. *Journal of the American Geriatrics Society*, 57, 1073-1079.
- Gannon MC, Nuttall FQ, Saeed A, Jordan K and Hoover H, 2003. An increase in dietary protein improves the blood glucose response in persons with type 2 diabetes. *American Journal of Clinical Nutrition*, 78, 734-741.
- Gaudichon C, Mahe S, Benamouzig R, Luengo C, Fouillet H, Dare S, Van Oyccke M, Ferriere F, Rautureau J and Tome D, 1999. Net postprandial utilization of [15N]-labeled milk protein nitrogen is influenced by diet composition in humans. *Journal of Nutrition*, 129, 890-895.

- Gaudichon C, Bos C, Morens C, Petzke KJ, Mariotti F, Everwand J, Benamouzig R, Dare S, Tome D and Metges CC, 2002. Ileal losses of nitrogen and amino acids in humans and their importance to the assessment of amino acid requirements. *Gastroenterology*, 123, 50-59.
- Geinoz G, Rapin CH, Rizzoli R, Kraemer R, Buchs B, Slosman D, Michel JP and Bonjour JP, 1993. Relationship between bone mineral density and dietary intakes in the elderly. *Osteoporosis International*, 3, 242-248.
- Gin H, Rigalleau V and Aparicio M, 2000. Lipids, protein intake, and diabetic nephropathy. *Diabetes and Metabolism*, 26 Suppl 4, 45-53.
- Giordano M, Castellino P and DeFronzo RA, 1996. Differential responsiveness of protein synthesis and degradation to amino acid availability in humans. *Diabetes*, 45, 393-399.
- Grote V, von Kries R, Closa-Monasterolo R, Scaglioni S, Gruszfeld D, Sengier A, Langhendries JP and Koletzko B, 2010. Protein intake and growth in the first 24 months of life. *Journal of Pediatric Gastroenterology and Nutrition*, 51 Suppl 3, S117-118.
- Guillet C, Zangarelli A, Mishellany A, Rousset P, Sornet C, Dardevet D and Boirie Y, 2004. Mitochondrial and sarcoplasmic proteins, but not myosin heavy chain, are sensitive to leucine supplementation in old rat skeletal muscle. *Experimental Gerontology*, 39, 745-751.
- Hammond KA and Janes DN, 1998. The effects of increased protein intake on kidney size and function. *Journal of Experimental Biology*, 201, 2081-2090.
- Hannan MT, Tucker KL, Dawson-Hughes B, Cupples LA, Felson DT and Kiel DP, 2000. Effect of dietary protein on bone loss in elderly men and women: the Framingham Osteoporosis Study. *Journal of Bone and Mineral Research*, 15, 2504-2512.
- Harber MP, Schenk S, Barkan AL and Horowitz JF, 2005. Effects of dietary carbohydrate restriction with high protein intake on protein metabolism and the somatotrophic axis. *Journal of Clinical Endocrinology and Metabolism*, 90, 5175-5181.
- Health Council of the Netherlands, 2001. Dietary Reference Intakes: energy, proteins, fats and digestible carbohydrates. Publication no. 2001/19ER, The Hague, The Netherlands, 168 pp.
- Heaney RP, 1998. Excess dietary protein may not adversely affect bone. *Journal of Nutrition*, 128, 1054-1057.
- Hegsted DM, 1986. Calcium and osteoporosis. *Journal of Nutrition*, 116, 2316-2319.
- Hilbig A and Kersting M, 2006. Effects of age and time on energy and macronutrient intake in German infants and young children: results of the DONALD study. *Journal of Pediatric Gastroenterology and Nutrition*, 43, 518-524.
- Hong SO and Layman DK, 1984. Effects of leucine on in vitro protein synthesis and degradation in rat skeletal muscles. *Journal of Nutrition*, 114, 1204-1212.
- Houston DK, Nicklas BJ, Ding J, Harris TB, Tyllavsky FA, Newman AB, Lee JS, Sahyoun NR, Visser M and Kritchevsky SB, 2008. Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) Study. *American Journal of Clinical Nutrition*, 87, 150-155.
- Humayun MA, Elango R, Ball RO and Pencharz PB, 2007. Reevaluation of the protein requirement in young men with the indicator amino acid oxidation technique. *American Journal of Clinical Nutrition*, 86, 995-1002.
- Huybrechts I and De Henauw S, 2007. Energy and nutrient intakes by pre-school children in Flanders-Belgium. *British Journal of Nutrition*, 98, 600-610.
- Hytten FE and Chamberlain G, 1990. *Clinical Physiology in Obstetrics*. Blackwell Scientific Publications, Oxford, UK, 455 pp.

- Iglay HB, Apolzan JW, Gerrard DE, Eash JK, Anderson JC and Campbell WW, 2009. Moderately increased protein intake predominately from egg sources does not influence whole body, regional, or muscle composition responses to resistance training in older people. *Journal of Nutrition, Health and Aging*, 13, 108-114.
- IoM (Institute of Medicine), 1990. *Nutrition During Pregnancy: Part I: Weight Gain, Part II: Nutrient Supplements*. National Academy Press, Washington, D. C., USA, 468 pp.
- IoM (Institute of Medicine), 2005. *Dietary Reference Intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids*. National Academies Press, Washington D.C., USA, 1357 pp.
- IoM and NRC (Institute of Medicine and National Research Council), 2009. *Weight gain during pregnancy: reexamining the guidelines*. National Academies Press, Washington, D.C., USA, 868 pp.
- IUNA (Irish Universities Nutrition Alliance), a. National Children's Food Survey. Available from: <http://www.iuna.net>, accessed 19 January 2012.
- IUNA (Irish Universities Nutrition Alliance), b. The National Teens' Food Survey. Available from: <http://www.iuna.net/>, accessed 19 January 2012.
- IUNA (Irish Universities Nutrition Alliance), c. National Adult Nutrition Survey. Available from: <http://www.iuna.net/>, accessed 19 January 2012.
- Jackson AA, 1995. Salvage of urea-nitrogen and protein requirements. *Proceedings of the Nutrition Society*, 54, 535-547.
- Jayalakshmi VT, Venkatachalam PS and Gopalan C, 1959. Nitrogen balance studies in pregnant women in South India. *Indian Journal of Medical Research*, 47, 86-92.
- Jenkins DJ, Kendall CW, Vidgen E, Augustin LS, van Erk M, Geelen A, Parker T, Faulkner D, Vuksan V, Josse RG, Leiter LA and Connelly PW, 2001. High-protein diets in hyperlipidemia: effect of wheat gluten on serum lipids, uric acid, and renal function. *American Journal of Clinical Nutrition*, 74, 57-63.
- Johansson L and Sovoll K, 1999. Norkost, 1997. Landsomfattende kostholdsundersøkelse blant menn og kvinner i alderen 16-79 år [Norkost, 1997. Nationwide dietary survey among men and women aged 16-79 years] Rapport nr.2/1999. Statens råd for ernæring og fysisk aktivitet, 84 pp.
- Johnson LR, Barret KE, Gishan FK, Merchant JL, Said HM and Wood JD, 2006. *Physiology of the Gastrointestinal Tract*. Elsevier Inc, San Diego, USA, 2080 pp.
- Johnstone FD, Campbell DM and MacGillivray I, 1981. Nitrogen balance studies in human pregnancy. *Journal of Nutrition*, 111, 1884-1893.
- Jones DB, 1941. Factors for converting percentages of nitrogen in foods and feeds into percentages of protein. Circular 183, United States Department of Agriculture, Washington, D.C., USA, 22 pp.
- Jones RD and Maxwell CV, 1982. Growth, reproductive performance and nitrogen balance of gilts as affected by protein intake and stage of gestation. *Journal of Animal Science*, 55, 848-856.
- Juillet B, Fouillet H, Bos C, Mariotti F, Gausseres N, Benamouzig R, Tome D and Gaudichon C, 2008. Increasing habitual protein intake results in reduced postprandial efficiency of peripheral, anabolic wheat protein nitrogen use in humans. *American Journal of Clinical Nutrition*, 87, 666-678.
- Karabatas LM, Lombardo YB and Basabe JC, 1992. High-protein diet: effect on insulin secretion patterns from streptozotocin-diabetic rats and mice. *Acta Physiologica, Pharmacologica et Therapeutica Latinoamericana*, 42, 239-254.
- Katsanos CS, Kobayashi H, Sheffield-Moore M, Aarsland A and Wolfe RR, 2005. Aging is associated with diminished accretion of muscle proteins after the ingestion of a small bolus of essential amino acids. *American Journal of Clinical Nutrition*, 82, 1065-1073.
- Kerstetter JE, O'Brien KO and Insogna KL, 1998. Dietary protein affects intestinal calcium absorption. *American Journal of Clinical Nutrition*, 68, 859-865.



- Kerstetter JE, Looker AC and Insogna KL, 2000. Low dietary protein and low bone density. *Calcified Tissue International*, 66, 313.
- Kerstetter JE, O'Brien KO and Insogna KL, 2003. Low protein intake: the impact on calcium and bone homeostasis in humans. *Journal of Nutrition*, 133, 855S-861S.
- Kimball SR, Horetsky RL and Jefferson LS, 1998. Implication of eIF2B rather than eIF4E in the regulation of global protein synthesis by amino acids in L6 myoblasts. *Journal of Biological Chemistry*, 273, 30945-30953.
- Kimball SR, Shantz LM, Horetsky RL and Jefferson LS, 1999. Leucine regulates translation of specific mRNAs in L6 myoblasts through mTOR-mediated changes in availability of eIF4E and phosphorylation of ribosomal protein S6. *Journal of Biological Chemistry*, 274, 11647-11652.
- King JC, Calloway DH and Margen S, 1973. Nitrogen retention, total body 40 K and weight gain in teenage pregnant girls. *Journal of Nutrition*, 103, 772-785.
- King RH and Brown WG, 1993. Interrelationships between dietary protein level, energy intake, and nitrogen retention in pregnant gilts. *Journal of Animal Science*, 71, 2450-2456.
- Kitagawa T, Owada M, Urakami T and Yamauchi K, 1998. Increased incidence of non-insulin dependent diabetes mellitus among Japanese schoolchildren correlates with an increased intake of animal protein and fat. *Clinical Pediatrics*, 37, 111-115.
- Klahr S, Levey AS, Beck GJ, Caggiula AW, Hunsicker L, Kusek JW and Striker G, 1994. The effects of dietary protein restriction and bloodpressure control on the progression of chronic renal disease. Modification of Diet in Renal Disease Study Group. *New England Journal of Medicine*, 330, 877-884.
- Knight EL, Stampfer MJ, Hankinson SE, Spiegelman D and Curhan GC, 2003. The impact of protein intake on renal function decline in women with normal renal function or mild renal insufficiency. *Annals of Internal Medicine*, 138, 460-467.
- Kobe H, Krzisnik C and Mis NF, 2011. Under- and over-reporting of energy intake in Slovenian adolescents. *Journal of Nutrition Education and Behavior*, 1-10.
- Koletzko B, von Kries R, Closa R, Escribano J, Scaglioni S, Giovannini M, Beyer J, Demmelmair H, Gruszfeld D, Dobrzanska A, Sengier A, Langhendries JP, Rolland Cachera MF and Grote V, 2009. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *American Journal of Clinical Nutrition*, 89, 1836-1845.
- Koopman R, Wagenmakers AJ, Manders RJ, Zorenc AH, Senden JM, Gorselink M, Keizer HA and van Loon LJ, 2005. Combined ingestion of protein and free leucine with carbohydrate increases postexercise muscle protein synthesis in vivo in male subjects. *American Journal of Physiology, Endocrinology and Metabolism*, 288, E645-653.
- Kortebein P, Ferrando A, Lombeida J, Wolfe R and Evans WJ, 2007. Effect of 10 days of bed rest on skeletal muscle in healthy older adults. *JAMA*, 297, 1772-1774.
- Kristiansen AL and Andersen LF, 2009. Småbarnskost – 2 år. Landsomfattende kostholdsundersøkelse blant 2 år gamle barn - Småbarnskost 2007 [Nationwide diet survey among 2-year-old children]. IS-1731, Helsedirektoratet, Mattilsynet og Universitetet i Oslo, 94 pp.
- Kurpad AV, Raj T, El-Khoury A, Beaumier L, Kuriyan R, Srivatsa A, Borgonha S, Selvaraj A, Regan MM and Young VR, 2001. Lysine requirements of healthy adult Indian subjects, measured by an indicator amino acid balance technique. *American Journal of Clinical Nutrition*, 73, 900-907.
- Kyttälä P, Ovaskainen M, Kronberg-Kippilä C, Erkkola M, Tapanainen H, Tuokkola J, Veijola R, Simell O, Knip M and Virtanen SM, 2008. The Diet of Finnish Preschoolers. B32/2008, National Public Health Institute, 158 pp.
- Kyttälä P, Erkkola M, Kronberg-Kippilä C, Tapanainen H, Veijola R, Simell O, Knip M and Virtanen SM, 2010. Food consumption and nutrient intake in Finnish 1-6-year-old children. *Public Health Nutrition*, 13, 947-956.

- Lacroix M, Gaudichon C, Martin A, Morens C, Mathe V, Tome D and Huneau JF, 2004. A long-term high-protein diet markedly reduces adipose tissue without major side effects in Wistar male rats. *American Journal of Physiology, Regulatory, Integrative and Comparative Physiology*, 287, R934-942.
- Larsen TM, Dalskov SM, van Baak M, Jebb SA, Papadaki A, Pfeiffer AF, Martinez JA, Handjieva-Darlenska T, Kunesova M, Pihlsgard M, Stender S, Holst C, Saris WH and Astrup A, 2010. Diets with high or low protein content and glycemic index for weight-loss maintenance. *New England Journal of Medicine*, 363, 2102-2113.
- Lau EM, Kwok T, Woo J and Ho SC, 1998. Bone mineral density in Chinese elderly female vegetarians, vegans, lacto-vegetarians and omnivores. *European Journal of Clinical Nutrition*, 52, 60-64.
- Layman DK and Grogan CK, 1986. Leucine stimulation of skeletal muscle protein synthesis. *Federation Proceedings*, 45, 232.
- Layman DK, Shiue H, Sather C, Erickson DJ and Baum J, 2003. Increased dietary protein modifies glucose and insulin homeostasis in adult women during weight loss. *Journal of Nutrition*, 133, 405-410.
- Leenders M and van Loon LJ, 2011. Leucine as a pharmaconutrient to prevent and treat sarcopenia and type 2 diabetes. *Nutrition Reviews*, 69, 675-689.
- Leenders M, Verdijk LB, van der Hoeven L, van Kranenburg J, Hartgens F, Wodzig WKWH, Saris WHM and van Loon LJC, 2011. Prolonged leucine supplementation does not augment muscle mass or affect glycemic control in elderly type 2 diabetic men. *Journal of Nutrition*, 141, 1070-1076.
- Lentine K and Wrone EM, 2004. New insights into protein intake and progression of renal disease. *Current Opinion in Nephrology and Hypertension*, 13, 333-336.
- Leung WW, Busson F and Jardin C, 1968. Food composition table for use in Africa. A research project sponsored jointly by U.S. Department of Health, Education, and Welfare, Public Health Service, Health Services and Mental Health Administration, National Center for Chronic Disease Control, Nutrition Program, Bethesda (MD), USA, and Food Consumption and Planning Branch, Nutrition Division, Food and Agriculture Organization of the United Nations, Rome, Italy. 306 pp.
- Li JB and Jefferson LS, 1978. Influence of amino acid availability on protein turnover in perfused skeletal muscle. *Biochimica et Biophysica Acta*, 544, 351-359.
- Linn T, Santosa B, Gronemeyer D, Aygen S, Scholz N, Busch M and Bretzel RG, 2000. Effect of long-term dietary protein intake on glucose metabolism in humans. *Diabetologia*, 43, 1257-1265.
- Liu Z, Jahn LA, Wei L, Long W and Barrett EJ, 2002. Amino acids stimulate translation initiation and protein synthesis through an Akt-independent pathway in human skeletal muscle. *Journal of Clinical Endocrinology and Metabolism*, 87, 5553-5558.
- Locatelli F, Alberti D, Graziani G, Buccianti G, Redaelli B and Giangrande A, 1991. Prospective, randomised, multicentre trial of effect of protein restriction on progression of chronic renal insufficiency. Northern Italian Cooperative Study Group. *Lancet*, 337, 1299-1304.
- Lopes C, Oliveira A, Santos AC, Ramos E, Gaio AR, Severo M and Barros H, 2006. Consumo alimentar no Porto. Faculdade de Medecina da Universidade do Porto, 172 pp.
- Macotela Y, Emanuelli B, Bang AM, Espinoza DO, Boucher J, Beebe K, Gall W and Kahn CR, 2011. Dietary leucine--an environmental modifier of insulin resistance acting on multiple levels of metabolism. *PLoS ONE*, 6, e21187.
- Manios Y, 2006. Design and descriptive results of the "Growth, Exercise and Nutrition Epidemiological Study In preSchoolers": the GENESIS study. *BMC Public Health*, 6, 32.
- Maroni BJ and Mitch WE, 1997. Role of nutrition in prevention of the progression of renal disease. *Annual Review of Nutrition*, 17, 435-455.
- Masanés R, Fernández-López J-A, Alemany M, Remesar X and Rafecas I, 1999. Effect of dietary protein content on tissue protein synthesis rates in Zucker lean rats. *Nutrition Research*, 19, 1017-1026.

- Matthys C, De Henauw S, Devos C and De Backer G, 2003. Estimated energy intake, macronutrient intake and meal pattern of Flemish adolescents. *European Journal of Clinical Nutrition*, 57, 366-375.
- McNurlan MA, Fern EB and Garlick PJ, 1982. Failure of leucine to stimulate protein synthesis in vivo. *Biochemical Journal*, 204, 831-838.
- Mensink GBM, Heseker H, Richter A, Stahl A and Vohmann C, 2007. Forschungsbericht: Ernährungsstudie als KiGGS-Modul (EsKiMo) [Research report: nutritional study as KiGGS module]. 143 pp.
- Merten C, Ferrari P, Bakker M, Boss A, Hearty A, Leclercq C, Lindtner O, Tlustos C, Verger P, Volatier JL and Arcella D, 2011. Methodological characteristics of the national dietary surveys carried out in the European Union as included in the European Food Safety Authority (EFSA) Comprehensive European Food Consumption Database. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment*, 28, 975-995.
- Meyer HE, Pedersen JI, Loken EB and Tverdal A, 1997. Dietary factors and the incidence of hip fracture in middle-aged Norwegians. A prospective study. *American Journal of Epidemiology*, 145, 117-123.
- Michaelsen KF, Hoppe C, Roos N, Kaestel P, Stougaard M, Lauritzen L, Molgaard C, Girma T and Friis H, 2009. Choice of foods and ingredients for moderately malnourished children 6 months to 5 years of age. *Food and Nutrition Bulletin*, 30, S343-404.
- Millward DJ, Fereday A, Gibson NR and Pacy PJ, 2000. Human adult amino acid requirements: [1-<sup>13</sup>C]leucine balance evaluation of the efficiency of utilization and apparent requirements for wheat protein and lysine compared with those for milk protein in healthy adults. *American Journal of Clinical Nutrition*, 72, 112-121.
- Mitch WE and Clark AS, 1984. Specificity of the effects of leucine and its metabolites on protein degradation in skeletal muscle. *Biochemical Journal*, 222, 579-586.
- Mojtahedi M, de Groot LC, Boekholt HA and van Raaij JM, 2002. Nitrogen balance of healthy Dutch women before and during pregnancy. *American Journal of Clinical Nutrition*, 75, 1078-1083.
- Mordier S, Deval C, Bechet D, Tassa A and Ferrara M, 2000. Leucine limitation induces autophagy and activation of lysosome-dependent proteolysis in C2C12 myotubes through a mammalian target of rapamycin-independent signaling pathway. *Journal of Biological Chemistry*, 275, 29900-29906.
- Moreira P, Santos S, Padrao P, Cordeiro T, Bessa M, Valente H, Barros R, Teixeira V, Mitchell V, Lopes C and Moreira A, 2010. Food patterns according to sociodemographics, physical activity, sleeping and obesity in Portuguese children. *International Journal of Environmental Research and Public Health*, 7, 1121-1138.
- Morens C, Gaudichon C, Fromentin G, Marsset-Baglieri A, Bensaid A, Larue-Achagiotis C, Luengo C and Tome D, 2001. Daily delivery of dietary nitrogen to the periphery is stable in rats adapted to increased protein intake. *American Journal of Physiology, Endocrinology and Metabolism*, 281, E826-836.
- Morens C, Bos C, Pueyo ME, Benamouzig R, Gausseres N, Luengo C, Tome D and Gaudichon C, 2003. Increasing habitual protein intake accentuates differences in postprandial dietary nitrogen utilization between protein sources in humans. *Journal of Nutrition*, 133, 2733-2740.
- Munger RG, Cerhan JR and Chiu BC, 1999. Prospective study of dietary protein intake and risk of hip fracture in postmenopausal women. *American Journal of Clinical Nutrition*, 69, 147-152.
- Mussolino ME, Looker AC, Madans JH, Langlois JA and Orwoll ES, 1998. Risk factors for hip fracture in white men: the NHANES I Epidemiologic Follow-up Study. *Journal of Bone and Mineral Research*, 13, 918-924.
- Nagasawa T, Kido T, Yoshizawa F, Ito Y and Nishizawa N, 2002. Rapid suppression of protein degradation in skeletal muscle after oral feeding of leucine in rats. *Journal of Nutritional Biochemistry*, 13, 121-127.
- Nair KS, Matthews DE, Welle SL and Braiman T, 1992. Effect of leucine on amino acid and glucose metabolism in humans. *Metabolism: Clinical and Experimental*, 41, 643-648.

- Nair KS and Short KR, 2005. Hormonal and signaling role of branched-chain amino acids. *Journal of Nutrition*, 135, 1547S-1552S.
- Newgard CB, An J, Bain JR, Muehlbauer MJ, Stevens RD, Lien LF, Haqq AM, Shah SH, Arlotto M, Slentz CA, Rochon J, Gallup D, Ilkayeva O, Wenner BR, Yancy WS, Jr., Eisensohn H, Musante G, Surwit RS, Millington DS, Butler MD and Svetkey LP, 2009. A branched-chain amino acid-related metabolic signature that differentiates obese and lean humans and contributes to insulin resistance. *Cell Metabolism*, 9, 311-326.
- NNR (Nordic Nutrition Recommendations), 2004. Integrating nutrition and physical activity. Nordic Council of Ministers, Copenhagen, Denmark, 436 pp.
- Noatsch A, Petzke KJ, Millrose MK and Klaus S, 2010. Body weight and energy homeostasis was not affected in C57BL/6 mice fed high whey protein or leucine-supplemented low-fat diets. *European Journal of Nutrition*, 50, 479-488.
- Nygren J and Nair KS, 2003. Differential regulation of protein dynamics in splanchnic and skeletal muscle beds by insulin and amino acids in healthy human subjects. *Diabetes*, 52, 1377-1385.
- Ocke MC, van Rossum CTM, Fransen HP, Buurma EJM, de Boer EJ, Brants HAM, Niekerk EM, van der Laan JD, Drijvers JJMM and Ghameshlou Z, 2008. Dutch National Food Consumption Survey - Young Children 2005/2006. RIVM Report 350070001/2008, National Institute for Public Health and the Environment, 105 pp.
- Øverby NC and Andersen LF, 2002. Ungkost, 2000. Landsomfattende kostholdundersøkelse blant elever i 4.- og 8. klasse i Norge [Nationwide nutrition survey among students in 4 - and 8th class in Norway]. Sosial – og helsedirektoratet, avdeling for ernærings, 40 pp.
- Pacy PJ, Price GM, Halliday D, Quevedo MR and Millward DJ, 1994. Nitrogen homeostasis in man: the diurnal responses of protein synthesis and degradation and amino acid oxidation to diets with increasing protein intakes. *Clinical Science (London)*, 86, 103-116.
- Paddon-Jones D, Sheffield-Moore M, Zhang XJ, Volpi E, Wolf SE, Aarsland A, Ferrando AA and Wolfe RR, 2004. Amino acid ingestion improves muscle protein synthesis in the young and elderly. *American Journal of Physiology. Endocrinology and Metabolism*, 286, E321-328.
- Paddon-Jones D and Rasmussen BB, 2009. Dietary protein recommendations and the prevention of sarcopenia. *Current Opinion in Clinical Nutrition and Metabolic Care*, 12, 86-90.
- Pannemans DL, Halliday D and Westerterp KR, 1995a. Whole-body protein turnover in elderly men and women: responses to two protein intakes. *American Journal of Clinical Nutrition*, 61, 33-38.
- Pannemans DL, Halliday D, Westerterp KR and Kester AD, 1995b. Effect of variable protein intake on whole-body protein turnover in young men and women. *American Journal of Clinical Nutrition*, 61, 69-74.
- Pannemans DL, Wagenmakers AJ, Westerterp KR, Schaafsma G and Halliday D, 1998. Effect of protein source and quantity on protein metabolism in elderly women. *American Journal of Clinical Nutrition*, 68, 1228-1235.
- Pannemans DLE, Wagenmakers AJM, Westerterp KR, Schaafsma G and Halliday D, 1997. The effect of an increase of protein intake on whole body protein turnover in elderly women is tracer dependent. *Journal of Nutrition*, 127, 1788-1794.
- Patti ME, Brambilla E, Luzi L, Landaker EJ and Kahn CR, 1998. Bidirectional modulation of insulin action by amino acids. *Journal of Clinical Investigation*, 101, 1519-1529.
- Paturi M, Tapanainen H, Reinivuo H and Pietinen P, 2008. The National FINDiet 2007 Survey. Report B23/2008, KTL-National Public Health Institute, 230 pp.
- Pedersen AN, Fagt S, Groth MV, Christensen T, Biltoft-Jensen A, Matthiessen J, Lyhne Andersen N, Kørup K, Hartkopp H, Hess Ygil K, Hinsch HJ, Saxholt E and Trolle E, 2010. Danskernes kostvaner 2003-2008. Hovedresultater [Dietary habits in Denmark 2003-2008. Main results]. DTU Fødevarerinstitutionen, 200 pp.

- Pellett PL and Young VR, 1980. Nutritional evaluation of protein foods. United Nations University Press, Tokyo, Japan, 167 pp.
- Pencharz PB and Ball RO, 2003. Different approaches to define individual amino acid requirements. *Annual Review of Nutrition*, 23, 101-116.
- Piatti PM, Monti F, Fermo I, Baruffaldi L, Nasser R, Santambrogio G, Librenti MC, Galli-Kienle M, Pontiroli AE and Pozza G, 1994. Hypocaloric high-protein diet improves glucose oxidation and spares lean body mass: comparison to hypocaloric high-carbohydrate diet. *Metabolism: Clinical and Experimental*, 43, 1481-1487.
- Pietinen P, Paturi M, Reinivuo H, Tapanainen H and Valsta LM, 2010. FINDIET 2007 Survey: energy and nutrient intakes. *Public Health Nutrition*, 13, 920-924.
- Pipe NG, Smith T, Halliday D, Edmonds CJ, Williams C and Coltart TM, 1979. Changes in fat, fat-free mass and body water in human normal pregnancy. *British Journal of Obstetrics and Gynaecology*, 86, 929-940.
- Pomerleau J, McKee M, Robertson A, Kadziauskiene K, Abaravicius A, Vaask S, Pudule I and Grinberga D, 2001. Macronutrient and food intake in the Baltic republics. *European Journal of Clinical Nutrition*, 55, 200-207.
- Price GM, Halliday D, Pacy PJ, Quevedo MR and Millward DJ, 1994. Nitrogen homeostasis in man: influence of protein intake on the amplitude of diurnal cycling of body nitrogen. *Clinical Science (London)*, 86, 91-102.
- Promislow JH, Goodman-Gruen D, Slymen DJ and Barrett-Connor E, 2002. Protein consumption and bone mineral density in the elderly : the Rancho Bernardo Study. *American Journal of Epidemiology*, 155, 636-644.
- Rand WM and Young VR, 1999. Statistical analysis of nitrogen balance data with reference to the lysine requirement in adults. *Journal of Nutrition*, 129, 1920-1926.
- Rand WM, Pellett PL and Young VR, 2003. Meta-analysis of nitrogen balance studies for estimating protein requirements in healthy adults. *American Journal of Clinical Nutrition*, 77, 109-127.
- Renteria-Flores JA, Johnston LJ, Shurson GC and Gallaher DD, 2008. Effect of soluble and insoluble fiber on energy digestibility, nitrogen retention, and fiber digestibility of diets fed to gestating sows. *Journal of Animal Science*, 86, 2568-2575.
- Rieu I, Sornet C, Bayle G, Prugnaud J, Pouyet C, Balage M, Papet I, Grizard J and Dardevet D, 2003. Leucine-supplemented meal feeding for ten days beneficially affects postprandial muscle protein synthesis in old rats. *Journal of Nutrition*, 133, 1198-1205.
- Rodler I, Bíró L, Greiner E, Zajkás G, Szórád I, Varga A, Domonkos A, Ágoston H, Balázs A, Mozsáry E, Vitrai J, Hermann D, Boros J, Németh R and Kéki Z, 2005. Táplálkozási vizsgálat Magyarországon, 2003–2004. Energia- és makrotápanyagbevitel [Dietary survey in Hungary, 2003–2004. Energy and macro-nutrient intake]. *Orvosi Hetilap [Hungarian Medical Journal]*, 146, 1781–1789.
- Rose WC, 1957. The amino acid requirements of adult man. *Nutrition Abstracts and Reviews, Series A: Human and Experimental*, 27, 631-647.
- Samaha FF, Iqbal N, Seshadri P, Chicano KL, Daily DA, McGrory J, Williams T, Williams M, Gracely EJ and Stern L, 2003. A low-carbohydrate as compared with a low-fat diet in severe obesity. *New England Journal of Medicine*, 348, 2074-2081.
- Satterlee LD, Marshall HF and Tennyson JM, 1979. Measuring protein quality. *Journal of the American Oil Chemists Society*, 56, 103-109.
- SCF (Scientific Committee for Food), 1993. Report on nutrient and energy intakes for the European Community, 31st Series. Food - Science and Techniques, European Commission, 255 pp.
- SCF (Scientific Committee on Food), 2003. Report on the Revision of Essential Requirements of Infant Formulae and Follow-on Formulae. SCF/CS/NUT/IF/65 Final, 213 pp.

- Schoknecht PA and Pond WG, 1993. Short-term ingestion of a high protein diet increases liver and kidney mass and protein accretion but not cellularity in young pigs. *Proceedings of the Society for Experimental Biology and Medicine*, 203, 251-254.
- Schwenk WF and Haymond MW, 1987. Effects of leucine, isoleucine, or threonine infusion on leucine metabolism in humans. *American Journal of Physiology*, 253, E428-434.
- Serra-Majem L, Ribas-Barba L, Salvador G, Jover L, Raido B, Ngo J and Plasencia A, 2007. Trends in energy and nutrient intake and risk of inadequate intakes in Catalonia, Spain (1992-2003). *Public Health Nutrition*, 10, 1354-1367.
- Sette S, Le Donne C, Piccinelli R, Arcella D, Turrini A and Leclercq C, 2010. The third Italian National Food Consumption Survey, INRAN-SCAI 2005-06 - Part 1: Nutrient intakes in Italy. *Nutrition, Metabolism and Cardiovascular Diseases*, 1-11.
- Shah SH, Crosslin DR, Haynes CS, Nelson S, Turer CB, Stevens RD, Muehlbauer MJ, Wenner BR, Bain JR, Laferrere B, Gorroochurn P, Teixeira J, Brantley PJ, Stevens VJ, Hollis JF, Appel LJ, Lien LF, Batch B, Newgard CB and Svetkey LP, 2012. Branched-chain amino acid levels are associated with improvement in insulin resistance with weight loss. *Diabetologia*, 55, 321-330.
- Sharman MJ, Kraemer WJ, Love DM, Avery NG, Gomez AL, Scheett TP and Volek JS, 2002. A ketogenic diet favorably affects serum biomarkers for cardiovascular disease in normal-weight men. *Journal of Nutrition*, 132, 1879-1885.
- Sherwin RS, 1978. Effect of starvation on the turnover and metabolic response to leucine. *Journal of Clinical Investigation*, 61, 1471-1481.
- Skov AR, Toubro S, Bulow J, Krabbe K, Parving HH and Astrup A, 1999a. Changes in renal function during weight loss induced by high vs low-protein low-fat diets in overweight subjects. *International Journal of Obesity and Related Metabolic Disorders*, 23, 1170-1177.
- Skov AR, Toubro S, Ronn B, Holm L and Astrup A, 1999b. Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *International Journal of Obesity and Related Metabolic Disorders*, 23, 528-536.
- Skov AR, Haulrik N, Toubro S, Molgaard C and Astrup A, 2002. Effect of protein intake on bone mineralization during weight loss: a 6-month trial. *Obesity Research*, 10, 432-438.
- Symons TB, Schutzler SE, Cocke TL, Chinkes DL, Wolfe RR and Paddon-Jones D, 2007. Aging does not impair the anabolic response to a protein-rich meal. *American Journal of Clinical Nutrition*, 86, 451-456.
- Szponar L, Sekuła W, Rychlik E, Oltarzewski M and Figurska K, 2003. *Badania indywidualnego spożycia żywności i stanu odżywienia w gospodarstwach domowych [Household food consumption and anthropometric survey]*. Instytut Żywności i Żywienia, Warsaw, Poland, 900 pp.
- Taillandier D, Aourousseau E, Combaret L, Guezennec CY and Attaix D, 2003. Regulation of proteolysis during reloading of the unweighted soleus muscle. *International Journal of Biochemistry and Cell Biology*, 35, 665-675.
- Teegarden D, Lyle RM, McCabe GP, McCabe LD, Proulx WR, Michon K, Knight AP, Johnston CC and Weaver CM, 1998. Dietary calcium, protein, and phosphorus are related to bone mineral density and content in young women. *American Journal of Clinical Nutrition*, 68, 749-754.
- Tessari P, Tsalikian E, Schwenk WF, Nissen SL and Haymond MW, 1985. Effects of [<sup>15</sup>N]leucine infused at low rates on leucine metabolism in humans. *American Journal of Physiology*, 249, E121-130.
- Thalacker-Mercer AE, Fleet JC, Craig BA and Campbell WW, 2010. The skeletal muscle transcript profile reflects accommodative responses to inadequate protein intake in younger and older males. *Journal of Nutritional Biochemistry*, 21, 1076-1082.
- Theil PK, Jorgensen H and Jakobsen K, 2002. Energy and protein metabolism in pregnant sows fed two levels of dietary protein. *Journal of Animal Physiology and Animal Nutrition*, 86, 399-413.

- Tian Y, Liu J, Zhang Y, Piao J, Gou L, Li M, Ji Y and Yang X, 2011. Examination of Chinese habitual dietary protein requirements of Chinese young female adults by indicator amino acid method. *Asia Pacific Journal of Clinical Nutrition*, 20, 390-396.
- Tischler ME, Desautels M and Goldberg AL, 1982. Does leucine, leucyl-tRNA, or some metabolite of leucine regulate protein synthesis and degradation in skeletal and cardiac muscle? *Journal of Biological Chemistry*, 257, 1613-1621.
- Tome D and Bos C, 2000. Dietary protein and nitrogen utilization. *Journal of Nutrition*, 130, 1868S-1873S.
- Tremblay F and Marette A, 2001. Amino acid and insulin signaling via the mTOR/p70 S6 kinase pathway. A negative feedback mechanism leading to insulin resistance in skeletal muscle cells. *Journal of Biological Chemistry*, 276, 38052-38060.
- Tsunehara CH, Leonetti DL and Fujimoto WY, 1990. Diet of second-generation Japanese-American men with and without non-insulin-dependent diabetes. *American Journal of Clinical Nutrition*, 52, 731-738.
- Tucker KL, Hannan MT and Kiel DP, 2001. The acid-base hypothesis: diet and bone in the Framingham Osteoporosis Study. *European Journal of Nutrition*, 40, 231-237.
- USDA/ARS (United States Department of Agriculture Agricultural Research Service), 2009. USDA National Nutrient Database for Standard Reference, Release 22. Nutrient Data Laboratory Home Page. Available from: <http://www.ars.usda.gov/ba/bhnrc/ndl>, accessed 19 January 2012.
- USDA/HHS (U.S. Department of Agriculture and U.S. Department of Health and Human Services), 2010. *Dietary Guidelines for Americans, 2010*. 112 pp.
- van Buuren S, Schönbeck Y and van Dommelen P, 2012. CT/EFSA/NDA/2010/01: Collection, collation and analysis of data in relation to reference heights and reference weights for female and male children and adolescents (0-18 years) in the EU, as well as in relation to the age of onset of puberty and the age at which different stages of puberty are reached in adolescents in the EU. Draft Technical Report submitted to EFSA, 57 pp.
- van Rossum CTM, Fransen HP, Verkaik-Kloosterman J, Buurma-Rethans EJM and Ocké MC, 2011. Dutch National Food Consumption Survey 2007-2010: Diet of children and adults aged 7 to 69 years. RIVM Report number: 350050006/2011, National Institute for Public Health and the Environment, 148 pp.
- Volek JS, Sharman MJ, Love DM, Avery NG, Gomez AL, Scheett TP and Kraemer WJ, 2002. Body composition and hormonal responses to a carbohydrate-restricted diet. *Metabolism: Clinical and Experimental*, 51, 864-870.
- Volpi E, Lucidi P, Cruciani G, Monacchia F, Reboldi G, Brunetti P, Bolli GB and De Feo P, 1996. Contribution of amino acids and insulin to protein anabolism during meal absorption. *Diabetes*, 45, 1245-1252.
- Wang J, Alexander JT, Zheng P, Yu HJ, Dourmashkin J and Leibowitz SF, 1998. Behavioral and endocrine traits of obesity-prone and obesity-resistant rats on macronutrient diets. *American Journal of Physiology*, 274, E1057-1066.
- Wang TJ, Larson MG, Vasan RS, Cheng S, Rhee EP, McCabe E, Lewis GD, Fox CS, Jacques PF, Fernandez C, O'Donnell CJ, Carr SA, Mootha VK, Florez JC, Souza A, Melander O, Clish CB and Gerszten RE, 2011. Metabolite profiles and the risk of developing diabetes. *Nature Medicine*, 17, 448-453.
- Waterlow JC, 1995. Whole-body protein turnover in humans--past, present, and future. *Annual Review of Nutrition*, 15, 57-92.
- Waterlow JC, 1996. The requirements of adult man for indispensable amino acids. *European Journal of Clinical Nutrition*, 50 Suppl 1, S151-176.
- Weigle DS, Breen PA, Matthys CC, Callahan HS, Meeuws KE, Burden VR and Purnell JQ, 2005. A high-protein diet induces sustained reductions in appetite, ad libitum caloric intake, and body weight despite compensatory changes in diurnal plasma leptin and ghrelin concentrations. *American Journal of Clinical Nutrition*, 82, 41-48.

- Westerterp-Plantenga MS, Lejeune MP, Nijs I, van Ooijen M and Kovacs EM, 2004. High protein intake sustains weight maintenance after body weight loss in humans. *International Journal of Obesity and Related Metabolic Disorders*, 28, 57-64.
- WHO/FAO/UNU (World Health Organization/Food and Agriculture Organization of the United Nations/United Nations University), 2007. Protein and amino acid requirements in human nutrition. Report of a Joint WHO/FAO/UNU Expert Consultation. WHO Technical Report Series, No 935, 284 pp.
- Widdowson EM, Southgate DAT and Hey EN, 1979. Body composition of the fetus and infant. In: *Nutrition of the fetus and infant*. Ed Visser H. Martinus Nijhoff Publishers, London, UK, 169-177.
- Wiegmann TB, Zlomke AM, MacDougall ML and Kipp DE, 1990. Controlled changes in chronic dietary protein intake do not change glomerular filtration rate. *American Journal of Kidney Diseases*, 15, 147-154.
- Wolfe RR, Miller SL and Miller KB, 2008. Optimal protein intake in the elderly. *Clinical Nutrition*, 27, 675-684.
- Yancy WS, Jr., Olsen MK, Guyton JR, Bakst RP and Westman EC, 2004. A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and hyperlipidemia: a randomized, controlled trial. *Annals of Internal Medicine*, 140, 769-777.
- Young VR and Marchini JS, 1990. Mechanisms and nutritional significance of metabolic responses to altered intakes of protein and amino acids, with reference to nutritional adaptation in humans. *American Journal of Clinical Nutrition*, 51, 270-289.
- Young VR and Borgonha S, 2000. Nitrogen and amino acid requirements: the Massachusetts Institute of Technology amino acid requirement pattern. *Journal of Nutrition*, 130, 1841S-1849S.
- Zeller KR, 1991. Low-protein diets in renal disease. *Diabetes Care*, 14, 856-866.
- Zernicke RF, Salem GJ, Barnard RJ, Woodward JS, Jr., Meduski JW and Meduski JD, 1995. Adaptations of immature trabecular bone to exercise and augmented dietary protein. *Medicine and Science in Sports and Exercise*, 27, 1486-1493.



## APPENDICES

### APPENDIX 1: MAIN FOOD CONTRIBUTORS TO DIETARY PROTEIN INTAKE (%) OF ADULTS (18-64 YEARS) IN EUROPEAN COUNTRIES AS ESTIMATED WITH THE EFSA COMPREHENSIVE EUROPEAN FOOD CONSUMPTION DATABASE

(EFSA, 2011b; Merten et al., 2011)\*

	Meat and meat products (including edible offal)	Milk and dairy products	Fish and other seafood	Eggs and egg products	Grains and grain-based products	Legumes, nuts and oilseeds	Starchy roots and tubers	Vegetables and vegetable products (including fungi)	Fruit and fruit products	Composite food (including frozen products)	Snacks, desserts, and other foods (including amphibians, reptiles, snails and insects)	Herbs, spices and condiments	Sugar and confectionary	Non-alcoholic beverages (including milk-based beverages)	Fruit and vegetable juices	Alcoholic beverages
<b>Austria</b>	40	19	4	3	19	2	1	3	2	0	1	1	1	1	1	1
<b>Belgium</b>	31	18	6	2	26	1	3	2	1	5	1	1	1	1	0	1
<b>Bulgaria</b>	30	15	6	4	30	5	3	4	1	0	0	0	0	0	0	0
<b>Czech Republic</b>	35	15	3	3	28	2	2	4	1	2	1	1	0	0	0	2
<b>Denmark</b>	32	25	4	3	23	1	3	2	1	0	1	0	1	1	0	1
<b>Estonia</b>	37	19	6	5	18	2	5	3	1	1	1	1	0	1	1	1
<b>Finland</b>	30	28	6	3	19	2	2	2	1	0	0	1	1	2	0	1
<b>France</b>	39	19	7	2	20	2	2	2	1	0	1	1	1	1	0	0
<b>Hungary</b>	37	15	2	4	29	4	3	3	1	0	0	1	0	0	0	0
<b>Ireland</b>	42	15	4	2	19	1	6	2	1	3	1	0	1	1	0	1
<b>Italy</b>	28	20	9	3	28	2	1	4	1	1	1	1	0	1	0	0
<b>Latvia</b>	39	10	4	2	18	1	4	1	1	17 #	0	1	1	1	0	0
<b>The Netherlands</b>	30	23	2	1	21	1	3	2	1	7	2	1	1	1	1	1
<b>Poland</b>	42	13	3	4	23	1	5	4	1	0	0	0	0	1	0	0
<b>Spain §</b>	32	19	13	3	17	4	2	4	1	2	1	0	0	0	0	1
<b>United Kingdom</b>	33	18	7	3	22	3	4	2	1	0	1	2	1	1	0	1

\* Figures may not add up to 100 % due to rounding.

# The use of items from the “Composite food (including frozen products)” category in FoodEx (the food classification system applied to the development of the EFSA Comprehensive European Food Consumption Database) was discouraged. Most countries managed to split the majority of their composite foods into their ingredients, but in Latvia 10 % of the composite foods or home-made dishes were not broken down into their ingredients (EFSA, 2011a). §Results for Spain II (dietary survey acronym: AESAN)

## APPENDIX 2A: POPULATION, METHODS AND PERIOD OF DIETARY ASSESSMENT IN CHILDREN AND ADOLESCENTS IN EUROPEAN COUNTRIES

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Austria</b>	Boys and girls aged 7-9 years	3-day record	2007-2008	(Elmadfa et al., 2009a; Elmadfa et al., 2009b)
	Boys and girls aged 10-14 years	3-day record	2007-2008	(Elmadfa et al., 2009a; Elmadfa et al., 2009b)
	Boys and girls aged 14-19 years	24-hour recall	2003-2004	(Elmadfa et al., 2009a; Elmadfa et al., 2009b). <i>Mainly from a large Viennese sample.</i>
<b>Belgium</b>	Boys and girls aged 2.5-3 years	3-day record	2002-2003	(Huybrechts and De Henauw, 2007). <i>Data collected in Flanders.</i>
	Boys and girls aged 4-6.5 years	3-day record	2002-2003	(Huybrechts and De Henauw, 2007). <i>Data collected in Flanders.</i>
	Boys and girls aged 13-15 years	7-day record	1997	(Matthys et al., 2003). <i>Data collected in the region of Ghent in Flanders.</i>
	Boys and girls aged 15-18 years	2 x 24-hour recall	2004	(De Vriese et al., 2006)
<b>Bulgaria</b>	Boys and girls aged 1-3 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Boys and girls aged 3-6 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Boys and girls aged 6-10 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Boys and girls aged 10-14 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Boys and girls aged 14-18 years	24-hour recall	1998	(Abrasheva et al., 1998)
<b>Czech Republic</b>	Boys and girls aged 4-6 years	48-hour recall	2007	(Elmadfa et al., 2009a)
	Boys and girls aged 7-9 years	48-hour recall	2007	(Elmadfa et al., 2009a)
<b>Denmark</b>	Boys and girls aged 1-3 years	7-day record	1995	(Andersen et al., 1996)
	Boys and girls aged 4-5 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Boys and girls aged 6-9 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Boys and girls aged 10-13 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Boys and girls aged 14-17 years	7-day record	2003-2008	(Pedersen et al., 2010)
<b>Finland</b>	Children aged 1 year	3-day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
	Children aged 2 years	3-day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
	Children aged 3 years	3 day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
	Children aged 4 years	3-day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
	Children aged 6 years	3-day record	2003-2005	(Kyttälä et al., 2008; Kyttälä et al., 2010)
<b>France</b>	Boys and girls aged 4-6 years	3 x 24-hour recall	2006-2007	(Elmadfa et al., 2009a)
	Boys and girls aged 7-9 years	3 x 24-hour recall	2006-2007	(Elmadfa et al., 2009a)
	Boys and girls aged 10-14 years	3 x 24-hour recall	2006-2007	(Elmadfa et al., 2009a)
	Boys and girls aged 15-18 years	3 x 24-hour recall	2006-2007	(Elmadfa et al., 2009a)
<b>Germany</b>	Infants aged 12 months	3-day record	1989-2003	(Hilbig and Kersting, 2006).
	Children aged 18 months	3-day record	1989-2003	(Hilbig and Kersting, 2006).
	Children aged 2 years	3-day record	1989-2003	(Hilbig and Kersting, 2006).
	Children aged 3 years	3-day record	1989-2003	(Hilbig and Kersting, 2006).
	Boys and girls aged 6 years	3-day record	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 7-9 years	3-day record	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 10-11 years	3-day record	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 12 years	Dietary history (over the last 4 weeks)	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 13-14 years	Dietary history (over the last 4 weeks)	2006	(Elmadfa et al., 2009a; Mensink et al., 2007)
	Boys and girls aged 15-17 years	Dietary history (over the last 4 weeks)	2006	(Mensink et al., 2007)

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Greece</b>	Boys and girls aged 12-24 mo	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006)
	Boys and girls aged 25-36 mo	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006)
	Boys and girls aged 37-48 mo	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006)
	Boys and girls aged 49-60 mo	3-day record (weighed food records and 24-hour recall or food diaries)	2003-2004	(Manios, 2006)
<b>Hungary</b>	Boys and girls aged 11-14 years	3-day record	2005-2006	(Biro et al., 2007). <i>Data collected in Budapest.</i>
<b>Ireland</b>	Boys and girls aged 5-8 years	7-day record	2003-2004	(IUNA (Irish Universities Nutrition Alliance), a)
	Boys and girls aged 9-12 years	7-day record	2003-2004	(IUNA (Irish Universities Nutrition Alliance), a)
	Boys and girls aged 13-14 years	7-day record	2005-2006	(IUNA (Irish Universities Nutrition Alliance), b)
	Boys and girls aged 15-17 years	7-day record	2005-2006	(IUNA (Irish Universities Nutrition Alliance), b)
<b>Italy</b>	Boys and girls aged 0-<3 years	consecutive 3-day food record	2005-2006	(Sette et al., 2010)
	Boys and girls aged 3-<10 years	consecutive 3-day food record	2005-2006	(Sette et al., 2010)
	Boys and girls aged 10-<18 years	consecutive 3-day food record	2005-2006	(Sette et al., 2010)
<b>The Netherlands</b>	Infants aged 9 months	2-day record (independent days)	2002	(de Boer et al., 2006)
	Infants aged 12 months	2-day record (independent days)	2002	(de Boer et al., 2006)
	Children aged 18 months	2-day record (independent days)	2002	(de Boer et al., 2006)
	Boys and girls aged 2-3 years	2-day record (independent days)	2005-2006	(Ocke et al., 2008)
	Boys and girls aged 4-6 years	2-day record (independent days)	2005-2006	(Ocke et al., 2008)
	Boys and girls aged 7-8 years	2 non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)
	Boys and girls aged 9-13 years	2 non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)
	Boys and girls aged 14-18 years	2 non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)
<b>Norway</b>	Children aged 2 years	Food Frequency Questionnaire	2007	(Kristiansen and Andersen, 2009)
	Boys and girls aged 4 years	4-day record	2000	(Elmadfa et al., 2009a; Øverby and Andersen, 2002)
	Boys and girls aged 9 years	4-day record	2000	(Elmadfa et al., 2009a; Øverby and Andersen, 2002)
	Boys and girls aged 13 years	4-day record	2000	(Elmadfa et al., 2009a; Øverby and Andersen, 2002)
	Boys and girls aged 16-19 years	Food Frequency Questionnaire	1997	(Johansson and Sovoll, 1999)
<b>Poland</b>	Boys and girls aged 1-3 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 4-6 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 7-9 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 10-12 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 13-15 years	24-hour recall	2000	(Szponar et al., 2003)
	Boys and girls aged 16-18 years	24-hour recall	2000	(Szponar et al., 2003)
<b>Portugal</b>	Boys and girls aged 5-10 years	Food Frequency Questionnaire	2006-2007	(Moreira et al., 2010)
<b>Slovenia</b>	Boys and girls aged 14-16 years	Food Frequency Questionnaire	2003-2005	(Kobe et al., 2011)
<b>Spain</b>	Boys and girls aged 10-14 years	2 non-consecutive 24-hour dietary recalls	2002-2003	(Elmadfa et al., 2009a). <i>Data collected in Catalonia.</i>
	Boys and girls aged 15-18 years	2 non-consecutive 24-hour dietary recalls	2002-2003	(Elmadfa et al., 2009a). <i>Data collected in Catalonia.</i>

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Sweden</b>	Boys and girls aged 4 years	4-day record	2003	(Enghardt-Barbieri et al., 2006)
	Boys and girls aged 8-9 years	4-day record	2003	(Enghardt-Barbieri et al., 2006)
	Boys and girls aged 11-12 years	4-day record	2003	(Enghardt-Barbieri et al., 2006)
<b>United Kingdom</b>	Boys and girls aged 1.5-3 years	4-day food diary	2008-2010	(Bates et al., 2011)
	Boys and girls aged 4-10 years	4-day food diary	2008-2010	(Bates et al., 2011)
	Boys and girls aged 11-18 years	4-day food diary	2008-2010	(Bates et al., 2011)

mo: months

**APPENDIX 2B: PROTEIN INTAKE OF CHILDREN AGED ~1-3 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	N	Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
			mean	SD	P5 - P95	mean	SD	P5 - P95	mean	SD	P5 - P95
<b>Infants and/or young children (both sexes)</b>											
<b>Bulgaria</b>	1-3 years	154	11.7	2.2		39.4	14.8		3.03	1.19	
<b>Germany</b>	12 mo	432 <sup>1,2</sup>	13.2	2.2							
	18 mo	478 <sup>1</sup>	13.9	2.1							
	2 years	458 <sup>1</sup>	13.6	2.2							
	3 years	427 <sup>1</sup>	12.9	2.0							
<b>Italy</b>	0-<3 years	52	14.7	4.4	5.7-21.6	41.5	18.0	7.7-71.3	3.64	1.24	1.46-5.58
<b>The Netherlands</b>	9 mo	333	11.8	1.4	10.2-13.7 <sup>3</sup>	28.8	6.2	21.4-27.0 <sup>3</sup>			
	12 mo	306	13.7	2.5	10.8-17.0 <sup>3</sup>	36.5	8.3	26.8-47.6 <sup>3</sup>			
	18 mo	302	15.0	2.1	12.4-17.7 <sup>3</sup>	43.1	6.5	34.9-51.5 <sup>3</sup>			
<b>United Kingdom</b>	1.5-3 years	219	15.3	2.4	11.5-20.9 <sup>4</sup>	42.6	11.1	21.2-63.8 <sup>4</sup>			
<b>Young children</b>											
<b>Males</b>											
<b>Belgium</b>	2.5-3	102	16.2	2.4		62.5	11.3				
<b>Denmark</b>	1-3	129	13		10-16	52		36-74			
<b>Finland</b>	1 <sup>2</sup>	257	15			35	11				
	2	112	16			43	12				
	3	236	16			49	12				
<b>Greece</b>	12-24 mo	100	16.3	1.8		52.2	10.7				
	25-36 mo	274	16.6	2.1		57.8	11.7				
	37-48 mo	488	16.6	2.2		59.8	12.7				
<b>The Netherlands</b>	2-3	327	13		11-16	44		31-60			
<b>Norway</b>	2	829				50.8	14.9				
<b>Poland</b>	1-3	70	13.3	3.9		46.4	21.3				
<b>Females</b>											
<b>Belgium</b>	2.5-3	95	16.7	1.6		57.7	11.3				
<b>Denmark</b>	1-3	149	14		11-16	54		31-69			
<b>Finland</b>	1 <sup>2</sup>	198	16			34	8				
	2	118	17			44	11				
	3	235	15			46	44				
<b>Greece</b>	12-24 mo	107	16.2	1.7		50.5	9.6				
	25-36 mo	226	16.5	2.3		55.2	12.6				
	37-48 mo	434	16.5	2.1		56.9	12.6				
<b>The Netherlands</b>	2-3	313	13		11-16	43		31-57			
<b>Norway</b>	2	826				48.6	14.9				
<b>Poland</b>	1-3	48	13.3	2.9		41.2	13.4				

<sup>1</sup>Number of 3-day records; <sup>2</sup>Breast-fed infants not included; <sup>3</sup>P10-P90; <sup>4</sup>P2.5-P97.5; mo: months

**APPENDIX 2C: PROTEIN INTAKE OF CHILDREN AGED ~4-6 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	N	Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
			mean	SD	P5 - P95	mean	SD	P5 - P95	mean	SD	P5 - P95
<b>Males</b>											
Belgium	4-6.5	236	15.4	2.2		58.5	10.0				
Czech Republic	4-6	641	14.0	2.2							
Denmark	4-5	81	14	2.0	11-18	63	13	44-85			
Finland	4	307	15			53	13				
	6	364	16			61	13				
France	4-6	164	15.5	0.2							
Germany	6	106	13.3	1.9	10.3-17.1	55.3	10.8	39.5-76.9			
Greece	49-60 mo	356	16.4	2.5		60.5	15.2				
The Netherlands	4-6	327	13		10-16	51		35-70			
Norway	4	206	14.2	2.3		52.4	14.5				
Poland	4-6	82	11.1	2.3		50.9	16.0				
Sweden	4	302	14.4	2.2	10.9-18.1	55	13	35-77			
United Kingdom	4-10	210	14.4	2.1	11.0-19.1 <sup>1</sup>	57.2	13.6	34.7-92.4 <sup>1</sup>			
<b>Females</b>											
Belgium	4-6.5	228	15.1	2.0		52.9	10.5				
Czech Republic	4-6	446	14.0	2.2							
Denmark	4-5	78	14	2.0	12-18	58	14	35-80			
Finland	4	307	15			49	11				
	6	349	15			53	12				
France	4-6	162	15.0	0.2							
Germany	6	102	13.6	2.0	11.0-18.5	50.6	12.4	32.1-68.1			
Greece	49-60 mo	389	16.3	2.3		57.8	13.7				
The Netherlands	4-6	312	13		10-16	46		32-60			
Norway	4	185	14.0	2.2		49.5	11.9				
Poland	4-6	84	12.0	2.8		49.4	18.4				
Sweden	4	288	14.4	2.1	11.3-18.1	51	11	34-71			
United Kingdom	4-10	213	14.3	2.3	10.4-19.5 <sup>1</sup>	53.9	12.6	31.8-81.6 <sup>1</sup>			
<b>Both sexes</b>											
Bulgaria	3-6	199	11.9	2.2		50.5	16.5		2.80	1.11	
Italy	3-<10	193	15.7	2.3	12.5-19.5	74.1	18.5	46.9-109.4	3.05	1.02	1.57-4.73

<sup>1</sup>P2.5-P97.5; mo: months

**APPENDIX 2D: PROTEIN INTAKE OF CHILDREN AGED ~7-9 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	N	Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
			mean	SD	P5 - P95	mean	SD	P5 - P95	mean	SD	P5 - P95
<b>Males</b>											
Austria	7-9	146	14.4	2.7							
Czech Republic	7-9	940	14.5	2.4							
Denmark	6-9	172	14	2.1	10-18	73	19	48-102			
France	7-9	160	14.7	0.2							
Germany	7-9	321	13.5	2.1	10.4-17.4	62.0	14.0	40.6-87.0			
Ireland	5-8	145	13.6	2.0	10.6-17.1	55.3	15.8	33.8-82.8			
The Netherlands	7-8	153	12.9 <sup>1</sup>		9.6-16.6	61 <sup>1</sup>		39-88			
Norway	9	402	14	2		73	21				
Poland	7-9	101	11.7	2.8		62.1	22.7				
Portugal	5-10	985	18.1	2.9							
Sweden	8-9	444	15.4	2.3	11.9-19.6	72	17	48-101			
<b>Females</b>											
Austria	7-9	134	13.5	2.7							
Czech Republic	7-9	765	14.5	2.4							
Denmark	6-9	151	14	2.0	11-17	63	14	43-90			
France	7-9	144	15.0	0.3							
Germany	7-9	308	13.6	2.7	9.5-18.5	55.5	14.9	35.6-81.3			
Ireland	5-8	151	13.7	2.1	10.3-17.1	51.9	12.8	34.7-73.0			
The Netherlands	7-8	151	12.4 <sup>1</sup>		9.8-16.8	60 <sup>1</sup>		39-85			
Norway	9	408	14	3		63	20				
Poland	7-9	103	11.3	2.5		53.1	18.7				
Portugal	5-10	991	18.5	3.5							
Sweden	8-9	445	15.4	2.2	12.1-19.2	65	15	43-92			
<b>Both sexes</b>											
Bulgaria	6-10	235	12.4	2.3		68.2	22.1		2.54	0.96	

<sup>1</sup>Median (mean not available)

**APPENDIX 2E: PROTEIN INTAKE OF CHILDREN AGED ~10-14 YEARS AND OVER IN EUROPEAN COUNTRIES**

Country	Age (years)	N	Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
			mean	SD	P5 - P95	mean	SD	P5 - P95	mean	SD	P5 - P95
<b>Males</b>											
Austria	10-14	248	14.6	3.2							
Belgium	13-15	74	14.7	2.1							
Bulgaria	10-14	167	12.5	2.3		80.5	27.0		1.97	0.75	
Denmark	10-13	164	15	2.3	11-18	79	20	49-109			
France	10-14	160	15.5	0.2							
Germany	10-11	199	13.8	2.3	10.3-18.1	64.4	16.2	43.1-94.5			
	12	114	13.3	1.9	10.5-16.5	82.5	29.1	46.2-135.5			
	13-14	214	13.7	2.3	10.3-17.4	94.0	33.9	47.5-159.6			
Hungary	11-14	124	14.6	2.0		89.7	18.9		1.99	0.59	
Ireland	9-12	148	13.6	2.4	9.5-18.0	64.2	15.8	40.9-90.9			
	13-14	95	15.3	2.6	10.7-19.9	81.7	23.0	49.3-125.7			
Italy	10-<18	108	15.6	1.9	12.9-19.2	99.3	26.2	62.8-147.1	1.82	0.59	1.02-3.22
The Netherlands	9-13	351	13.1 <sup>1</sup>		9.8-16.8	75 <sup>1</sup>		50-106			
Norway	13	590	15.0	3.0							
Poland	10-12	128	11.5	2.9		69.0	25.2				
	13-15	118	11.8	3.2		90.0	35.0				
Spain	10-14	66	16.9	2.1							
Sweden	11-12	517	15.9	2.7	11.8-20.5	72	19	44-106			
United Kingdom	11-18	238	14.9	2.9	8.8-20.5 <sup>2</sup>	73.7	20.7	33.7-116.3 <sup>2</sup>			
<b>Females</b>											
Austria	10-14	239	14.1	3.0							
Belgium	13-15	89	15.3	2.5							
Bulgaria	10-14	180	12.3	2.7		66.6	21.9		1.72	0.71	
Denmark	10-13	196	14	2.2	11-18	65	18	36-91			
France	10-14	144	15.6	0.2							
Germany	10-11	198	13.7	2.4	10.3-18.0	60.7	15.3	32.2-86.4			
	12	103	13.1	1.9	9.6-16.3	70.4	23.7	36.4-121.0			
	13-14	230	13.1	2.2	9.7-17.0	73.0	21.7	40.5-115.3			
Hungary	11-14	111	13.9	1.9		75.4	15.3		1.73	0.60	
Ireland	9-12	150	13.5	2.2	9.8-17.2	55.6	13.4	35.8-80.5			
	13-14	93	14.1	2.2	10.3-17.8	59.2	16.3	32.2-87.1			
Italy	10-<18	139	15.8	2.2	12.2-19.7	81.8	20.1	49.4-118.7	1.74	0.56	0.97-2.94
The Netherlands	9-13	352	13.0 <sup>1</sup>		9.6-17.1	64 <sup>1</sup>		42-90			
Norway	13	515	14.0	3.0							
Poland	10-12	121	11.2	2.5		58.0	19.6				
	13-15	134	12.1	2.9		69.8	28.1				
Spain	10-14	53	17.6	1.9							
Sweden	11-12	499	15.4	2.7	11.1-20.2	62	17	37-91			
United Kingdom	11-18	215	14.2	2.6	8.6-19.8 <sup>2</sup>	57.3	14.9	19.7-84.4 <sup>2</sup>			

<sup>1</sup>Median (mean not available); <sup>2</sup>P2.5-P97.5



**APPENDIX 2F: PROTEIN INTAKE OF ADOLESCENTS AGED ~15-18 YEARS AND OVER IN EUROPEAN COUNTRIES**

Country	Age (years)	N	Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
			mean	SD	P5 – P95	mean	SD	P5 - P95	mean	SD	P5 - P95
<b>Males</b>											
Austria	14-19	1,527	16.1	4.0							
Belgium	15-18	405	13.8	2.1							
Bulgaria	14-18	178	12.5	2.7		86.9	28.1		1.44	0.48	
Denmark	14-17	101	15	2.3	11-19	88	28	46-135			
France	15-18	181	15.7	0.3							
Germany	15-17	294	13.9	2.5	10.6-17.1	116.1	48.2	62.3-201.0			
Ireland	15-17	129	15.2	2.4	11.9-19.9	88.2	24.7	52.3-144.9			
The Netherlands	14-18	352	13.4 <sup>1</sup>		10.1-17.2	86 <sup>1</sup>		59-119			
Norway	16-19	92	14.3			114					
Poland	16-18	130	12.4	2.8		105.6	38.4				
Slovenia	14-16	1,085	14.8								
Spain	15-18	61	17.8	2.6							
<b>Females</b>											
Austria	14-19	1,422	14.7	4.1							
Belgium	15-18	401	13.7	2.1							
Bulgaria	14-18	190	12.5	3.0		65.6	23.2		1.26	0.49	
Denmark	14-17	134	14	2.2	11-18	61	21	28-98			
France	15-18	222	15.6	0.2							
Germany	15-17	317	12.9	2.3	9.6-16.7	75.0	32.3	37.8-125.6			
Ireland	15-17	124	14.4	2.8	9.7-19.8	61.1	19.6	32.0-98.0			
The Netherlands	14-18	354	13.7 <sup>1</sup>		10.2-18.0	67 <sup>1</sup>		45-94			
Norway	16-19	86	15.3			80					
Poland	16-18	122	12.1	2.9		66.3	28.6				
Slovenia	14-16	1,346	13.5								
Spain	15-18	57	18.0	2.5							

<sup>1</sup>Median (mean not available)

### APPENDIX 3A: POPULATION, METHODS AND PERIOD OF DIETARY ASSESSMENT IN ADULTS IN EUROPEAN COUNTRIES

Country	Population	Dietary assessment method	Year of survey	Reference
<b>Austria</b>	Males and females aged 19-64 years	24-hour recall	2005-2006	(Elmadfa et al., 2009a; Elmadfa et al., 2009b)
	Males and females aged 65 and over	3-day record	2007-2008	(Elmadfa et al., 2009a; Elmadfa et al., 2009b)
<b>Belgium</b>	Males and females aged 19-59 years	2 x 24-hour recall	2004-2005	(De Vriese et al., 2006)
	Males and females aged 60-74 years	2 x 24-hour recall	2004-2005	(De Vriese et al., 2006)
	Males and females aged 75 and over	2 x 24-hour recall	2004-2005	(De Vriese et al., 2006)
<b>Bulgaria</b>	Males and females aged 18-30 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Males and females aged 30-60 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Males and females aged 60-75 years	24-hour recall	1998	(Abrasheva et al., 1998)
	Males and females aged >75 years	24-hour recall	1998	(Abrasheva et al., 1998)
<b>Czech Republic</b>	Males and females aged 19-64 years	24-hour recall	2000-2001	(Cifkova and Skodova, 2004; Elmadfa et al., 2009a)
<b>Denmark</b>	Males and females aged 18-75 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Males and females aged 18-24 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Males and females aged 25-34 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Males and females aged 35-44 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Males and females aged 45-54 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Males and females aged 55-64 years	7-day record	2003-2008	(Pedersen et al., 2010)
	Males and females aged 65-75 years	7-day record	2003-2008	(Pedersen et al., 2010)
<b>Estonia</b>	Males and females aged 19-64 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Males and females aged 19-34 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Males and females aged 35-49 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Males and females aged 50-64 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
<b>Finland</b>	Males and females aged 25-64 years	48-hour recall	2007	(Paturi et al., 2008; Pietinen et al., 2010)
	Males and females aged 25-34 years	48-hour recall	2007	(Paturi et al., 2008)
	Males and females aged 35-44 years	48-hour recall	2007	(Paturi et al., 2008)
	Males and females aged 45-54 years	48-hour recall	2007	(Paturi et al., 2008)
	Males and females aged 55-64 years	48-hour recall	2007	(Paturi et al., 2008)
	Males and females aged 65-75 years	48-hour recall	2007	(Paturi et al., 2008)
<b>France</b>	Males and females aged 19-64 years	3 x 24-hour recall	2006-2007	(Elmadfa et al., 2009a)
	Males and females aged 65-74 years	3 x 24-hour recall	2006-2007	(Elmadfa et al., 2009a)
<b>Germany</b>	Males and females aged 19-80 years	24-hour recall + Dietary History	2005-2006	(Anonymous, 2008)
	Males and females aged 19-24 years	24-hour recall + Dietary History	2005-2006	(Anonymous, 2008)
	Males and females aged 25-34 years	24-hour recall + Dietary History	2005-2006	(Anonymous, 2008)
	Males and females aged 35-50 years	24-hour recall + Dietary History	2005-2006	(Anonymous, 2008)
	Males and females aged 51-64 years	24-hour recall + Dietary History	2005-2006	(Anonymous, 2008)
	Males and females aged 65-80 years	24-hour recall + Dietary History	2005-2006	(Anonymous, 2008; Elmadfa et al., 2009a)

Country	Population	Dietary assessment method	Year of survey	Reference
Greece	Males and females aged 19-64 years	Food frequency questionnaire + 24-hour recall in sub group	1994-1999	(Elmadfa et al., 2009a)
	Males and females aged 65 and over	Food frequency questionnaire	1994-1999	(Elmadfa et al., 2009a)
Hungary	Males and females aged 18-59	3-day record	2003-2004	(Elmadfa et al., 2009a; Rodler et al., 2005)
	Males and females aged 18-34 years	3-day record	2003-2004	(Elmadfa et al., 2009a; Rodler et al., 2005)
	Males and females aged 35-59 years	3-day record	2003-2004	(Elmadfa et al., 2009a; Rodler et al., 2005)
	Males and females aged 60 and over	3-day record	2003-2004	(Elmadfa et al., 2009a; Rodler et al., 2005)
Ireland	Males and females aged 18-64 years	4-day record	2008-2010	(IUNA (Irish Universities Nutrition Alliance), c)
	Males and females aged 18-35 years	4-day record	2008-2010	(IUNA (Irish Universities Nutrition Alliance), c)
	Males and females aged 36-50 years	4-day record	2008-2010	(IUNA (Irish Universities Nutrition Alliance), c)
	Males and females aged 51-64 years	4-day record	2008-2010	(IUNA (Irish Universities Nutrition Alliance), c)
	Males and females aged 65-90 years	4-day record	2008-2010	(IUNA (Irish Universities Nutrition Alliance), c)
Italy	Males and females aged 18-<65years	Consecutive 3-day food record	2005-2006	(Sette et al., 2010)
	Males and females aged 65 and over	Consecutive 3-day food record	2005-2006	(Sette et al., 2010)
Latvia	Males and females aged 19-64 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Males and females aged 19-34 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Males and females aged 35-49 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
	Males and females aged 50-64 years	24-hour recall	1997	(Elmadfa et al., 2009a; Pomerleau et al., 2001)
Lithuania	Males and females aged 19-64 years	24-hour recall	2007	(Elmadfa et al., 2009a)
The Netherlands	Males and Females aged 19-30 years	2 non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)
	Males and Females aged 31-50 years	2 non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)
	Males and Females aged 51-69 years	2 non-consecutive 24-hour dietary recalls	2007-2010	(van Rossum et al., 2011)
Norway	Males and females aged 19-64 years	Food frequency questionnaire	1997	(Elmadfa et al., 2009a)
	Males and females aged 20-29 years	Food frequency questionnaire	1997	(Johansson and Sovoll, 1999)
	Males and females aged 30-39 years	Food frequency questionnaire	1997	(Johansson and Sovoll, 1999)
	Males and females aged 40-49 years	Food frequency questionnaire	1997	(Johansson and Sovoll, 1999)
	Males and females aged 50-59 years	Food frequency questionnaire	1997	(Johansson and Sovoll, 1999)
	Males and females aged 65 and over	Food frequency questionnaire	1997	(Elmadfa et al., 2009a)
	Males and females aged 60-69 years	Food frequency questionnaire	1997	(Johansson and Sovoll, 1999)
	Males and females aged 70-79 years	Food frequency questionnaire	1997	(Johansson and Sovoll, 1999)
Poland	Males and females aged 19-25 years	24-hour recall	2000	(Szponar et al., 2003)
	Males and females aged 26-60 years	24-hour recall	2000	(Szponar et al., 2003)
	Males and females aged 61 and over	24-hour recall	2000	(Szponar et al., 2003)
Portugal	Males and females aged 18-≥65 years	Food frequency questionnaire	1999-2003	(Elmadfa et al., 2009a). <i>Data collected in Porto.</i>
	Males and females aged 18-39 years	Food frequency questionnaire	1999-2003	(Lopes et al., 2006)
	Males and females aged 40-49 years	Food frequency questionnaire	1999-2003	(Lopes et al., 2006)
	Males and females aged 50-64 years	Food frequency questionnaire	1999-2003	(Lopes et al., 2006)

Country	Population	Dietary assessment method	Year of survey	Reference
	Males and females aged 65 and over	Food frequency questionnaire	1999-2003	(Elmadfa et al., 2009a; Lopes et al., 2006). <i>Data collected in Porto.</i>
<b>Romania</b>	Males and females aged 19-64 years	Personal interview	2006	(Elmadfa et al., 2009a)
	Males and females aged 65 and over	Personal interview	2006	(Elmadfa et al., 2009a)
<b>Slovenia</b>	Males and females aged 18-65 years	Food frequency questionnaire	2007-2008	(Gabrijelčič Blenkuš et al., 2009)
<b>Spain</b>	Males and females aged 18-24 years	2 non-consecutive 24-hour recalls	2002-2003	(Serra-Majem et al., 2007). <i>Data collected in Catalonia.</i>
	Males and females aged 25-44 years	2 non-consecutive 24-hour recalls	2002-2003	(Serra-Majem et al., 2007). <i>Data collected in Catalonia.</i>
	Males and females aged 45-64 years	2 non-consecutive 24-hour recalls	2002-2003	(Serra-Majem et al., 2007). <i>Data collected in Catalonia.</i>
	Males and females aged 65-75 years	2 non-consecutive 24-hour recalls	2002-2003	(Serra-Majem et al., 2007). <i>Data collected in Catalonia.</i>
<b>Sweden</b>	Males and females aged 17-74 years	7-day record	1997-1998	(Becker and Pearson, 2002)
	Males and females aged 17-24 years	7-day record	1997-1998	(Becker and Pearson, 2002)
	Males and females aged 25-34 years	7-day record	1997-1998	(Becker and Pearson, 2002)
	Males and females aged 35-44 years	7-day record	1997-1998	(Becker and Pearson, 2002)
	Males and females aged 45-54 years	7-day record	1997-1998	(Becker and Pearson, 2002)
	Males and females aged 55-64 years	7-day record	1997-1998	(Becker and Pearson, 2002)
	Males and females aged 65-74 years	7-day record	1997-1998	(Becker and Pearson, 2002)
<b>United Kingdom</b>	Males and females aged 19-64 years	4-day food diary	2008-2010	(Bates et al., 2011)
	Males and females aged 65 years and over	4-day food diary	2008-2010	(Bates et al., 2011)

**APPENDIX 3B: PROTEIN INTAKE OF ADULTS AGED ~19-65 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	N			Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
		mean	SD	P5 - P95	mean	SD	P5 - P95	mean	SD	P5 - P95			
<b>Males</b>													
Austria	19-64	778	16.8	4.9									
Belgium	19-59	413	16.0	3.1									
Czech Republic	19-64	1,046	14.1	4.0									
Denmark	18-75	1,569	14	2.3	11-17 <sup>1</sup>	87	25	57-118 <sup>1</sup>					
Estonia	19-64	900	14.7	4.7					1.0	0.6			
Finland	25-64	730	16.8	3.7		89	31						
France	19-64	852	16.3	0.1									
Germany	19-64	4,912	14.6	3.2									
Greece	19-64	8,365	14.1	1.7									
Hungary	18->60	473	14.7	2.0		102.0	23.6						
Ireland	18-64	634	17.1	3.8		100.2	28.4						
Italy	18-<65	1,068	15.7	2.2	12.6-19.3	92.6	25.3	56.2-136.1	1.20	0.36	0.71-1.83		
Latvia	19-64	1,065	13.7	4.2					1.1	0.6			
Lithuania	19-64	849	16.5	5.2									
Norway	19-64	1,050	16.0	2.0									
Portugal	18->65	917	17.6	2.4	13.4-21.6	103.0	24.5	66.3-146.6					
Romania	19-64	177	17.8	3.8									
Slovenia	18-65	n.a.	14.6										
Sweden	17-74	589	16	2	13-19	90	23	55-130					
United Kingdom	19-64	346	16.5	4.8	10.1-25.3 <sup>2</sup>	88.1	35.7	44.9-151.3 <sup>2</sup>					
<b>Females</b>													
Austria	19-64	1,345	15.4	2.8									
Belgium	19-59	460	16.7	3.4									
Czech Republic	19-64	1,094	14.7	7.7									
Denmark	18-75	1,785	15	2.4	12-18 <sup>1</sup>	67	18	46-91 <sup>1</sup>					
Estonia	19-64	1,115	15.0	4.4					0.9	0.5			
Finland	25-64	846	17.2	4.14		67	21						
France	19-64	1,499	17.0	0.1									
Germany	19-64	6,016	14.4	2.6									
Greece	19-64	12,034	14.4	1.7									
Hungary	18->60	706	14.6	1.9		79.7	18.0						
Ireland	18-64	640	16.7	3.8		70.4	19.8						
Italy	18-<65	1,245	15.9	2.3	12.4-19.9	76.0	19.5	45.4-108.6	1.25	0.36	0.71-1.90		
Latvia	19-64	1,235	13.7	4.8					0.9	0.5			
Lithuania	19-64	1,087	16.7	6.2									
Norway	19-64	1,146	16.0	3.0									
Portugal	18->65	1,472	19.0	2.4	15.2-22.9	98.2	24.4	61.4-142.7					
Romania	19-64	341	17.1	3.6									
Slovenia	18-65	n.a.	14.2										
Sweden	17-74	626	16	2	13-20	73	17	47-102					
United Kingdom	19-64	461	16.5	4.1	10.3-26.6 <sup>2</sup>	65.4	18.1	32.1-101.7 <sup>2</sup>					

<sup>1</sup>P10-P90; <sup>2</sup>P2.5-P97.5; n.a.: not available

**APPENDIX 3C: PROTEIN INTAKE OF ADULTS AGED ~19-34 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	N	Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
			mean	SD	P5 – P95	mean	SD	P5 - P95	mean	SD	P5 - P95
<b>Males</b>											
<b>Bulgaria</b>	18-30	208	12.6	2.7		86.2	34.9		1.20	0.47	
<b>Denmark</b>	18-24	105	15	2.4	11-19	96	28	50-147			
	25-34	234	14	2.3	11-19	93	25	58-137			
<b>Estonia</b>	19-34	396	14.3	4.6					1.1	0.6	
<b>Finland</b>	25-34	137	16.5	3.5		95	35				
<b>Germany</b>	19-24	510				101.8	1.84 <sup>1</sup>	51.4-189.0			
	25-34	690				99.0	1.50 <sup>1</sup>	53.2-168.0			
<b>Hungary</b>	18-34	136	14.8	2.0		108.6	23.6				
<b>Ireland</b>	18-35	276	16.8	4.3		105.0	32.1				
<b>Latvia</b>	19-34	337	13.5	4.1					1.2	0.6	
<b>The Netherlands</b>	19-30	356	13.9 <sup>2</sup>		10.5-17.8	94 <sup>2</sup>		66-128			
<b>Norway</b>	20-29	248	15.0			109					
	30-39	269	15.6			103					
<b>Poland</b>	19-25	191	12.8	2.7		114.3	37.3				
<b>Portugal</b>	18-39	179	17.8	2.4	14.0-21.7	109.9	23.9	73.1-147.2			
<b>Spain</b>	18-24	127	18.0			104.1					
	25-44	326	18.8			101.2					
<b>Sweden</b>	17-24	67	15	2	12-19	92	27	48-144			
	25-34	128	15	2	12-18	91	21	58-129			
<b>Females</b>											
<b>Bulgaria</b>	18-30	204	13.1	3.4		62.4	23.1		1.10	0.44	
<b>Denmark</b>	18-24	150	14	2.2	11-18	66	18	41-98			
	25-34	340	15	2.4	11-18	70	18	42-99			
<b>Estonia</b>	19-34	459	14.6	4.5					1.0	0.5	
<b>Finland</b>	25-34	180	17.0	4.7		69	23				
<b>Germany</b>	19-24	510				65.2	1.00 <sup>1</sup>	35.8-106.5			
	25-34	972				69.6	0.73 <sup>1</sup>	40.4-108.5			
<b>Hungary</b>	18-34	176	14.4	1.9		81.5	17.4				
<b>Ireland</b>	18-35	255	15.7	3.3		68.6	20.5				
<b>Latvia</b>	19-34	342	13.3	5.0					1.0	0.5	
<b>The Netherlands</b>	19-30	347	14.5 <sup>2</sup>		10.8-18.9	71 <sup>2</sup>		49-99			
<b>Norway</b>	20-29	268	15.1			76					
	30-39	289	15.8			76					
<b>Poland</b>	19-25	211	12.8	3.3		61.2	27.0				
<b>Portugal</b>	18-39	299	19.1	2.6	15.0-23.1	101.5	25.0	62.5-143.9			
<b>Spain</b>	18-24	182	18.5			83.7					
	25-44	376	19.4			80.8					
<b>Sweden</b>	17-24	70	15	2	12-20	70	19	35-103			
	25-34	132	16	2	12-20	73	16	49-103			

<sup>1</sup>SE; <sup>2</sup>Median (mean not available)

**APPENDIX 3D: PROTEIN INTAKE OF ADULTS AGED ~35-64 YEARS IN EUROPEAN COUNTRIES**

Country	Age (years)	N	Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
			mean	SD	P5 - P95	mean	SD	P5 - P95	mean	SD	P5 - P95
<b>Males</b>											
<b>Bulgaria</b>	30-60	224	12.3	2.5		84.7	29.1		1.07	0.37	
<b>Denmark</b>	35-44	318	14	2.0	11-18	93	27	55-134			
	45-54	336	14	2.2	11-18	86	23	50-125			
	55-64	336	14	2.4	11-19	82	24	49-129			
<b>Estonia</b>	35-49	319	14.7	4.8					1.0	0.5	
	50-64	185	15.4	4.7					1.0	0.5	
<b>Finland</b>	35-44	177	16.6	3.5		91	32				
	45-54	190	17.1	3.8		91	27				
	55-64	226	16.7	3.8		84	28				
<b>Germany</b>	35-50	2,079				93.9	0.74 <sup>1</sup>	51.0-151.5			
	51-64	1,633				85.7	0.69 <sup>1</sup>	47.5-136.6			
<b>Hungary</b>	35-59	199	14.7	2.0		104.5	22.6				
<b>Ireland</b>	36-50	205	17.4	3.4		99.0	25.8				
	51-64	153	17.2	3.5		93.1	22.3				
<b>Latvia</b>	35-49	372	13.8	4.5					1.1	0.6	
	50-64	356	13.8	4.0					1.0	0.5	
<b>The Netherlands</b>	31-50	348	15.0 <sup>2</sup>		11.4-19.0	98 <sup>2</sup>		70-133			
	51-69	351	16.2 <sup>2</sup>		12.5-20.4	95 <sup>2</sup>		66-129			
<b>Norway</b>	40-49	256	16.0			97					
	50-59	196	16.2			92					
<b>Poland</b>	26-60	865	13.6	3.2		103.6	41.4				
<b>Portugal</b>	40-49	197	17.3	2.5	13.2-21.5	105.0	22.8	70.9-146.7			
	50-64	295	17.6	2.3	13.7-21.7	102.8	24.9	66.9-147.8			
<b>Spain</b>	45-64	265	20.0			95.4					
<b>Sweden</b>	35-44	143	16	2	13-19	91	22	57-133			
	45-54	118	16	2	12-20	91	23	56-129			
	55-64	68	16	2	13-20	85	20	49-118			
<b>Females</b>											
<b>Bulgaria</b>	30-60	224	12.5	2.8		60.9	24.7		0.95	0.42	
<b>Denmark</b>	35-44	412	15	2.4	11-19	71	18	44-104			
	45-54	359	15	2.4	11-19	65	16	39-92			
	55-64	326	15	2.5	11-19	63	16	40-94			
<b>Estonia</b>	35-49	376	15.2	4.5					0.9	0.5	
	50-64	280	15.3	4.3					0.8	0.4	
<b>Finland</b>	35-44	211	17.0	4.0		69	21				
	45-54	232	17.3	3.9		66	18				
	55-64	223	17.5	3.6		64	21				
<b>Germany</b>	35-50	2,694				68.9	0.41 <sup>1</sup>	39.3-106.7			
	51-64	1,840				67.3	0.49 <sup>1</sup>	38.7-105.2			
<b>Hungary</b>	35-59	295	14.7	2.0		81.6	17.5				
<b>Ireland</b>	36-50	232	17.0	4.0		70.6	20.2				
	51-64	153	17.9	3.7		73.1	17.8				
<b>Latvia</b>	35-49	396	13.7	4.4					0.9	0.5	
	50-64	496	14.0	4.9					0.8	0.4	
<b>The Netherlands</b>	31-50	351	15.6 <sup>2</sup>		11.7-20.1	75 <sup>2</sup>		52-103			
	51-69	353	16.3 <sup>2</sup>		12.4-21.0	74 <sup>2</sup>		51-102			
<b>Norway</b>	40-49	289	16.5			74					
	50-59	196	16.9			74					
<b>Poland</b>	26-60	1,035	13.1	3.5		63.9	27.1				
<b>Portugal</b>	40-49	340	18.9	2.3	15.4-22.7	101.6	23.0	66.0-144.6			
	50-59	494	19.1	2.4	15.5-23.0	99.9	24.5	63.9-140.1			
<b>Spain</b>	45-64	337	20.2			76.4					
<b>Sweden</b>	35-44	132	16	2	13-20	71	15	47-98			
	45-54	153	16	2	13-21	73	17	49-102			
	55-64	81	17	2	14-21	75	16	49-99			

<sup>1</sup>SE; <sup>2</sup>Median (mean not available)

**APPENDIX 3E: PROTEIN INTAKE OF ADULTS AGED ~65 YEARS AND OVER IN EUROPEAN COUNTRIES**

Country	Age (years)	N	Protein (E%)			Protein (g/d)			Protein (g/kg bw x d <sup>-1</sup> )		
			mean	SD	P5 - P95	mean	SD	P5 - P95	mean	SD	P5 - P95
<b>Males</b>											
Austria	65+	147	14.9	3.1							
Belgium	60-74	416	16.9	2.7							
	75+	389	16.0	3.2							
Bulgaria	60-75	186	12.7	2.6		75.8	31.2		1.00	0.41	
	76+	101	12.6	2.2		66.9	23.0		0.94	0.30	
Denmark	65-75	240	14	2.6	10-18	76	22	44-113			
Finland	65-74	229	17.4	3.8		78	27				
France	65-74	130	16.5	0.3							
Germany	65-80	1,469	14.5	2.6		77.8	0.59 <sup>1</sup>	45.0-119.7			
Greece	65+	2,508	14.1	1.7							
Hungary	60+	138	14.7	2.1		91.9	21.7				
Ireland	65+	106	17.7	3.7		77.6	34.3				
Italy	65+	202	15.5	2.0	12.2-18.8	88.2	21.4	55.6-124.5	1.15	0.30	0.70-1.67
Norway	65+	176	16.0	2.0							
	60-69	131	16.3			83					
	70-79	106	16.5			85					
Poland	61+	226	13.5	3.3		83.8	33.6				
Portugal	65+	246	17.5	2.4	13.0-21.4	96.7	24.5	63.2-142.2			
Romania	65+	177	17.2	3.5							
Spain	65-75	122	19.5			77.6					
Sweden	65-74	65	16	2	13-20	87	24	53-131			
United Kingdom	65+	96	16.3	3.4	10.8-23.1 <sup>2</sup>	79.7	27.0	33.9-123.1 <sup>2</sup>			
<b>Females</b>											
Austria	65+	202	15.0	2.5							
Belgium	60-74	406	16.7	2.8							
	75+	355	17.0	3.8							
Bulgaria	60-75	194	13.3	3.2		63.1	23.4		0.94	0.35	
	76+	113	13.1	2.8		58.5	20.8		0.97	0.41	
Denmark	65-75	198	14	2.6	11-19	62	17	36-95			
Finland	65-74	234	17.6	3.4		60	18				
France	65-74	219	17.5	0.3							
Germany	65-80	1,562	14.4	2.7		61.6	0.45 <sup>1</sup>	34.9-91.6			
Greece	65+	3,600	14.4	1.8							
Hungary	60+	235	14.5	1.8		76.0	18.5				
Ireland	65+	120	18.0	2.9		69.4	17.8				
Italy	65+	316	15.7	2.4	12.4-19.9	71.4	18.8	41.0-100.7	1.12	0.32	0.63-1.69
Norway	65+	166	17.0	3.0							
	60-69	137	17.3			73					
	70-79	109	16.5			67					
Poland	61+	365	13.3	3.5		64.2	26.0				
Portugal	65+	339	18.7	2.3	15.0-22.6	89.5	23.2	57.2-133.4			
Romania	65+	341	16.3	3.0							
Spain	65-75	122	20.2			68.0					
Sweden	65-74	58	16	3	12-22	75	20	41-119			
United Kingdom	65+	128	17.1	2.9	12.4-23.2 <sup>2</sup>	64.2	13.9	38.0-100.6 <sup>2</sup>			

<sup>1</sup>SE; <sup>2</sup>P2.5-P97.5



#### **APPENDIX 4: CALCULATION OF PRI FOR INFANTS, CHILDREN AND ADOLESCENTS**

The PRI for infants from six months onwards and for children is calculated as follows:

$PRI = AR + 1.96 SD_{combined}$ , with the  $SD_{combined}$  calculated from the formula:

$$SD_{combined} = (\sqrt{[CV_{maintenance} \times maintenance\ requirement]^2 + [CV_{growth} \times growth\ requirement]^2}).$$

$CV_{maintenance}$  is 0.12, the maintenance requirement is given in Tables 9 and 10,  $CV_{growth}$  can be calculated from the SD for growth given in Table 29 of the WHO/FAO/UNU report (WHO/FAO/UNU, 2007), and the growth requirement is the rate of protein deposition (see Tables 9 and 10) divided by the efficiency of dietary protein utilisation.

## GLOSSARY / ABBREVIATIONS

<b>AFSSA</b>	Agence Française de Sécurité Sanitaire des Aliments (French Food Safety Agency)
<b>ANSES</b>	Agence Nationale de Sécurité Sanitaire de l'alimentation, de l'environnement et du travail (French Agency for Food, Environmental and Occupational Health and Safety)
<b>AR</b>	Average Requirement
<b>BCAA</b>	Branched-chain amino acid
<b>BMI</b>	Body mass index
<b>BV</b>	Biological value
<b>bw</b>	Body weight
<b>CI</b>	Confidence interval
<b>CIQUAL</b>	Centre d'information sur la qualité des aliments (French data centre on food quality)
<b>COMA</b>	Committee on Medical Aspects of Food and Nutrition Policy
<b>CV</b>	Coefficient of variation
<b>d</b>	day
<b>D-A-CH</b>	Deutschland-Austria-Confoederatio Helvetica
<b>DNA</b>	Deoxyribonucleic acid
<b>DoH</b>	Department of Health
<b>DRV</b>	Dietary Reference Value
<b>EAR</b>	Estimated Average Requirement
<b>EC</b>	European Commission
<b>EFSA</b>	European Food Safety Authority
<b>EU</b>	European Union
<b>f</b>	female
<b>FAO</b>	Food and Agriculture Organisation
<b>GFR</b>	Glomerular filtration rate
<b>Health ABC Study</b>	Health, Aging, and Body Composition Study
<b>IGF</b>	Insulin-like growth factor
<b>IGFBP</b>	IGF-binding protein

<b>IoM</b>	U.S. Institute of Medicine
<b>m</b>	male
<b>mo</b>	months
<b>mTOR</b>	Mammalian target of rapamycin
<b>N</b>	Nitrogen
<b>n.a.</b>	Not available
<b>NH<sub>3</sub></b>	Ammonia
<b>NH<sub>4</sub><sup>+</sup></b>	Ammonium
<b>NNR</b>	Nordic Nutrition Recommendations
<b>NPN</b>	Non-protein nitrogen
<b>NPPU</b>	Net post-prandial protein utilisation
<b>NPU</b>	Net protein utilisation
<b>PD-CAAS</b>	Protein digestibility-corrected amino acid score
<b>PER</b>	Protein efficiency ratio
<b>PRI</b>	Population Reference Intake
<b>RDA</b>	Recommended Dietary Allowance
<b>RNA</b>	Ribonucleic acid
<b>SCF</b>	Scientific Committee on Food
<b>SD</b>	Standard deviation
<b>SE</b>	Standard error
<b>UL</b>	Tolerable Upper Intake Level
<b>UNU</b>	United Nations University
<b>WHO</b>	World Health Organisation
<b>y</b>	year