



Effect of the nematophagous fungus *Pochonia chlamydosporia* on soil content of ascarid eggs and infection levels in exposed hens

Thapa, Sundar; Thamsborg, Stig M; Wang, Rui; Meyling, Nicolai V; Dalgaard, Tina S; Petersen, Heidi Huus; Mejer, Helena

Published in:
Parasites & Vectors

Link to article, DOI:
[10.1186/s13071-018-2898-1](https://doi.org/10.1186/s13071-018-2898-1)

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Thapa, S., Thamsborg, S. M., Wang, R., Meyling, N. V., Dalgaard, T. S., Petersen, H. H., & Mejer, H. (2018). Effect of the nematophagous fungus *Pochonia chlamydosporia* on soil content of ascarid eggs and infection levels in exposed hens. *Parasites & Vectors*, 11, [319]. <https://doi.org/10.1186/s13071-018-2898-1>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH

Open Access



Effect of the nematophagous fungus *Pochonia chlamydosporia* on soil content of ascarid eggs and infection levels in exposed hens

Sundar Thapa^{1*}, Stig M. Thamsborg¹, Rui Wang², Nicolai V. Meyling³, Tina S. Dalgaard⁴, Heidi H. Petersen⁵ and Helena Mejer¹

Abstract

Background: The nematophagous fungus *Pochonia chlamydosporia* can degrade ascarid (e.g. *Ascaridia galli*) eggs in agar and soil *in vitro*. However, it has not been investigated how this translates to reduced infection levels in naturally exposed chickens. We thus tested the infectivity of soil artificially contaminated with *A. galli* (and a few *Heterakis gallinarum*) eggs and treated with *P. chlamydosporia*. Sterilised and non-sterilised soils were used to examine any influence of natural soil biota.

Methods: Unembryonated eggs were mixed with sterilised (S)/non-sterilised (N) soil, either treated with the fungus (F) or left as untreated controls (C) and incubated (22 °C, 35 days) to allow eggs to embryonate and fungus to grow. Egg number in soil was estimated on days 0 and 35 post-incubation. Hens were exposed to the soil (SC/SF/NC/NF) four times over 12 days by mixing soil into the feed. On day 42 post-first-exposure (p.f.e.), the hens were euthanized and parasites were recovered. Serum *A. galli* IgY level and ascarid eggs per gram of faeces (EPG) were examined on days -1 and 36 (IgY) or 40 p.f.e. (EPG).

Results: Egg recovery in SF soil was substantially lower than in SC soil, but recovery was not significantly different between NF and NC soils. SF hens had a mean worm count of 76 whereas the other groups had means of 355–453. Early mature/mature *A. galli* were recovered from SF hens whereas hens in the other groups harboured mainly immature *A. galli*. *Heterakis gallinarum* counts were low overall, especially in SF. The SF post-exposure IgY response was significantly lower while EPG was significantly higher compared to the other groups.

Conclusions: *Pochonia chlamydosporia* was very effective in reducing ascarid egg numbers in sterilised soil and thus worm burdens in the exposed hens. However, reduced exposure of hens shifted *A. galli* populations toward a higher proportion of mature worms and resulted in a higher faecal egg excretion within the study period. This highlights a fundamental problem in ascarid control: if not all eggs in the farm environment are inactivated, the resulting low level infections may result in higher contamination levels with associated negative long-term consequences.

Keywords: *Ascaridia galli*, *Heterakis gallinarum*, Fungus, Biological control

* Correspondence: sundar@sund.ku.dk

¹Section for Parasitology and Aquatic Pathobiology, Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Dyrølægevej 100, 1870 Frederiksberg C, Denmark
Full list of author information is available at the end of the article



Background

Ascaridia galli and *Heterakis* spp., collectively known as ascarids, are economically important intestinal nematodes of chickens worldwide. *Ascaridia galli* can impair the health [1–3], productivity [4–7] and welfare of chickens [8]. Moreover, *A. galli* can reduce the vaccine efficacy against Newcastle disease [9, 10] and increase the susceptibility of chickens to other infectious diseases such as fowl cholera [11]. On rare occasions, *A. galli* can leave the host's intestine, migrate up the oviduct and become enclosed inside one of the hen's eggs, which is of aesthetic concern to the consumers [12, 13]. Compared to *A. galli*, *Heterakis* spp. are less pathogenic, but they can act as a vector for *in ovo* transmission of the protozoan *Histomonas meleagridis* to turkeys and chickens [14]. *Histomonas meleagridis* is pathogenic [15, 16] and re-emerging in layer flocks in many European countries, mainly after the ban of the prophylactic use of chemotherapeutics in the European Union (EU) member countries [17–20].

Recent studies have shown that *A. galli* and *Heterakis* spp. are highly prevalent in European organic laying hen flocks [21–23]. Both nematodes have a simple life-cycle that involves a pre-parasitic development phase (i.e. free-living nematode eggs) in the environment such as litter and soil and a parasitic phase in the chicken's intestine following ingestion of infective eggs [24, 25]. Ascarid eggs have thick shells [26, 27] and they can survive in the outdoor environment for up to 2–4 years [28, 29], but no effective means of inactivating eggs during or after embryonation in the farm yards and pastures are currently available. At present, control of ascarid infections by farmers therefore solely relies on flock treatment with commercial anthelmintics. Of the anthelmintics, only flubendazole and fenbendazole are available for use in layers in the EU [30, 31]. As hens are rapidly reinfected due to continuous exposure to eggs present in the surroundings and do not appear to acquire protective immunity [32–34], repeated treatments are thus necessary. However, overuse of these drugs may over time enhance the risk of selecting for anthelmintic resistance.

To be able to also combat the parasites in the environment, there is an increasing interest in using naturally occurring soil microfungi as is done to control agricultural nematode pests [35]. An isolate of *Pochonia chlamydosporia* (syn. *Verticillium chlamydosporium*) (Ascomycota: Hypocreales), a microfungus of global occurrence [36–38], that can mechanically and enzymatically degrade the egg shell components (protein and chitin) has already been developed as a biocontrol agent against plant-parasitic nematode eggs [39]. The same isolate [40] and two other isolates [41] of *P. chlamydosporia* have subsequently been shown to kill ~70% and > 80%,

respectively, of *A. galli* eggs in laboratory agar assays. However, the effect in a soil based assay was relatively lower (~45%) [42]. In the latter study, the fungal effectiveness was evaluated based on eggs recovered from soil before and after the fungal treatment. However, it is unknown if the recovered eggs judged visually as viable are indeed infective and whether the reduced contamination in the soil assay translates to lower worm burdens in chickens exposed to the fungus-treated soil. In addition, literature indicates that population composition of *A. galli* in chickens can be dose-dependent as shown by a reduced number of inhibited larvae in the host's intestine and a shorter prepatent period in lightly infected compared to heavily infected chickens [43, 44]. It is thus important to use an *in vivo* infection model to assess how changes in exposure level as a result of fungal treatment of soil may modulate worm population dynamics within the host as this may in turn alter on-farm transmission dynamics.

The overall aim of this study was to evaluate the *in vivo* infectivity of soil experimentally contaminated with ascarid eggs and treated with *P. chlamydosporia*. Both sterilised and non-sterilised soils were used to include any effect inherent to the natural soil biota and thus to evaluate the potential of *P. chlamydosporia* as an on-farm biocontrol agent.

Methods

Experimental design

Unembryonated ascarid eggs were added to Petri dishes with sterilised (S) or non-sterilised (N) soil, half of which were treated with spores of the fungus *P. chlamydosporia* (F) while the other half were untreated (C). The Petri dishes were incubated at 22 °C for 35 days to allow the fungus to grow and the eggs to reach infectivity. After incubation, three subgroups of 10 hens per soil treatment were exposed in-feed four times to the soil from one of the four treatments (SC, SF, NF and NC). The hens were euthanized on day 42 post-first-exposure (p.f.e.) and examined for *A. galli* and *Heterakis* spp. To estimate the number of eggs in the Petri dishes, recovery of eggs was tested before [i.e. day 0 post-incubation (p.i.)] and after incubation (day 35 p.i.) for each of the four soil treatments.

Origin and isolation of ascarid eggs

Before collecting faeces, the infection status of *A. galli* and *Heterakis* spp. in a Danish organic layer farm with a flock size of 3000 hens was examined through necropsy of 18 randomly selected hens. The prevalence was 89% for *A. galli* and 100% for *Heterakis* spp., and mean \pm S.E. worm burdens were 40 ± 9 and 80 ± 21 worms, respectively. Ascarid eggs (*A. galli* and *Heterakis* spp.) were isolated from fresh hen faeces collected from the ground using only the top part of the faeces as described by

Thapa et al. [29]. The eggs were stored in sterile demineralised water at 5 °C for 6 days. Before use, a subsample of eggs was embryonated in 0.1 N H₂SO₄ at 25 °C for 15 days to assess percentage embryonation (i.e. ability to develop larva) of the egg batch [42] which was 95 ± 1% (mean ± SE).

Preparation of fungal inoculum

Parboiled rice initially soaked for 1 h in demineralised water and autoclaved (121 °C, 15 min) inside a polypropylene bag (Labsolute®, 300 g rice per bag) was inoculated with 10 ml *P. chlamydosporia* Biotype 10 spore suspension [5.5×10^7 conidia and 1.6×10^4 chlamydospores in 0.05% Triton® X-100 (Merck KGaA, Darmstadt, Germany) harvested from 5-week-old culture in Sabouraud's dextrose agar]. After incubating the rice at 25 °C for 25 days in darkness, 30 ml 0.05% Triton® X-100 was added to each 10 g of rice granules in centrifuge tubes and shaken gently to separate spores from the rice. The mixture was filtered through a 900 µm sieve to remove rice particles and centrifuged (1831× g, 3 min) three times after re-suspension in 0.05% Triton® X-100. Spore concentration was adjusted to 4×10^8 conidia and 1.5×10^5 chlamydospores per ml suspension. Spore germination was determined [42] to be 96% and 91% for the conidia and chlamydospores, respectively.

Preparation of soil

In April 2016, 15 kg sandy loam soil (pH 6.8) was collected from a Danish experimental plot that was established in 2002 and treated annually (2003–2015) with source separated organic household waste compost (CH) and sown with spring cereals [45, 46]. After removing plant material and stones, the soil was sieved (3 mm) and thoroughly homogenised. Three kilograms of soil was sterilised by autoclaving (121 °C, 30 min) inside a polypropylene bag (Labsolute®, 200 g soil per bag, treatment S) while another 3 kg soil was kept without autoclaving (treatment N).

Fungal treatment of soil and eggs

For both soils (S, N), 44 replicate Petri dishes (14.5 × 2 cm) each containing 46 g soil and 2 ml ascarid egg suspension with approximately 8000 ± 260 eggs (mean ± SE) in sterile water were prepared. Both soil types were randomised into control (C) and fungus treatment (F). To all SF and NF dishes, 2 ml fungal suspension containing approximately 8×10^8 conidia and 3×10^5 chlamydospores of *P. chlamydosporia* in 0.05% Triton® X-100 was added whereas all SC and NC dishes received 2 ml 0.05% Triton® X-100 without fungus. The SC and SF dishes received an additional 635 µl of sterile water to balance the total moisture level between the S and N soils. The dishes were sealed with Parafilm 'M' and the initial

(i.e. pre-incubation) soil moisture level per dish (23–24% of the total soil weight) was estimated (105 °C, 24 h). Five random dishes per treatment were used to estimate day 0 p.i. egg recovery, while the remaining dishes were incubated at 22 °C for 35 days in darkness. The weight of each incubated dish was recorded at days 0 and 35 p.i. to determine soil moisture loss (%). On day 35 p.i., the dishes were opened and 4 ml sterile water was added to each dish, re-sealed with parafilm and stored at 10 °C for up to 18 days. On day 7 post-storage at 10 °C (i.e. after incubation was terminated), egg recovery (i.e. exposure level) was estimated in five random dishes per treatment (see section on recovery of eggs from soil). Soil from one random dish was selected and administered in the feed to a corresponding subgroup of hens (see section on animal exposure to parasites). This exposure was repeated on days 11, 15 and 19 after incubation was terminated.

Recovery of eggs from soil

Fifty millilitres of 0.5 M NaOH was added to each of the 40 dishes ($n = 5$ per treatment on day 0 and 35 p.i.) that was then stored at 5 °C for 16 h. The soil was washed through 212 and 20 µm sieves and the material on the latter was divided into four 50 ml tubes, centrifuged at 253× g for 7 min and the eggs were recovered as described by Thapa et al. [29]. For each dish, the egg quantity and development stage (unembryonated, pre-larvated, larvated or degenerated) was examined in a 20% subsample at 100× magnification [29].

Experimental animals and housing

One hundred thirty pullets (ISA Warren, 18-weeks-old), raised indoors without previous anthelmintic treatment, were obtained from a commercial breeder. On arrival (day -15 p.f.e.), 10 randomly selected pullets were euthanized and examined for ascarid infections of the breeder farm-origin (see section on recovery of worms). All necropsied pullets were found positive for tissue phase *A. galli* larvae (~0.5 mm long) with an overall mean ± SE worm burden of 194 ± 97 *A. galli*, while only two birds harboured luminal *Heterakis* spp. giving an overall burden of 1 ± 1 worm per hen. The remaining pullets ($n = 120$), after random allocation into 12 indoor pens ($c.2.8$ m², 10 pullets per pen), were therefore treated with flubendazole (Verminator®, 1.43 mg flubendazole per kg live weight daily) in the feed from days -13 to -6 p.f.e. The individual body weight of all birds was measured on days -1 and 36 p.f.e. The birds were given pelleted feed (17.5% crude protein, 4.5% crude fat) in two meals (110 g feed per bird per day) and water *ad libitum*. Crushed oyster shells were offered daily. Wood-chips and straw were

used as bedding material. Pens were enriched with a perch and nests, and cleaned thoroughly once weekly.

Animal exposure to parasites

The 12 pens were allocated to the four treatment groups (SC, SF, NC and NF) in triplets (i.e. three subgroups per treatment group). The hens were exposed to ascarid contaminated soil on days 0, 4, 8 and 12 p.f.e. to mimic a moderate trickle infection. On each exposure, entire soil from one Petri dish was transferred to a 500 ml container with 150 g feed of the morning meal and 50 ml tapwater, mixed thoroughly and spread in a tray (58 × 21 × 3 cm) in each pen. The feed was eaten within 10–15 min and the remainder of the meal was then given in the same tray.

Recovery of worms

The hens were euthanized by stunning and cervical dislocation on day 42 p.f.e. The *A. galli* worms in the small intestinal lumen were isolated using an agar-gel method [47] and collected using a 20 µm sieve. The tissue phase larvae of *A. galli* (day -15 and 42 p.f.e.) and *Heterakis* spp. (day -15 p.f.e.) were isolated from the intestinal/caecal tissue by pepsin (1:3000 IU)-HCl (30%) digestion [47] and collected on a 20 µm sieve. To recover luminal *Heterakis* spp. (day -15 and day 42 p.f.e.), the caeca were opened and stored in tap water at 5 °C. After 48 h, the caeca and contents were washed on a 20 µm sieve. All worm samples were stored in 70% ethanol and examined using a dissection microscope (30–40× magnification). All *A. galli* worms were categorised as < 0.5, 0.5–1.5, 1.5–3.0, 3.0–5.0 or 5.0–8.0 cm, whereas *Heterakis* spp. were categorised as < 0.5 or ≥ 0.5 cm. Moreover, *Heterakis* species were determined based on the length of the spicules [48, 49] of 50 randomly selected male worms (1 worm per hen and representing all experimental groups) after exposing each worm to a drop of 10% lactic acid in water (weight/weight).

Faecal egg counts

Individual faecal samples from all birds were collected on days -1 and 40 p.f.e. Ascarid eggs per gram faeces (EPG) was determined by a concentration McMaster technique (minimum detection limit: 20 EPG) using a flotation fluid of 500 g glucose monohydrate per litre of saturated NaCl solution (specific gravity: 1.27) [50].

Ascaridia galli antibody (IgY) levels

To determine the systemic antibody response as an indirect assessment of parasite exposure, individual blood samples from all birds were collected on days -1 and 36 p.f.e. from a wing vein. Serum was separated by centrifugation at 1000× *g* for 15 min and stored at -20 °C. The *A. galli* IgY level was determined by ELISA according to Norup et al. [51] using crude adult *A. galli* somatic

antigens and one replicate serum sample per animal per sampling day. A dilution series of a highly positive serum was used as standard and the highest concentration was set at the relative value 2.


Statistical analyses

All statistical analyses were performed using SAS 9.4 (Cary, NC, USA). The main and interaction effects of soil sterility (S, N), fungal treatment (C, F) and incubation time (days 0, 35 p.i.) on egg recovery from soil were analysed using a generalised linear model fitted with negative binomial distribution of errors (NBD) (procedure GENMOD). Soil moisture loss during incubation was analysed with a linear model (procedure GLM) with percent moisture loss as the outcome and soil sterility (S, N), fungal treatment (C, F) and their interaction as predictors. Body weight at day 0 p.f.e. and weight gain (days 0 to 36 p.f.e.) in relation to soil sterility (S, N) and fungal treatment (C, F) were analysed separately with a linear-mixed model (procedure MIXED) with subgroup (i.e. pen) as a random effect. Worm burden (total, *A. galli*, *H. gallinarum*), proportion (%) of *A. galli* in the intestinal tissue, proportion (%) of *A. galli* in each length category (< 0.5, 0.5–1.5, 1.5–3.0, 3.0–5.0, 5.0–8.0 cm) and at day 40 p.f.e. EPG was analysed with a generalised linear mixed model (procedure GLIMMIX, NBD) that included soil sterility (S, N), fungal treatment (C, F) and their interaction as fixed effects and subgroup as a random effect. The log-transformed IgY titre was analysed with a linear-mixed model (procedure MIXED) with soil sterility (S, N), fungal treatment (C, F), sampling time (days -1, 36 p.f.e.) and their interaction as fixed effects, subgroup as a random effect and individual bird as a repeated measurement. At group level, the linear relationships between worm burden (total ascarid or *A. galli*) and the IgY titre difference between pre- and post-exposures were examined using a Spearman method (procedure CORR). The goodness of fit of each GENMOD and GLIMMIX model was assessed with the ratio of Pearson's χ^2 and corresponding degrees of freedom. The normality of residuals of each GLM and MIXED model was examined by a q-q plot and a histogram, and homogeneity of residual variance assessed by residual plots. For each model, the *post-hoc* significant differences were determined with the differences of least squares means (Tukey-Kramer's adjustment for multiple comparisons, $P < 0.05$).

Results

Recovery of eggs from soil

On day 0 p.i., the mean number of eggs recovered from the SC, SE, NC and NF soils were 8702–9673 with no significant differences between the treatments ($P > 0.9950$ in all cases) (Fig. 1). Irrespective of treatment, > 97% of the




recovered eggs were unembryonated. On day 35 p.i., the mean egg number in the SC, SF, NC and NF soils was 5535 (36% reduction), 521 (94% reduction), 4176 (57% reduction) and 3201 eggs (65% reduction), respectively (Fig. 1). In the sterilised soil, the fungal treatment resulted in a significant reduction in egg recovery when compared to the control ($P < 0.0001$). In contrast, there was no such difference in the non-sterilised soil ($P = 0.5480$). This meant that there was a strong significant ($\chi^2 = 70.72$, $df = 4$, $P < 0.0001$) interaction between soil sterility, fungal treatment and incubation time on egg recovery. Regardless of treatment, ~94% of the recovered eggs at day 35 p.i. contained a slender larva that resembled the infective stage. The mean \pm S.E. moisture loss in the sterilised soil ($28 \pm 1.8\%$) was slightly but significantly higher than in the non-sterilised soil ($21 \pm 1.4\%$) ($F_{(1, 64)} = 8.53$, $P = 0.0048$).

Clinical observations and performance

On day 0 p.f.e., the overall mean live weight of hens in the four groups was 1.53–1.60 kg with no significant effect of soil sterility ($F_{(1, 106)} = 2.29$, $P = 0.1330$), fungal treatment ($F_{(1, 106)} = 1.91$, $P = 0.1701$) and their interaction ($F_{(1, 106)} = 0.82$, $P = 0.3684$). By day 36 p.f.e., the overall mean weight gain of hens in the four groups was -76 to 90 g, but there was no significant effect of soil sterility ($F_{(1, 106)} = 3.39$, $P = 0.0686$), fungal treatment ($F_{(1, 106)} = 0.04$, $P = 0.8367$) and their interaction ($F_{(1, 106)} = 2.72$, $P = 0.1022$). Most hens started laying eggs from days 3–7 p.f.e. During the study, the hens showed no overt signs of illness but two hens from one of the three NC subgroups died, possibly due to cannibalism.

Worm burdens

The overall mean worm burdens of *A. galli* and *H. gallinarum* in hens in the four groups are shown in Fig. 2a and b, respectively. All 118 hens were *A. galli* positive, while 115 birds were *H. gallinarum* positive. The left and right spicules of *H. gallinarum* males had a mean \pm SE length of $2086 \pm 26 \mu\text{m}$ (range: 1554–2417 μm) and $723 \pm 8 \mu\text{m}$ (range: 402–850 μm), respectively. The interaction between the soil sterility and fungal treatment strongly influenced the total ascarid worm burden ($F_{(1, 106)} = 100.38$, $P < 0.0001$) and the individual worm burdens of both *A. galli* ($F_{(1, 106)} = 96.85$, $P < 0.0001$) and *H. gallinarum* ($F_{(1, 106)} = 10.07$, $P = 0.0020$). Group SF hens thus had significantly lower worm burdens of both *A. galli* ($P < 0.0001$ in all cases) and *H. gallinarum* ($P \leq 0.0001$ in all cases) compared to the three other groups that all had comparable *A. galli* ($P > 0.3120$ in all cases) and *H. gallinarum* worm burdens ($P > 0.9989$ in all cases) (Fig. 2). *Heterakis gallinarum* represented 6% of the total ascarids





for SF hens and 2–3% for the three other groups. With reference to the estimated cumulative egg dose of 2214 (SC), 208 (SF), 1670 (NC) and 1281 eggs (NF) that each hen was theoretically exposed to on four exposures, the overall establishments of total ascarid were 20% (SC), 36% (SF), 21% (NC) and 33% (NF).

Parasite population composition

The overall mean proportion of *A. galli* recovered from the intestinal lumen and intestinal wall is shown in Fig. 3. In general, *A. galli* were more prevalent in the intestinal lumen (61–78%) than in the intestinal tissue (22–39%). Fungal treatment had a significant effect on the relative distribution of tissue phase and luminal phase *A. galli* ($F_{(1, 106)} = 9.85, P = 0.0022$). This resulted in a significantly higher proportion ($37 \pm 2\%$, mean \pm S.E.) of tissue phase *A. galli* in hens not exposed to fungal treatments compared to the hens exposed to fungal treatments ($27 \pm 2\%$). There were no significant effects of soil sterility ($F_{(1, 106)} = 3.05, P = 0.0660$) as well as the interaction between fungal treatment and soil sterility ($F_{(1, 106)} = 0.50, P = 0.4815$) on the *A. galli* distribution between intestinal lumen and tissue.

The proportion of *A. galli* (of the total *A. galli* worm burden) within each length category was significantly related to the interaction between soil sterility and fungal treatment (< 0.5 cm: $F_{(1, 106)} = 29.48, P < 0.0001$; 0.5–1.5 cm: $F_{(1, 106)} = 9.89, P = 0.0022$; 1.5–3.0 cm: $F_{(1, 106)} = 29.98, P < 0.0001$; 3.0–5.0 cm: $F_{(1, 106)} = 16.58, P < 0.0001$; 5.0–8.0 cm: $F_{(1, 106)} = 7.24, P = 0.0083$) (Fig. 4). Compared to the groups SC, NC and NF hens, the group SF hens



hosted a significantly lower proportion of *A. galli* < 0.5 cm ($P < 0.0001$ in all cases) and significantly higher proportions of the three largest length categories ($P < 0.0225$ in all cases, except $P = 0.0508$ for SF vs NF in the category 5.0–8.0 cm). The SC, NC and NF hens hosted nearly equal proportions of all five *A. galli* length categories ($P > 0.0625$ in all cases, except $P = 0.0076$ for SC vs NC in the category 1.5–3.0 cm). Irrespective of group, all tissue phase *A. galli* larvae were < 0.5 cm (~ 0.5 mm). In groups SC, NC and NF hens, the luminal *A. galli* worms within the category < 0.5 cm (i.e. 5 mm) were approximately 0.5–1.0 mm whereas those in group SF hens ranged ~ 0.5 –4.9 mm.



For *H. gallinarum*, the highest mean \pm S.E. proportion of worms > 0.5 cm was hosted by the group SF hens ($52 \pm 9\%$) followed by NC ($38 \pm 6\%$), NF ($31 \pm 5\%$) and SC hens ($25 \pm 6\%$). However, the effect of soil sterility, fungal treatment and their interactions on *H. gallinarum* population composition was not possible to analyse using the same statistical model that was used for *A. galli* because many hens had only < 0.5 or ≥ 0.5 cm *H. gallinarum*.

Faecal egg counts

On day -1 p.f.e., all hens were negative for ascarid eggs. On day 40 p.f.e., 3, 57, 7 and 17% hens of groups SC, SF, NC and NF, respectively, had positive EPG. There was a significant interaction between fungal treatment and soil sterility regarding day 40 p.f.e. EPG ($F_{(1, 106)} = 5.17, P < 0.0250$) as the overall mean EPG in group SF hens was significantly higher than in groups SC ($P < 0.0001$) and NC hens ($P = 0.0241$) but comparable to group NF hens ($P = 0.4040$) (Fig. 5).

Ascaridia galli IgY titres

The overall group mean (+ SE) *A. galli* IgY titres in hens in the four groups on days -1 and 36 p.f.e. are shown in Fig. 6. All hens were seropositive at both time-points. The antibody titer was significantly affected by the interaction between sterility of soil, fungal treatment and sampling time ($F_{(4, 212)} = 14.02, P < 0.0001$). On day -1 p.f.e., the group mean \pm SE titres ranged between 734 ± 62 and 1071 ± 115 , with no significant differences between the groups ($P > 0.7380$ in all cases). By day 36 p.f.e., the IgY titre had increased significantly in all groups with an overall 8–11 fold increase in groups SC ($P < 0.0001$), NC ($P < 0.0001$) and NF ($P < 0.0001$), but only three fold increase in group SF hens ($P < 0.0001$). There were no significant correlations ($P > 0.05$) between IgY titre and individual worm burden (total ascarid, total *A. galli*) in all groups except SF where there were significant but weak correlations for the total ascarid ($r_{(30)} = 0.38, P = 0.0362$) and *A. galli* worm burden ($r_{(30)} = 0.40, P = 0.0275$).



Discussion

The present study has for the first time shown that ascarid transmission to hens exposed to egg contaminated soil can be reduced after the soil has been treated with the fungus *P. chlamydosporia*, but only in sterilised soil. The reduced exposure resulted in a higher rate of development into adult worms and thus more patent infections compared to the more heavily infected control hens.

The fungus *P. chlamydosporia* Biotype 10 substantially reduced the egg recovery in the sterilised soil whereas in non-sterilised soil there was no additional effect when compared to the corresponding controls. This limited effect of *P. chlamydosporia* in the non-sterilised soil is in line with previous findings for egg-degrading fungi in general [38, 42, 52, 53]. The currently available literature indicates that native soil biota can reduce the establishment of a newly added fungus [54–58]. This is probably because the new fungus must compete for the soil resources or overcome antagonism by native established soil biota such as other fungi [59–61], bacteria [62–65], protozoa [66], free-living nematodes [67, 68], mites and dipteran larvae [69]. In future studies, application of fungi in nutrient-rich substrates (e.g. rice or barley kernels, decomposed resources etc.) could be explored as this may help increase fungal establishment in soil [55, 70, 71]. Ascarid eggs are sensitive to desiccation [72] and after incubation, we found a slightly higher moisture loss in the sterilised soil compared to the non-sterilised soil. However, results indicate that this had no major impact as the

moisture loss in the control and the fungus-treated sterilised soil was not significantly different and both the highest moisture loss and highest egg recovery from a single Petri dish was found in the sterilised soil.

The differences in soil egg numbers (i.e. exposure levels) after fungal treatment was reflected *in vivo* by parasite burden, establishment rate and population composition within the host. The least exposed group had the lowest ascarid worm burdens but a higher parasite establishment rate compared to the three other groups that were more heavily exposed. Similar findings have been reported for an *A. galli* trickle infection in chickens [32], *H. gallinarum* single infection in red-necked pheasants [73] and *Oesophagostomum dentatum* single and trickle infections in pigs [74–76]. In contrast, Permin et al. [77] found no differences in *A. galli* burdens following a single dose of 100, 500 or 2500 eggs. This may be because they only quantified the luminal worms and many larvae in the two higher dose groups may potentially have been, at least temporarily, arrested in the intestinal mucosa [43, 44]. In the current study, we thus found an increased proportion of tissue phase *A. galli* presumably at the third larval stage (L3) [43, 44] in the three high exposure groups, and larger worms and patent infections primarily in the lowest exposure group. The absence of patent infections in most heavily infected hens supports that faecal egg counts can severely underestimate immature worm burdens and exposure levels [44, 74, 76].

The current results showed that low exposure may at least, in the short-term, lead to mature *A. galli* populations in contrast to predominantly immature infections at higher exposure levels. Reduced exposure and lower worm burdens are both desirable to lower the overall impact of ascarids on chicken health and productivity but seem to favour the establishment of patent infections. Density-dependent worm maturation was previously documented for *H. gallinarum* in chickens and ring-necked pheasants where heavily infected birds hosted significantly smaller female worms compared to lightly infected birds [73, 78]. It is unknown if, given time, at least some of our arrested larvae, presumably L3 in the intestinal tissue and L4 in the intestinal lumen [1, 43], would have reached maturity as our hens were only followed for 30 days after the last exposure. However, Ikeme [44] found the development of nearly all L3 to be arrested for up to 13 weeks post-last-exposure in birds that received a high infection dose. It is therefore very important to use sensitive recovery techniques to minimize the risk of overlooking high immature worm burdens. The precise mechanisms responsible for density-dependent effects are not fully understood. The combination of intraspecific competition among worms for limited resources (e.g. space, nutrients) in the host gut

and the effect of host immune responses on parasite population seem important [79].

The above findings highlight the basic complication of any control strategy that cannot inactivate all parasite eggs in the environment. Initially there may be a lowered impact on the hosts present at the time due to lowered exposure, but if the result is associated with altered infection dynamics, and thus an earlier onset of patency, environmental recontamination might be higher than if there had been no intervention. This is further complicated as freshly deposited eggs take weeks to months to develop to infectivity depending on weather and season [30]. This goes to show that designing and implementing control strategies on a farm must take parasite biology and ecology into account to not only offer temporary relief, but also be effective long-term.

There is a close phylogenetic relationship between *A. galli* and *Heterakis* spp. [80] with corresponding production of cross-reacting antibodies [81]. However, the current contribution in the IgY titre due to *H. gallinarum* is expected to be negligible due to the much lower worm burdens compared to *A. galli*. The individual antibody levels appeared to increase with increasing exposure level and worm burden. However, individual antibody levels seemed uninvolved in any immune-related short-term regulation of *A. galli* populations. This is in agreement with previous findings of a very weak or a complete lack of correlation between systemic/egg-yolk IgY level and *A. galli*/*H. gallinarum* worm burden [3, 81]. A similar lack of association between porcine blood IgG level and worm burdens has been reported for *Ascaris suum* [82, 83] and *Trichuris suis* [84]. Furthermore, we also found that previous exposure did not protect against subsequent *A. galli* reinfection, which is in line with other studies [32, 85]. Others have reported increased mRNA expression of Th2 cytokines IL-4 and IL-13 in the intestinal tissues and spleen of *A. galli* infected hosts [3, 86, 87]. Both cytokines play a role in mediating protective immunity against several helminth parasites [88, 89] but it appears that *A. galli* may evade host immune responses to avoid expulsion as suggested for *O. dentatum* in pigs [90, 91]. This could be a reason why *A. galli* prevalence in laying hens kept in non-cage systems (barn, free-range and organic) seems to increase over time during an egg laying period of approximately one year [34].

To the best of our knowledge, there are no optimal/standardized protocols to establish patent *A. galli* infections in chickens. Experimental infection procedures vary greatly in relation to infection material (source, embryonation medium, temperature and duration of embryonation) and host factors (age, breed, etc.) [10, 32, 92–96]. This makes it extremely difficult to compare results between different studies. Many experiments performed earlier by our group could

not establish patent *A. galli* infections when chickens were either infected with a single dose of 500 eggs [97, 98] or trickle infected twice weekly with 25–100 eggs per infection over a period of six weeks [32, 85]. The lowest infection dose used in the latter studies is very similar to the lowest exposure level of the current study. We have therefore made some modifications in the current protocol in relation to the previous failures. Hens were exposed to ascarid eggs at 20–22 weeks of age. This corresponded to the period when most hens started to lay eggs and it has been suggested that hens during this period are more susceptible and may have an increased establishment of *A. galli* due to hormonal changes in the birds [92]. Furthermore, we embryonated (i.e. incubated) ascarid eggs at 22 °C for only five weeks compared to the six week incubation protocol in the earlier studies. This was because chicken ascarid eggs develop fully within four weeks of incubation at 22 °C [29] and we thus provided only one additional week for the developed eggs to mature to infectivity.

Conclusions

The Biotype 10 strain of *P. chlamydosporia* was only effective in inactivating ascarid eggs and thereby reducing the infection level of sterilised soil and thus total worm burdens of the exposed hens. The consequence was that the proportion of early mature/mature *A. galli* increased and faecal eggs counts were higher than in all the other groups where hens were exposed to a higher number of infective eggs in the soil. This underlines an inherent dilemma and complexity of ascarid control in that hosts may suffer less short-term, but current reduced exposure may lead to long-term higher environmental re-contamination if not all eggs can be eliminated. This needs to be considered in future biological and other control strategies in poultry.

Abbreviations

EPG: eggs per gram of faeces; FEC: faecal egg counts; SE: standard error

Acknowledgements

The authors gratefully acknowledge Rothamsted Research Ltd. (United Kingdom) for kindly providing *Pochonia chlamydosporia* Biotype 10. We are very thankful to Lise-Lotte Christensen, Suraj Dhakal, Lene R. Dal, Sara Almeida, Sophie Stolzenbach, Tina V. A. Hansen and Eline P. Hansen for the technical assistance.

Funding

This research and the PhD fellowship of ST were financed by the Faculty of Health and Medical Sciences, University of Copenhagen, Denmark.

Availability of data and materials

The datasets generated and analysed in the current study can be available from the corresponding author upon reasonable request.

Authors' contributions

ST, SMT, HM and NVM designed the study. ST, RW, HM and TSD conducted the study. ST analysed the data and drafted the manuscript. All authors discussed the results, contributed in the manuscript and read and approved the final manuscript.

Ethics approval

The animal experiment was approved by the Animal Experiments Inspectorate, The Danish Ministry of Food, Agriculture and Fisheries (Permit Number: 2015-15-0201-00760). The animals were treated according to the Danish ethical guidelines.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Section for Parasitology and Aquatic Pathobiology, Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Dyrhøjevej 100, 1870 Frederiksberg C, Denmark. ²College of Veterinary Medicine, Inner Mongolia Agricultural University, Hohhot 010018, People's Republic of China. ³Section for Organismal Biology, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark. ⁴Section for Immunology and Microbiology, Department of Animal Science, Aarhus University, Blichers Allé 20, Building P25, 3334, 8830 Tjele, Denmark. ⁵Section for Diagnostics and Scientific Advice, National Veterinary Institute, Technical University of Denmark, Kemitorvet, 2800 Kgs. Lyngby, Denmark.

Received: 15 February 2018 Accepted: 14 May 2018

Published online: 29 May 2018

References

- Ikeme MM. Observations on the pathogenicity and pathology of *Ascaridia galli*. *Parasitology*. 1971;63:169–79.
- Ramadan HH, Abou Znada NY. Some pathological and biochemical studies on experimental ascariasis in chickens. *Nahrung*. 1991;35:71–84.
- Schwarz A, Gaulty M, Abel H, Daş G, Humburg J, Rohn K, et al. Immunopathogenesis of *Ascaridia galli* infection in layer chicken. *Dev Comp Immunol*. 2011;35:774–84.
- Reid WM, Carmon JL. Effects of numbers of *Ascaridia galli* in depressing weight gains in chicks. *J Parasitol*. 1958;44:183–6.
- Kilpinen O, Roepstorff A, Permin A, Nørgaard-Nielsen G, Lawson LG, Simonsen HB. Influence of *Dermanyssus gallinae* and *Ascaridia galli* infections on behaviour and health of laying hens (*Gallus gallus domesticus*). *Br Poult Sci*. 2005;46:26–34.
- Phiri IK, Phiri AM, Ziela M, Chota A, Masuku M, Monrad J. Prevalence and distribution of gastrointestinal helminths and their effects on weight gain in free-range chickens in Central Zambia. *Trop Anim Health Prod*. 2007;39:309–15.
- Skallerup P, Luna LA, Johansen MV, Kyvsgaard NC. The impact of natural helminth infections and supplementary protein on growth performance of free-range chickens on smallholder farms in El Sauce, Nicaragua. *Prev Vet Med*. 2005;69:229–44.
- Gaulty M, Duss C, Erhardt G. Influence of *Ascaridia galli* infections and anthelmintic treatments on the behaviour and social ranks of laying hens (*Gallus gallus domesticus*). *Vet Parasitol*. 2007;146:271–80.
- Hørning G, Rasmussen S, Permin A, Bisgaard M. Investigations on the influence of helminth parasites on vaccination of chickens against Newcastle disease virus under village conditions. *Trop Anim Health Prod*. 2003;35:415–24.
- Pleidrup J, Dalgaard TS, Norup LR, Permin A, Schou TW, Skovgaard K, et al. *Ascaridia galli* infection influences the development of both humoral and cell-mediated immunity after Newcastle disease vaccination in chickens. *Vaccine*. 2014;32:383–92.
- Dahl C, Permin A, Christensen JP, Bisgaard M, Muhairwa AP, Petersen KMD, et al. The effect of concurrent infections with *Pasteurella multocida* and *Ascaridia galli* on free range chickens. *Vet Microbiol*. 2002;86:313–24.
- Reid WM, Mabon JL, Harshbarger WC. Detection of worm parasites in chicken eggs by candling. *Poult Sci*. 1973;52:2316–24.
- Fioretti DP, Veronesi F, Diaferia M, Franciosini MP, Proietti PC. *Ascaridia galli*: a report of erratic migration. *Ital J Anim Sci*. 2005;4:310–2.

14. Lee DL. The structure and development of *Histomonas meleagridis* (Mastigamoebidae: Protozoa) in the female reproductive tract of its intermediate host, *Heterakis gallinarum* (Nematoda). *Parasitology*. 1969;59:877–84.
15. Windisch M, Hess M. Experimental infection of chickens with *Histomonas meleagridis* confirms the presence of antibodies in different parts of the intestine. *Parasite Immunol*. 2010;32:29–35.
16. McDougald LR. Blackhead disease (histomoniasis) in poultry: a critical review. *Avian Dis*. 2005;49:462–76.
17. Dolka B, Żbikowski A, Dolka I, Szeleszczuk P. Histomonosis - an existing problem in chicken flocks in Poland. *Vet Res Commun*. 2015;39:189–95.
18. Grafl B, Liebhart D, Windisch M, Ibesich C, Hess M. Seroprevalence of *Histomonas meleagridis* in pullets and laying hens determined by ELISA. *Vet Rec*. 2011;168:160.
19. Stokholm NM, Permin A, Bisgaard M, Christensen JP. Causes of mortality in commercial organic layers in Denmark. *Avian Dis*. 2010;54:1241–50.
20. Esquent C, De Herdt P, De Bosschere H, Ronsmans S, Ducatelle R, Van Erum J. An outbreak of histomoniasis in free-range layer hens. *Avian Pathol*. 2003;32:305–8.
21. Thapa S, Hinrichsen LK, Brenninkmeyer C, Gunnarsson S, Heerkens JLT, Verwer C, et al. Prevalence and magnitude of helminth infections in organic laying hens (*Gallus gallus domesticus*) across Europe. *Vet Parasitol*. 2015;214:118–24.
22. Jansson DS, Nyman A, Vågsholm I, Christensson D, Göransson M, Fossum O, et al. Ascarid infections in laying hens kept in different housing systems. *Avian Pathol*. 2010;39:525–32.
23. Kaufmann F, Daş G, Sohnrey B, Gauly M. Helminth infections in laying hens kept in organic free range systems in Germany. *Livest Sci*. 2011;141:182–7.
24. Ackert JE. The morphology and life history of the fowl nematode *Ascaridia lineata* (Schneider). *Parasitology*. 1931;23:360–79.
25. Clapham PA. On the life-history of *Heterakis gallinae*. *J Helminthol*. 1933;11:67–86.
26. Wharton D. Nematode egg-shells. *Parasitology*. 1980;81:447–63.
27. Christensen RO, Earle HH, Butler RL, Creel HH. Studies on the eggs of *Ascaridia galli* and *Heterakis gallinae*. *Trans Am Microsc Soc*. 1942;61:191.
28. Farr MM. Further observations on the survival of the protozoan parasite, *Histomonas meleagridis* and eggs of poultry nematodes in feces of infected birds. *Cornell Vet*. 1961;51:3–13.
29. Thapa S, Thamsborg SM, Meyling NV, Dhakal S, Mejer H. Survival and development of chicken ascarid eggs in temperate pastures. *Parasitology*. 2017;144:1243–52.
30. Committee for Medicinal Products for Veterinary Use. Flubendazole (extrapolation to poultry), summary report (4). In: EMEA/CVMP/33128/2006-Final; 2006. http://www.ema.europa.eu/docs/en_GB/document_library/Maximum_Residue_Limits_-_Report/2009/11/WC500014292.pdf. Accessed 21 Mar 2017.
31. Committee for Medicinal Products for Veterinary Use. CVMP assessment report for Panacur AquaSol (EMEA/V/C/002008/X/00003). 2014. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Assessment_Report_-_Variation/veterinary/002008/WC500165605.pdf. Accessed 21 Mar 2017.
32. Ferdusy T, Luna-Olivares LA, Nejsum P, Thamsborg SM, Kyvsgaard NC. The use of genetically marked infection cohorts to study changes in establishment rates during the time course of a repeated *Ascaridia galli* infection in chickens. *Int J Parasitol*. 2015;45:393–8.
33. Tarbiat B, Jansson DS, Tydén E, Höglund J. Comparison between anthelmintic treatment strategies against *Ascaridia galli* in commercial laying hens. *Vet Parasitol*. 2016;226:109–15.
34. Zloch A, Kuchling S, Hess M, Hess C. Influence of alternative husbandry systems on postmortem findings and prevalence of important bacteria and parasites in layers monitored from end of rearing until slaughter. *Vet Rec*. 2018;182:350.
35. Braga FR, De Araújo JV. Nematophagous fungi for biological control of gastrointestinal nematodes in domestic animals. *Appl Microbiol Biotechnol*. 2014;98:71–82.
36. Domsch KH, Gams W, Anderson T-H. Compendium of soil fungi, Volume 1. 1st ed. London: Academic Press; 1980.
37. Lýsek H, Fassatióvá O, Cuervo Pineda N, Lorenzo Hernández N. Ovicidal fungi in soils of Cuba. *Folia Parasitol*. 1982;29:265–70.
38. Siddiqui IA, Atkins SD, Kerry BR. Relationship between saprotrophic growth in soil of different biotypes of *Pochonia chlamydsoporia* and the infection of nematode eggs. *Ann Appl Biol*. 2009;155:131–41.
39. Manzanilla-López HR, Esteves I, Finetti-Sialer MM, Hirsch PR, Ward E, Devonshire J, et al. *Pochonia chlamydsoporia*: advances and challenges to improve its performance as a biological control agent of sedentary endo-parasitic nematodes. *J Nematol*. 2013;45:1–7.
40. Thapa S, Meyling NV, Katakam KK, Thamsborg SM, Mejer H. A method to evaluate relative ovicidal effects of soil microfungi on thick-shelled eggs of animal-parasitic nematodes. *Biocontrol Sci Technol*. 2015;25:756–67.
41. Braga FR, Araújo JV, Araujo JM, Frassy LN, Tavela AO, Soares FEF, et al. *Pochonia chlamydsoporia* fungal activity in a solid medium and its crude extract against eggs of *Ascaridia galli*. *J Helminthol*. 2012;86:348–52.
42. Thapa S, Mejer H, Thamsborg SM, Lekfeldt JDS, Wang R, Jensen B, et al. Survival of chicken ascarid eggs exposed to different soil types and fungi. *Appl Soil Ecol*. 2017;121:143–51.
43. Herd RP, McNaught DJ. Arrested development and the histotropic phase of *Ascaridia galli* in the chicken. *Int J Parasitol*. 1975;5:401–6.
44. Ikeme MM. Effects of different levels of nutrition and continuing dosing of poultry with *Ascaridia galli* eggs on the subsequent development of parasite populations. *Parasitology*. 1971;63:233–50.
45. Poulsen PHB, Al-Soud WA, Bergmark L, Magid J, Hansen LH, Sørensen SJ. Effects of fertilization with urban and agricultural organic wastes in a field trial - prokaryotic diversity investigated by pyrosequencing. *Soil Biol Biochem*. 2013;57:784–93.
46. López-Rayó S, Laursen KH, Lekfeldt JDS, Delle Grazie F, Magid J. Long-term amendment of urban and animal wastes equivalent to more than 100 years of application had minimal effect on plant uptake of potentially toxic elements. *Agric Ecosyst Environ*. 2016;231:44–53.
47. Ferdusy T, Nejsum P, Roepstorff A, Thamsborg SM, Kyvsgaard NC. *Ascaridia galli* in chickens: intestinal localization and comparison of methods to isolate the larvae within the first week of infection. *Parasitol Res*. 2012;111:2273–9.
48. Lapage G. Mönning's veterinary helminthology and entomology. London: Bailliere, Tindall and Cox; 1956.
49. Soulsby E. Textbook of veterinary clinical parasitology, Volume 1. Helminths. Oxford: Blackwell Scientific Publications; 1965.
50. Roepstorff A, Nansen P. Epidemiology, diagnosis and control of helminth parasites of swine. *FAO Animal Health Manual*. Rome: Food and Agriculture Organization of the United Nations; 1998.
51. Norup LR, Dalgaard TS, Pleidrup J, Permin A, Schou TW, Jungersen G, et al. Comparison of parasite-specific immunoglobulin levels in two chicken lines during sustained infection with *Ascaridia galli*. *Vet Parasitol*. 2013;191:187–90.
52. Bourne JM, Kerry BR, De Leij FAAM. Methods for the study of *Verticillium chlamydsoporium* in the rhizosphere. *J Nematol*. 1994;26:587–91.
53. Monfort E, Lopez-Llorca LV, Jansson HB, Salinas J. *In vitro* soil receptivity assays to egg-parasitic nematophagous fungi. *Mycol Prog*. 2006;5:18–23.
54. Lockwood JL, Lingappa BT. Fungitoxicity of sterilised soil inoculated with soil microflora. *Phytopathology*. 1963;58:917–20.
55. Lockwood JL. Fungistasis in soils. *Biol Rev*. 1977;52:1–43.
56. Toyota K, Ritz K, Young IM. Microbiological factors affecting the colonisation of soil aggregates by *Fusarium oxysporum* f. sp. *raphani*. *Soil Biol Biochem*. 1996;28:1513–21.
57. Mauchline TH, Kerry BR, Hirsch PR. Quantification in soil and the rhizosphere of the nematophagous fungus *Verticillium chlamydsoporium* by competitive PCR and comparison with selective plating. *Appl Environ Microbiol*. 2002;68:1846–53.
58. de Boer W, Verheggen P, Klein Gunnewiek PJ, Kowalchuk G, van Veen JA. Microbial community composition affects soil fungistasis. *Appl Environ Microbiol*. 2003;69:835–44.
59. Chen SY, Chen FJ. Fungal parasitism of *Heterodera glycines* eggs as influenced by egg age and pre-colonization of cysts by other fungi. *J Nematol*. 2003;35:271–7.
60. Howell CR. Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: The history and evolution of current concepts. *Plant Dis*. 2003;87:4–10.
61. Harman GE, Howell CR, Viterbo A, Chet I, Lorito M. *Trichoderma* species-opportunistic, avirulent plant symbionts. *Nat Rev Microbiol*. 2004;2:43–56.
62. Lopez Llorca LV, Boag B. Inhibition of *Verticillium suchlasporium* and other nematophagous fungi by bacteria colonising *Heterodera avenae* females. *Nematol Mediterr*. 1990;18:233–7.
63. Ballhausen MB, van Veen JA, Hundscheid MPJ, de Boer W. Methods for baiting and enriching fungus-feeding (Mycophagous) rhizosphere bacteria. *Front Microbiol*. 2015;6:1416.

64. Rudnick MB, van Veen JA, de Boer W. Baiting of rhizosphere bacteria with hyphae of common soil fungi reveals a diverse group of potentially mycophagous secondary consumers. *Soil Biol Biochem.* 2015;88:73–82.
65. Li L, Mo M, Qu Q, Luo H, Zhang K. Compounds inhibitory to nematophagous fungi produced by *Bacillus* sp. strain H6 isolated from fungistatic soil. *Eur J Plant Pathol.* 2007;117:329–40.
66. Ekelund F, Rönn R. Notes on protozoa in agricultural soil with emphasis on heterotrophic flagellates and naked amoebae and their ecology. *FEMS Microbiol Rev.* 1994;15:321–53.
67. Ruess L, Dighton J. Cultural studies on soil nematodes and their fungal hosts. *Nematologica.* 1996;42:330–46.
68. Jaffee BA, Muldoon AE, Didden WAM. Enchytraeids and nematophagous fungi in soil microcosms. *Biol Fertil Soils.* 1997;25:382–8.
69. Ruess L, Lussenhop J. Trophic interactions of fungi and animals. In: Dighton J, White FJ, Oudemans P, editors. *The fungal community - its organization and role in the ecosystem.* 3rd ed. Boca Raton: CRC Press; 2005. p. 581–98.
70. Kerry B. The use of microbial agents for the biological control of plant parasitic nematodes. In: Jones DG, editor. *Exploitation of microorganisms.* London: Chapman & Hall; 1993. p. 81–104.
71. Luambano ND, Manzanilla-López RH, Kimenju JW, Powers SJ, Narla RD, Wanjohi WJ, et al. Effect of temperature, pH, carbon and nitrogen ratios on the parasitic activity of *Pochonia chlamydosporia* on *Meloidogyne incognita*. *Biol Control.* 2015;80:23–9.
72. Tarbiat B, Jansson DS, Höglund J. Environmental tolerance of free-living stages of the poultry roundworm *Ascaridia galli*. *Vet Parasitol.* 2015;209:101–7.
73. Tompkins DM, Hudson PJ. Regulation of nematode fecundity in the ring-necked pheasant (*Phasianus colchicus*): not just density dependence. *Parasitology.* 1999;118:417–23.
74. Christensen CM, Barnes EH, Nansen P, Roepstorff A, Slotved HC. Experimental *Oesophagostomum dentatum* infections in the pig: Worm populations at regular intervals during trickle infections with three dose levels of larvae. *Int J Parasitol.* 1995;25:1491–8.
75. Roepstorff A, Nilsson O, Oksanen A, Gjerde B, Richter SH, Örtengren E, et al. Intestinal parasites in swine in the Nordic countries: prevalence and geographical distribution. *Vet Parasitol.* 1998;76:305–19.
76. Roepstorff A, Bjørn H, Nansen P, Barnes EH, Christensen CM. Experimental *Oesophagostomum dentatum* infections in the pig: worm populations resulting from trickle infections with three dose levels of larvae. *Int J Parasitol.* 1996;26:399–408.
77. Permin A, Bojesen M, Nansen P, Bisgaard M, Frandsen F, Pearman M. *Ascaridia galli* populations in chickens following single infections with different dose levels. *Parasitol Res.* 1997;83:614–7.
78. Daş G, Gaulty M. Density related effects on lifetime fecundity of *Heterakis gallinarum* in chickens. *Parasit Vectors.* 2014;7:334.
79. Keymer A. Density-dependent mechanisms in the regulation of intestinal helminth populations. *Parasitology.* 1982;84:573–87.
80. Wang B-J, Gu X-B, Yang G-Y, Wang T, Lai W-M, Zhong Z-J, et al. Mitochondrial genomes of *Heterakis gallinae* and *Heterakis beramporia* support that they belong to the infraorder Ascaridomorpha. *Infect Genet Evol.* 2016;40:228–35.
81. Daş G, Hennies M, Sohnrey B, Rahimian S, Wongrak K, Stehr M, et al. A comprehensive evaluation of an ELISA for the diagnosis of the two most common ascarids in chickens using plasma or egg yolks. *Parasit Vectors.* 2017;10:187.
82. Lind P, Eriksen L, Nansen P, Nilsson O, Roepstorff A. Response to repeated inoculations with *Ascaris suum* eggs in pigs during the fattening period. II. Specific IgA, IgG, and IgM antibodies determined by enzyme-linked immunosorbent assay. *Parasitol Res.* 1993;79:240–4.
83. Roepstorff A, Eriksen L, Slotved HC, Nansen P. Experimental *Ascaris suum* infection in the pig: worm population kinetics following single inoculations with three doses of infective eggs. *Parasitology.* 1997;115:443–52.
84. Nejsum P, Thamsborg SM, Petersen HH, Kringel H, Fredholm M, Roepstorff A. Population dynamics of *Trichuris suis* in trickle-infected pigs. *Parasitology.* 2009;136:691–7.
85. Ferdushy T, Schou TW, Norup LR, Dalgaard TS, Thamsborg SM, Nejsum P, et al. Acquisition of resistance after continuous infection with *Ascaridia galli* in chickens. *Parasitology.* 2014;141:1603–10.
86. Degen WGJ, Van Daal N, Rothwell L, Kaiser P, Schijns VEJC. Th1/Th2 polarization by viral and helminth infection in birds. *Vet Microbiol.* 2005;105:163–7.
87. Dalgaard TS, Skovgaard K, Norup LR, Pleidrup J, Permin A, Schou TW, et al. Immune gene expression in the spleen of chickens experimentally infected with *Ascaridia galli*. *Vet Immunol Immunopathol.* 2015;164:79–86.
88. Anthony RM, Rutitzky LI, Urban JF, Stadecker MJ, Gause WC. Protective immune mechanisms in helminth infection. *Nat Rev Immunol.* 2007;7:975–87.
89. Finkelman FD, Wynn TA, Donaldson DD, Urban Jr JF. The role of IL-13 in helminth-induced inflammation and protective immunity against nematode infections. *Curr Opin Immunol.* 1999;11:420–6.
90. Andreassen A, Petersen HH, Kringel H, Iburg TM, Skovgaard K, Dawson H, et al. Immune and inflammatory responses in pigs infected with *Trichuris suis* and *Oesophagostomum dentatum*. *Vet Parasitol.* 2015;207:249–58.
91. Andreassen A, Skovgaard K, Klaver EJ, van Die I, Mejer H, Thamsborg SM, et al. Comparison of innate and Th1-type host immune responses in *Oesophagostomum dentatum* and *Trichuris suis* infections in pigs. *Parasite Immunol.* 2016;38:53–63.
92. Gaulty M, Homann T, Erhardt G. Age-related differences of *Ascaridia galli* egg output and worm burden in chickens following a single dose infection. *Vet Parasitol.* 2005;128:141–8.
93. Permin A, Nansen P, Bisgaard M, Frandsen F. *Ascaridia galli* infections in free-range layers fed on diets with different protein contents. *Br Poult Sci.* 1998;39:441–5.
94. Rahimian S, Daş G, Gaulty M. Maternal protection against *Ascaridia galli*? *Vet Parasitol.* 2017;233:43–7.
95. Ruhnke I, Andronicos NM, Swick RA, Hine B, Sharma N, Kheravii SK, et al. Immune responses following experimental infection with *Ascaridia galli* and necrotic enteritis in broiler chickens. *Avian Pathol.* 2017;46:602–9.
96. Marcos-Atxutegi C, Gandolfi B, Arangüena T, Sepúlveda R, Arévalo M, Simón F. Antibody and inflammatory responses in laying hens with experimental primary infections of *Ascaridia galli*. *Vet Parasitol.* 2009;161:69–75.
97. Ferdushy T, Luna-Olivares LA, Nejsum P, Roepstorff AK, Thamsborg SM, Kyvsgaard NC. Population dynamics of *Ascaridia galli* following single infection in young chickens. *Parasitology.* 2013;140:1078–84.
98. Luna-Olivares LA, Kyvsgaard NC, Ferdushy T, Nejsum P, Thamsborg SM, Roepstorff A, et al. The jejunal cellular responses in chickens infected with a single dose of *Ascaridia galli* eggs. *Parasitol Res.* 2015;114:2507–15.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://www.biomedcentral.com/submissions)

