



NuSTAR detection of 4s Hard X-ray Lags from the Accreting Pulsar GS 0834-430

Bachetti, Matteo; Miyasaka, Hiromasa; Harrison, Fiona; Fürst, Felix; Barret, Didier; Bellm, Eric C.; Boggs, Steven E.; Chakrabarty, Deepto; Chenevez, Jérôme; Christensen, Finn Erland

Published in:
E P J Web of Conferences

Link to article, DOI:
[10.1051/epjconf/20136406011](https://doi.org/10.1051/epjconf/20136406011)

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Bachetti, M., Miyasaka, H., Harrison, F., Fürst, F., Barret, D., Bellm, E. C., ... Zhang, W. (2014). NuSTAR detection of 4s Hard X-ray Lags from the Accreting Pulsar GS 0834-430. *E P J Web of Conferences*, 64, [06011]. <https://doi.org/10.1051/epjconf/20136406011>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

***NuSTAR* detection of 4s Hard X-ray Lags from the Accreting Pulsar GS 0834-430**

Matteo Bachetti^{1,2,a}, Hiromasa Miyasaka³, Fiona Harrison³, Felix Fürst³, Didier Barret^{1,2}, Eric C. Bellm³, Steven E. Boggs⁴, Deepto Chakrabarty⁵, Jerome Chenevez⁶, Finn E. Christensen⁶, William W. Craig^{4,7}, Brian W. Grefenstette³, Charles J. Hailey⁸, Kristin K. Madsen³, Lorenzo Natalucci⁹, Katja Pottschmidt¹⁰, Daniel Stern¹¹, John A. Tomsick⁴, Dominic J. Walton³, Jörn Wilms¹², and William Zhang¹³

¹ *Université de Toulouse; UPS-OMP; IRAP; Toulouse, France*

² *CNRS; Institut de Recherche en Astrophysique et Planétologie; 9 Av. colonel Roche, BP 44346, F-31028 Toulouse cedex 4, France*

³ *Cahill Center for Astronomy and Astrophysics, California Institute of Technology, Pasadena, CA 91125*

⁴ *Space Sciences Laboratory, University of California, Berkeley, CA 94720, USA*

⁵ *Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139*

⁶ *DTU Space, National Space Institute, Technical University of Denmark, Elektrovej 327, DK-2800 Lyngby, Denmark*

⁷ *Lawrence Livermore National Laboratory, Livermore, CA 94550, USA*

⁸ *Columbia Astrophysics Laboratory, Columbia University, New York, NY 10027, USA*

⁹ *Istituto di Astrofisica e Planetologia Spaziali, INAF, Via Fosso del Cavaliere 100, Roma I-00133, Italy*

¹⁰ *CRESST, UMBC, and NASA GSFC, Code 661, Greenbelt, MD 20771, USA*

¹¹ *Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA*

¹² *Dr. Karl-Remeis-Sternwarte and ECAP, Sternwartstr. 7, 96 049 Bamberg, Germany*

¹³ *NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA*

Abstract. The *NuSTAR* hard X-ray telescope observed the transient Be/X-ray binary GS 0834–430 during its 2012 outburst. The source is detected between 3 – 79 keV with high statistical significance, and we were able to perform very accurate spectral and timing analysis. The phase-averaged spectrum is consistent with that observed in many other magnetized accreting pulsars. We fail to detect cyclotron resonance scattering features in either phase-averaged nor phase-resolved spectra that would allow us to constrain the pulsar’s magnetic field. We detect a pulse period of ~ 12.29 s in all energy bands. The pulse profile can be modeled with a double Gaussian and shows a strong and smooth hard lag of up to 0.3 cycles in phase, or about 4s between the pulse at ~ 3 and $\gtrsim 30$ keV. This is the first report of such a strong lag in high-mass X-ray binary (HMXB) pulsars. Previously reported lags have been significantly smaller in phase and restricted to low-energies ($E < 10$ keV). We investigate the possible mechanisms that might produce such lags. We find the most likely explanation for this effect to be a complex beam geometry.

^ae-mail: matteo.bachetti@irap.omp.eu




Figure 1. (Left) Evolution of the pulse profile with energy. Starting from an energy resolution of ~ 0.4 keV and 48 phase bins, we averaged over an increasing number of bins at higher energies, following approximately a geometrical progression both in phase and in energy in order to attain a better S/N. Each profile is background-subtracted and normalized from 0 to 1. This normalized rate is color-coded, ranging from the lowest count-rates in black to the highest in white. The profiles are repeated twice in phase for clarity. The white dashed lines indicate roughly the phase of the peak and the dip, respectively, at the softest energies to guide the eye. The minimum and the peak of the profiles are shifting in phase with energy, showing a clear hard lag. (Right) Phase lags of the two peaks, as obtained by fitting each rebinned profile with two independent Gaussians, relative to the phase of the first peak in the lowest energy band. Black is the main peak, red the secondary. Full-color error bars are $1 - \sigma$ uncertainties, while light-colored error bars indicate the HWHM of the Gaussians. Dashed lines show the phase of the two peaks in the lowest energy band. The dash-dotted line indicates the fit of a Comptonization lag model to the secondary peak lags (see the details in [6]). The insets show the fit at 3 keV, at ~ 35 keV where the secondary peak becomes dominant, and at ~ 60 keV respectively.

1 NuSTAR observation of GS 0834–430

GS 0834–430 is an accreting Pulsar discovered in 1990 [1]. It is an example of Be/X-ray binary. It has a spin period of ~ 13 s, an orbital period of 108d with a wide orbit ($a_x \sin i \sim 128 l - s$) [2] around a B0-2 V-IIIe companion [3]. At the peak of its latest outburst [4], on UT 2012 July 11, the source was observed by NuSTAR [5] for about 31ks. In these pointings, at 10 keV the source is 10^3 times brighter than the background, and at 79 keV the source is still 10 times above the background. We could perform very high quality timing and spectral analysis up to 79 keV.

The pulse profile is detected significantly up to ~ 60 keV. The energy-resolved pulse profile shows clear phase lags between the pulse profiles at different energies. The shift is about 0.3 cycles in phase, or 4 s in time, between 3 and 30 keV. Fitting the profile with a double Gaussian, the two components seem to move together, almost linearly at first, then saturating above 20 keV (Fig. 1). The primary peak dominates at low energy, then the second becomes the stronger one above 30 keV.

The source spectrum varies considerably with pulse phase. We use an NPEX model to characterize the spectrum in 10 phase bins (Fig. 1) of similar hardness ratio. The model fits adequately the spectrum for all phase bins, and there are no signatures that suggest the presence of a cyclotron line. A weak (EW ~ 40 eV) iron line is present, mildly anti-correlated with flux. We find no evidence of cyclotron resonance scattering features (CRSF), often observed in accreting pulsars and yielding an approximate measure of the magnetic field in these sources [see 7, for a review].




Figure 2. Phase-averaged spectrum fit using the FDCUT continuum model. Black points represent data from module FPMA, red from FPMB. The top panel shows the counts spectrum, with the data-to-model ratio for FDCUT and NPEX shown on the middle and bottom panels respectively. The region around the iron line is shown in the inset.

2 Discussion

Time lags are observed in many accreting sources [see 8, for a review]. They can be produced through delays due, e.g., to Comptonization of the signal [e.g. 9], reflection from a medium far from the source [e.g. 10, 11], relativistic effects [e.g. 12], or propagation of signals in a disk [e.g. 13]. But 4s time lags are very difficult to reconcile with the most common geometries proposed for lag production (see the full discussion in [6]).

A CRSF can also produce a similar effect as the one we observe, producing an apparent delay in the spectral emission due to the different contributions of the feature in the various phases of the pulse profile [e.g. 14], but we do not find evidence for CRSFs in the spectrum in the relevant bands.

Similarly, it is possible that the effect is produced by the different emission components coming from the different regions of the accretion column. If the source has a luminosity between $\sim 10^{37}$ erg s^{-1} , as is the case for GS 0834–430 during this outburst, calculations show that a Compton shock is formed in the accretion column, but that the luminosity is not sufficient to produce a radiation dominated shock, which would decelerate the infalling material high above the neutron star surface [15]. In this intermediate regime, a complex beam profile is therefore produced: radiation, after being upscattered to high energies by inverse Compton scattering inside the accretion column, emerges through the walls of the accretion column (fan beam), as well as along the magnetic field lines (pencil beam). The fan and pencil beam components can in principle have different spectral properties. Given the right geometry, this could create the phase-shift effect seen in the *NuSTAR* data through a gradual mixing of a softer and harder continuum. These processes are complicated and strongly dependent on the single source physical structure, and this might explain why only this source shows such strong lags.

If this is the case, our observation represents a very interesting step in the understanding of these processes. More details can be found in [6].

References

- [1] R. Sunyaev, IAU Circ. **5122**, 2 (1990)
- [2] C.A. Wilson, M.H. Finger, B.A. Harmon, D.M. Scott, R.B. Wilson, L. Bildsten, D. Chakrabarty, T.A. Prince, ApJ **479**, 388 (1997)




Figure 3. (Left) The pulse profile in four different energy bands. A strong dependence with energy is clearly visible. Intervals in the top pulse profile show the phases used for phase-resolved spectroscopy. (Right) Hardness ratio between the 20 – 40 keV and 4 – 7 keV bands as a function of pulse phase (top panel). Intervals have been chosen based on evolution of the hardness ratio to fit NPEX model parameters. The lower three panels show the evolution of fit parameters as a function of pulse phase, i.e., the photon index of the negative powerlaw (α), the cutoff energy kT , the normalization of the negative powerlaw component A_n , and the normalization of the positive powerlaw component A_p . The pulse is repeated once for clarity.

- [3] G.L. Israel, S. Covino, S. Campana, V.F. Polcaro, P. Roche, L. Stella, A. di Paola, D. Lazzati, S. Mereghetti, E. Giallongo et al., *MNRAS* **314**, 87 (2000)
- [4] S.P. Drave, V. Sguera, A.J. Bird, M. Goossens, L. Sidoli, A. Bazzano, M. Fiocchi, *The Astronomer’s Telegram* **4218**, 1 (2012)
- [5] F.A. Harrison, W.W. Craig, F.E. Christensen, C.J. Hailey, W.W. Zhang, S.E. Boggs, D. Stern, W.R. Cook, K. Forster, P. Giommi et al., *ApJ* **770**, 103 (2013)
- [6] H. Miyasaka, M. Bachetti, F.A. Harrison, F. Fürst, D. Barret, E.C. Bellm, S.E. Boggs, D. Chakrabarty, J. Chenevez, F.E. Christensen et al., *ApJ* **775**, 65 (2013)
- [7] W. Coburn, P. Kretschmar, I. Kreykenbohm, V.A. McBride, R.E. Rothschild, R. Staubert, J. Wilms, *ASR* **38**, 2747 (2006)
- [8] J. Poutanen, *ASR* **28**, 267 (2001)
- [9] E.P. Liang, P.L. Nolan, *SSRv* **38**, 353 (1984)
- [10] M.M. Basko, R.A. Sunyaev, L.G. Titarchuk, *A&A* **31**, 249 (1974)
- [11] I.M. George, A.C. Fabian, *MNRAS* **249**, 352 (1991)
- [12] W. Cui, E.H. Morgan, L.G. Titarchuk, *ApJL* **504**, L27 (1998)
- [13] Y.E. Lyubarskii, *MNRAS* **292**, 679 (1997)
- [14] C. Ferrigno, M. Falanga, E. Bozzo, P.A. Becker, D. Klochkov, A. Santangelo, *A&A* **532**, 76 (2011)
- [15] P.A. Becker, D. Klochkov, G. Schönherr, O. Nishimura, C. Ferrigno, I. Caballero, P. Kretschmar, M.T. Wolff, J. Wilms, R. Staubert, *A&A* **544**, 123 (2012)