



Evaluation of the Danish cultivated sugarkelp as possible future source of ingredients such as minerals and pigments

Holdt, Susan Løvstad; Silva Marinho, Goncalo; Sloth, Jens Jørgen; Safafar, Hamed; Jakobsen, Jette; Angelidaki, Irini

Published in:

Book of Abstracts. DTU's Sustain Conference 2015

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Holdt, S. L., Silva Marinho, G., Sloth, J. J., Safafar, H., Jakobsen, J., & Angelidaki, I. (2015). Evaluation of the Danish cultivated sugarkelp as possible future source of ingredients such as minerals and pigments. In *Book of Abstracts. DTU's Sustain Conference 2015* [F-5] Lyngby: Technical University of Denmark (DTU).

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Evaluation of the Danish cultivated sugarkelp as possible future source of ingredients such as minerals and pigments

Susan L. Holdt^{*1}, Goncalo S. Marinho², Jens J. Sloth¹, Hamed Safafar¹, Jette Jakobsen¹, Irini Angelidaki²

1: DTU Food; 2: DTU Environment

*suho@food.dtu.dk

The seaweed *Saccharina latissima* (sugarkelp) is cultivated near a fish and mussel farm, Hjarnø Havbrug A/S near Horsens Fjord in Denmark (Figure 1). The sugarkelp is cultivated commercially for the bioremediation (waste water management) of especially nitrogen, and the valuable biomass then sold and used for various purposes such as food or potentially as a feed supplement. Generally, seaweed is known for their nutraceutical application, but also for being able to accumulate e.g. very high iodine concentrations and unwanted heavy metals (Holdt and Kraan, 2011). In this study, the year-round minerals (incl. trace and heavy metals), pigments and vitamins (vit A and E), were analysed to evaluate the nutritional value, possible risks and harvest time of the *S. latissima* biomass for optimized value and application.

Rope cultivated sugarkelp was sampled both in close proximity to a blue mussel and fish farm (IMTA; see Figure 1) and in a reference/control site, both outside Horsens fjord in Denmark. Sugarkelp biomass was measured from 1m rope droppers (n=3) at 2 m depth in 2013-2014 (deployed in February 2013). Biomass was weighed, followed by freeze drying, homogenizing, frozen before further chemical characterization by various methods for the specific analyses of biomass composition.

Surprisingly high concentrations of potassium (K) and calcium (Ca) were found in the sugarkelp, and also the other trace metals Cr, Fe, Mn, Co, P, Na, Zn, and Se were found. The unwanted elements such as Cu, Hg, As, and inorganic arsenic were below legislative threshold values, whereas a few samples of Cd and Pb were problematic in some seasons, but not considering the recommended daily intake. The iodine was found in so high levels (up to 5 g/kg) that this will be the limiting element for the daily recommended intake of sugarkelp. The pigment profile did not change during the year, however the concentration did, and with fucoxanthin as the most interesting. Generally the year-round variations were due season, and not between the two locations (reference and IMTA), so harvest time are important for optimized use, and may be conflicting with highest yields of sugarkelp.




Fig. 1 Left: Integrated Multi-trophic Aquaculture with the fed organisms (fish), filtrating (mussels) and assimilating (seaweed) organisms combined. Right: Year-round calcium concentration (g/kg dry weight) of the sugarkelp. Levels of calcium from milk and cheese are inserted for comparison (source: www.foodcomp.dk).

Reference: Holdt, S.L., Kraan, S. (2011). Bioactive compounds in seaweed: functional food applications and legislation. *Journal of Applied Phycology*, 23, 543-597