



Complete Genome Sequence of Escherichia coli Strain WG5

Imamovic, Lejla; Misiakou, Maria-Anna; van der Helm, Eric; Panagiotou, Gianni; Muniesa, Maite; Sommer, Morten Otto Alexander

Published in:
Genome Announcements

Link to article, DOI:
[10.1128/genomeA.01403-17](https://doi.org/10.1128/genomeA.01403-17)

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Imamovic, L., Misiakou, M-A., van der Helm, E., Panagiotou, G., Muniesa, M., & Sommer, M. O. A. (2018). Complete Genome Sequence of Escherichia coli Strain WG5. *Genome Announcements*, 6(2), [e01403-17]. <https://doi.org/10.1128/genomeA.01403-17>

General rights


Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Complete Genome Sequence of *Escherichia coli* Strain WG5

Lejla Imamovic,^a Maria-Anna Misiakou,^a Eric van der Helm,^a Gianni Panagiotou,^b  Maite Muniesa,^c Morten Otto Alexander Sommer^a

^aNovo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Lyngby, Denmark

^bLeibniz Institute for Natural Product Research and Infection Biology, Hans Knöll Institute, Jena, Germany

^cDepartment of Genetics, Microbiology and Statistics, Microbiology Section, Faculty of Biology, University of Barcelona, Barcelona, Spain

ABSTRACT *Escherichia coli* strain WG5 is a widely used host for phage detection, including somatic coliphages employed as standard ISO method 10705-1 (2000). Here, we present the complete genome sequence of a commercial *E. coli* WG5 strain.

Escherichia coli is a common commensal of the human and animal gut. *E. coli* strain WG5 is used as a host to detect somatic coliphages, proposed to be an indicator of human and animal fecal contamination of water, sediments, and sludge, as described in ISO method 10705-1 (2000) (1). This is an easily applicable and affordable method for water quality and contamination management in water treatment facilities (2). Other studies employ a similar method using strain WG5 as a host to detect temperate infectious phages from food (3, 4). The applicability of the method is based on the high sensitivity of the *E. coli* WG5 host to infection by somatic coliphages (5). *E. coli* WG5 (6) is a nalidixic acid-resistant mutant of *E. coli* C, also known as strain CN, and is publicly available in the ATCC (ATCC number 700078). *E. coli* strain WG5 possesses an attenuated host restriction-modification system and contains only the core part of the lipopolysaccharide (LPS), increasing susceptibility to phage infections (5, 7).

Bacterial DNA was isolated using a DNeasy blood and tissue kit (Qiagen), according to the manufacturer's protocols. DNA libraries were prepared using a TruSeq Nano kit (Illumina) and sequenced on a MiSeq platform (2 × 300 bp). In parallel, genomic DNA was used to prepare barcoded DNA with Native barcoding kit 1D (product number EXP-NBD103; Oxford Nanopore Technologies), together with Ligation sequencing kit 1D (SQK-LSK108; Oxford Nanopore Technologies). DNA was sequenced using R9.4 chemistry (FLO-MIN106; Oxford Nanopore Technologies), and the raw signal was base called using Albacore 1.2.6. The sequences were assembled *de novo* into a single contig using the Unicycler hybrid assembler (8), with default settings.

The WG5 strain has a circular complete genome of 4,592,887 bp. Genome annotation was acquired from NCBI Prokaryotic Genome Annotation Pipeline (9), which revealed 4,657 genes, 4,536 coding sequences, 22 rRNAs (5S, 16S, and 23S), 87 tRNAs, and 22 noncoding RNAs. Multilocus sequence types were identified using MSTL (10). No horizontally identified antibiotic resistance genes were detected by using ResFinder (11). The PHASTER prophage finder (12) identified three prophage regions, with two incomplete prophage regions and one questionable prophage region.

Accession number(s). The genome sequences have been deposited in GenBank under the accession number [CP024090](https://doi.org/10.1128/genomeA.01403-17). The version described in this paper is the first version.

ACKNOWLEDGMENTS

This research was funded by the EU H2020 ERC-20104-STG LimitMDR (grant 638902), the Danish Council for Independent Research Sapere Aude Program DFF-4004-00213, and the European Union Seventh Framework Programme (FP7-KBBE-2013-7-singlestage, no.

Received 8 November 2017 **Accepted** 14 November 2017 **Published** 11 January 2018

Citation Imamovic L, Misiakou M-A, van der Helm E, Panagiotou G, Muniesa M, Sommer MOA. 2018. Complete genome sequence of *Escherichia coli* strain WG5. *Genome Announc* 6:e01403-17. <https://doi.org/10.1128/genomeA.01403-17>.

Copyright © 2018 Imamovic et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Morten Otto Alexander Sommer, msom@bio.dtu.dk.

613745, Promys). M.O.A.S. acknowledges additional funding from the Novo Nordisk Foundation Center for Biosustainability and the Lundbeck Foundation. M.-A.M. acknowledges Spanish Ministerio de Innovación y Ciencia (AGL2016-75536-P), the Agencia Estatal de Investigación (AEI), and the European Regional Fund (ERF).

REFERENCES

1. ISO. 2000. Water quality—Detection and enumeration of bacteriophages—Part 2: enumeration of somatic coliphages. ISO 10705-2:2000. International Organization for Standardization, Geneva, Switzerland.
2. Claassen P, Hedimbi M, Basson K. 2015. Use of affordable materials to improve water quality in peri-urban settlements in Windhoek, Namibia. *AiM* 5:190–197. <https://doi.org/10.4236/aim.2015.53018>.
3. Imamovic L, Muniesa M. 2011. Quantification and evaluation of infectivity of Shiga toxin-encoding bacteriophages in beef and salad. *Appl Environ Microbiol* 77:3536–3540. <https://doi.org/10.1128/AEM.02703-10>.
4. Imamovic L, Jofre J, Schmidt H, Serra-Moreno R, Muniesa M. 2009. Phage-mediated Shiga toxin 2 gene transfer in food and water. *Appl Environ Microbiol* 75:1764–1768. <https://doi.org/10.1128/AEM.02273-08>.
5. Havelaar AH, Hogeboom WM. 1983. Factors affecting the enumeration of coliphages in sewage and sewage-polluted waters. *Antonie Van Leeuwenhoek* 49:387–397.
6. Grabow WOK, Coubrough P. 1986. Practical direct plaque assay for coliphages in 100-ml samples of drinking water. *Appl Environ Microbiol* 52:430–433.
7. Muniesa M, Jofre J. 2004. Factors influencing the replication of somatic coliphages in the water environment. *Antonie Van Leeuwenhoek* 86: 65–76. <https://doi.org/10.1023/B:ANTO.0000024909.75523.be>.
8. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
9. Tatusova T, DiCuccio M, Badretdin A, Chetverin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
10. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <https://doi.org/10.1128/JCM.06094-11>.
11. Kleinheinz KA, Joensen KG, Larsen MV. 2014. Applying the ResFinder and VirulenceFinder web-services for easy identification of acquired antibiotic resistance and *E. coli* virulence genes in bacteriophage and phage nucleotide sequences. *Bacteriophage* 4:e27943. <https://doi.org/10.4161/bact.27943>.
12. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.