



Geographic and socioeconomic diversity of food and nutrient intakes: a comparison of four European countries

Mertens, Elly; Kuijsten, Anneleen; Dofková, Marcela; Mistura, Lorenza; D'Addezio, Laura; Turrini, Aida; Dubuisson, Carine; Favret, Sandra; Havard, Sabrina; Trolle, Ellen

Published in:
European Journal of Nutrition

Link to article, DOI:
[10.1007/s00394-018-1673-6](https://doi.org/10.1007/s00394-018-1673-6)

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Mertens, E., Kuijsten, A., Dofková, M., Mistura, L., D'Addezio, L., Turrini, A., ... Geleijnse, J. M. (2018). Geographic and socioeconomic diversity of food and nutrient intakes: a comparison of four European countries. *European Journal of Nutrition*. <https://doi.org/10.1007/s00394-018-1673-6>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Geographic and socioeconomic diversity of food and nutrient intakes: a comparison of four European countries

Elly Mertens¹ · Anneleen Kuijsten¹ · Marcela Dofková² · Lorenza Mistura³ · Laura D'Addezio³ · Aida Turrini³ · Carine Dubuisson⁴ · Sandra Favret⁴ · Sabrina Havard⁴ · Ellen Trolle⁵ · Pieter van't Veer¹ · Johanna M. Geleijnse¹

Received: 28 July 2017 / Accepted: 22 March 2018
© The Author(s) 2018

Abstract

Purpose Public health policies and actions increasingly acknowledge the climate burden of food consumption. The aim of this study is to describe dietary intakes across four European countries, as baseline for further research towards healthier and environmentally-friendlier diets for Europe.

Methods Individual-level dietary intake data in adults were obtained from nationally-representative surveys from Denmark and France using a 7-day diet record, Italy using a 3-day diet record, and Czech Republic using two replicates of a 24-h recall. Energy-standardised food and nutrient intakes were calculated for each subject from the mean of two randomly selected days.

Results There was clear geographical variability, with a between-country range for mean fruit intake from 118 to 199 g/day, for vegetables from 95 to 239 g/day, for fish from 12 to 45 g/day, for dairy from 129 to 302 g/day, for sweet beverages from 48 to 224 ml/day, and for alcohol from 8 to 15 g/day, with higher intakes in Italy for fruit, vegetables and fish, and in Denmark for dairy, sweet beverages and alcohol. In all countries, intakes were low for legumes (<20 g/day), and nuts and seeds (<5 g/day), but high for red and processed meat (>80 g/day). Within countries, food intakes also varied by socio-economic factors such as age, gender, and educational level, but less pronounced by anthropometric factors such as overweight status. For nutrients, intakes were low for dietary fibre (15.8–19.4 g/day) and vitamin D (2.4–3.0 µg/day) in all countries, for potassium (2288–2938 mg/day) and magnesium (268–285 mg/day) except in Denmark, for vitamin E in Denmark (6.7 mg/day), and for folate in Czech Republic (212 µg/day).

Conclusions There is considerable variation in food and nutrient intakes across Europe, not only between, but also within countries. Individual-level dietary data provide insight into the heterogeneity of dietary habits beyond per capita food supply data, and this is crucial to balancing healthy and environmentally-friendly diets for European citizens.

Keywords Diet · Foods · Nutrients · Dietary guidelines · Europe · SUSFANS

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00394-018-1673-6>) contains supplementary material, which is available to authorized users.

✉ Elly Mertens
elly.mertens@wur.nl

¹ Division of Human Nutrition, Wageningen University, PO Box 8129, 6700 EV Wageningen, The Netherlands

² Center for Health, Nutrition and Food, National Institute of Public Health, Brno, Czech Republic

³ Council for Agricultural Research and Economics, Research Centre for Food and Nutrition, Via Ardeatina 546, 00178 Rome, Italy

Introduction

Poor dietary habits are the second-leading risk factor for deaths and disability-adjusted life-years (DALYs) globally, accounting for 10.3 million deaths and 229.1 million

⁴ French Agency for Food, Environmental and Occupational Health and Safety (Anses)/Risk Assessment Department (DER), 14 rue Pierre et Marie Curie, 94701 Maisons-Alfort Cedex, France

⁵ Division of Risk Assessment and Nutrition, National Food Institute, Technical University of Denmark, Søborg, Denmark

DALYs in 2016 [1]. Low intakes of whole grains, fruit and vegetables, and nuts and seeds, and high intakes of alcohol and sodium ranked among the leading risk factors for early death and disability in European populations. However, as westernisation of diets progressed, diets high in red and processed meat, followed by diets high in sugar-sweetened beverages and low in milk are becoming a growing public health concern.

Dietary patterns are shaped by cultural, environmental, technological and economic factors, and they have become more similar over time owing to a general rise in living standards and globalisation of the food sector [2, 3]. Also in Europe there is a growing similarity of diets, in which traditional diets of Northern and Mediterranean countries are converging towards a more Western diet, viewed by the increased share of fruit and vegetables in Northern countries and the increased share of animal-based products in Mediterranean countries [4–6]. Increase in animal-based products and excessive caloric intake have been thought as a key factor in nutrition transition, which warrants the need for public health action to promote healthier food patterns consistent with traditional cultural preferences, hence the development of food-based dietary guidelines.

Food-based dietary guidelines are evidence-based integrated messages aimed at the general population for maintaining health and the prevention of non-communicable diseases [7, 8]. Promoting the intake of whole grains, fruit and vegetables, low-fat dairy and fish, and limiting the intake of red and processed meat, sugar-sweetened food products, alcohol and salt is covered by most national food-based dietary guidelines [9], although recommended quantities may differ. Monitoring food consumption patterns and assessing adherence to dietary guidelines in a nationally representative sample is especially regarded as a key instrument for evaluating the effectiveness of public health action towards a healthier diet.

In recent years, public health policies and actions have increasingly acknowledged the climate burden of food production and consumption, hence the need to address the food-climate connection, as outlined in the SUSFANS project (Metrics, Models and Foresight for European Sustainable Food And Nutrition Security) [10]. Production and technological changes in the food system will, however, not be sustainable without a change in food consumption patterns. The SUSFANS project, therefore, elaborates on the status-quo of diets and the design of optimised diets that are environmentally Sustainable, Healthy, Affordable, Reliable and Preferred (SHARP). This paper is a first step to study European food consumption patterns in terms of food groups and nutrients using national dietary survey data carried out at the individual level in four countries. Intakes of food groups and nutrients were compared with current food-based dietary guidelines and

nutrient reference values, overall and in relevant population subgroups.

Populations and methods

Data sources

Individual-level dietary intake data from national dietary surveys representative for different European regions, i.e. Denmark (Scandinavia) [11], Czech Republic (Central East Europe) [12], Italy (Mediterranean) [13] and France (Western Europe) [14], were collated for adult population aged ≥ 18 years within the SUSFANS project [10]. These four countries were chosen to capture the wide range of foods and agricultural commodities, including their extreme intakes, that are incorporated in the diverse European food consumption patterns.

Survey characteristics

Survey characteristics are shown in Table 1. National representativeness was ensured using random sampling based on civil registration systems in Denmark [11], national census data in Czech Republic [12] and France [14], and national census data with telephone books in Italy [13] that served as sampling frame, and followed by appropriate weighing for socio-demographic parameters, as applied in Denmark [11, 15] and France [14]. Surveys were organised throughout the whole year, covering the four seasons of the year, and have dietary data on week- and weekend-days.

Method of dietary assessment

In the four study countries, dietary intake was assessed over two to seven 24-h periods, either consecutively for 3–7 days using a diet record, as applied in Denmark, Italy and France [11, 13, 14], or non-consecutively spaced over a 3–5 months sampling period using two replicates of 24-h recall, as applied in Czech Republic [12]. In the present analyses, dietary intake from two random days has been reported. To this end, two non-consecutive days were sampled in Denmark, Italy and France, whereas all available days were used in Czech Republic.

Food and nutrient intakes

Intakes of food groups and nutrients were calculated for each subject from the mean of the selected two days, and were standardised for energy using the density method. Densities were calculated as the absolute value divided by total energy intake, and multiplied by 2000 kcal. Harmonised food groups, including similar foods, have been elaborated

Table 1 Dietary surveys in four European countries, i.e. Denmark, Czech Republic, Italy and France, including adult population only

	Denmark	Czech Republic	Italy	France
Survey characteristics, including adult population only				
Survey, year	The Danish National Survey on Diet and Physical Activity 2005–2008 National Food Institute, Technical University of Denmark (DTU)	Czech National Food Consumption Survey 2003–2004 (SISP04) National Institute of Public Health	Italian National Food Consumption Survey INRAN-SCAI 2005–2006 National institute for Research on Food and Nutrition	Individual and National Study on Food Consumption INCA-2 2006–2007 Agence Française de Sécurité Sanitaires des Aliments (AFSSA)
Population	18–75 years	18–90 years	18–98 years	18–79 years
Method of dietary assessment ^a	7-day diet record on consecutive days	24-h recall on two non-consecutive days	3-day diet record on consecutive days	7-day diet record on consecutive days
Baseline characteristics of the study sample, including adult population only, <i>n</i> (%)				
Sample size (response rate)	2025 (54%)	1869 (54%)	2831 (33%)	2624 (60%)
Age, 18–64 years	1739 (85.9%)	1666 (89.1%)	2313 (81.7%)	2276 (86.7%)
Gender, men	777 (44.7%)	793 (47.6%)	1068 (46.2%)	936 (41.1%)
Educational level, low	248 (14.2%)	345 (20.7%)	692 (31.7%)	1039 (45.8%)
Overweight status, BMI \geq 25	739 (43.2%)	864 (51.9%)	828 (35.8%)	871 (38.7%)

BMI Body Mass Index

^aIncluded in the present study were for Czech Republic both day, for Denmark and France two randomly selected days, and for Italy the first and the last day of the national dietary survey

using the ‘Exposure Hierarchy’ of the food classification and description system FoodEx2 developed and revised in 2015 by the European Food Safety Authority (EFSA) [16, 17]. A main challenge to encounter when grouping the foods was the level of food disaggregation; disaggregation of foods into ingredients was only considered as necessary for composite/prepared foods provided that the food itself was not included in FoodEx2, but its ingredients are. Nutrient intakes were calculated from dietary sources only, i.e. excluding dietary supplements, using country-specific food composition tables [18–24]. Intakes of added sugar, plant and animal protein were calculated based on food selection. Added sugar was defined as the total sugar intake minus sugars naturally occurring in fruits, vegetables and dairy. Plant protein was defined as protein derived from cereals, legumes, nuts and seeds, and others (including potatoes, vegetables, fruits, etc.). Animal protein was defined as protein derived from meat and meat products, fish and fish products, egg and egg products, milk and milk products (including cream, cheese and butter). None of the data excluded under- and over-reporting, however, misreporting was identified using Goldberg equation [25] and adopted by Black [26] (Online Resource 1).

Dietary quality

Foods

To evaluate European populations’ energy-standardised food group intakes, reference values were set for the food groups that are important for disease risk reduction based on an inventory of the current food-based dietary guidelines of European countries. Minimum values were set for foods that are beneficial for health, such as fruits and vegetables, and maximum values for foods that are unfavourable for health, such as red and processed meat (see Box 1). Reference values were derived using the 2015 Dutch food-based dietary guidelines [8] as reference point, complemented by the food-based dietary guidelines of the four countries [27–30] in which the less restrictive reference values were chosen.

Box 1 A set of food-based dietary guidelines for European countries, including their exposure definition and reference values, developed for the SUSFANS project

	Exposure definition	Reference values ^a
Foods to increase		
Fruit	All kind of fruits (including fresh, dried, tinned or canned fruit products, but excluding fruit juice)	≥ 200 g/day
Vegetables	All kind of vegetables (including fresh, dried, tinned or canned vegetable products, but excluding potatoes, vegetable juices and vegetables from soup, sauces and ready-to-eat products)	≥ 200 g/day
Legumes	Kidney beans, pinto beans, white beans, black beans, garbanzo beans (chickpeas), lima beans, split peas, lentils, and edamame (green soybeans)	≥ 135 g/week (≥ 19 g/day)
Nuts and seeds	Walnuts, almonds, hazel, cashew, pistachio, macadamia, Brazil, pecan, pine nuts, flax seeds, sesame seeds, sunflower seeds, pumpkin seeds, poppy seeds, and peanut	≥ 15 g/day
Dairy products	Food products produced from the milk of mammals, including milk, yoghurt, fresh uncured cheese, quark, custard, milk puddings, excluding cheese and butter	≥ 300 g/day
Fish	All kind of fish and fish products	≥ 150 g/week (≥ 21 g/day)
Foods to decrease		
Red and processed meat	Red meat: all mammalian muscle meat, including beef, veal, pork, lamb, mutton, horse and goat, excluding rabbit meat; Processed meat: meat transformed through salting, curing, fermentations, smoking or other processed to enhance flavour or improve preservation (e.g. meat products as sandwich filling, ready-to-eat minced meat, sausages, etc.)	≤ 500 g/week (≤ 71 g/day)
Cheese	All types of cheese formed by coagulation of milk protein casein	≤ 150 g/week (≤ 21 g/day)
Sugar-sweetened beverages	Cold beverages with added sugars (sucrose, fructose or glucose), for example fruit juices, fruit nectars, soft drinks, ice teas, vitamin-water or sports drinks with added sugars	≤ 500 ml/week (≤ 71 ml/day)
Alcohol (Ethanol)	Ethanol content calculated from all kind of alcoholic beverages	≤ 10 g/day
Foods to replace ^b		
Whole grains	Whole grains (bran, germ and endosperm in their natural proportion) from cereals, pasta, bread, breakfast cereals and other grain sources	Replace white grains by whole grains
White meat	Meat from all kind of poultry, including rabbit meat	Replace red and processed meat by white meat
Soft margarine and oils	Soft margarine: soft-solid fats made from vegetables oils; Oils: liquid fats at room temperature derived from plants or fish	Replace butter and hard margarines by soft margarine and oils

^aReference values were derived from current food-based dietary guidelines, using the 2015 Dutch food-based dietary guidelines [8] as reference point, complemented by the food-based dietary guidelines of the four countries [34–37] in which the less restrictive reference values was chosen (Quantitative guideline)

^bFoods to replace' represent food groups for which insufficient convincing evidence was available to set a fixed cut-off point, however replacement of those food products by a healthier alternative is recommended (Qualitative guideline)

Nutrients

To evaluate European populations' energy-standardised nutrient intakes, nutrient density of the diet was quantified using Nutrient Rich Diet (NRD) score [31, 32], i.e. overall summary estimate of nutrient intakes based on the principles of the Nutrient Rich Food Index [33, 34]. The NRD algorithm was calculated as:

$$\text{NRD } X \cdot Y = \sum_{i=1}^{i=X} \frac{Q_{\text{nutrient } i}}{\text{DRV}_i} \times 100 - \sum_{j=1}^{j=Y} \frac{Q_{\text{nutrient } j}}{\text{MRV}_j} \times 100$$

where X is the number of qualifying nutrients, Y is the number of disqualifying nutrients, Q nutrient i or j is the average daily intake of nutrient i or j , DRV is the dietary reference value of qualifying nutrient i and MRV j is the maximum recommended value of the nutrient to limit j . DRVs are defined using reference values from EFSA [35], i.e. average requirement (AR), and adequate intake (AI) if AR cannot be set, and MRVs using reference values of World Health Organisation [36, 37] and Food and Agriculture Organisation [38].

In the present analyses, NRD9.3 and NRD15.3 were used. The NRD9.3, including nine nutrients for which

intake should be promoted (protein, dietary fibre, calcium, iron, potassium, magnesium, and vitamin A, C and E) and three nutrients for which intake should be limited (saturated fat (SFA), added sugar, and sodium), standardised for 2000 kcal/day diet and capped nutrient intake at 100% of DRV was primarily chosen, based on its validation among US populations [33, 34]. To capture more nutrients that are potentially relevant for European populations, we also used its extended version, i.e. NRD15.3 that additionally included mono-unsaturated fatty acids, zinc, vitamin D and B-vitamins (B1, B2, B12, folate), but excluded magnesium. A sub-score on the intake of qualifying nutrients is represented in NRD9 and NRD15, and that of disqualifying nutrients in NRDX.3, while the total score, i.e. NRD9.3 and NRD15.3, is a combination of both.

Estimating the dietary quality of European populations' diets

Percentages of the population that adhere to food-based dietary guidelines and percentages of the population with inadequate nutrient intakes were estimated using the AR cut-point method [39], without correction for within subject variability. This percentage would be interpreted as proxy figures for adherence and inadequacy, because of different survey's methodologies. When the DRV of the nutrient under study was defined as an AI (dietary fibre, potassium, magnesium, vitamin D, E and B12), this percentage of populations with intake below AI was only applicable for comparison between countries and population subgroups. Dietary intakes were characterised in the overall country-specific population of adults aged ≥ 18 years and in relevant population subgroups by age, gender, educational level, and overweight status. Subgroups by age included younger and middle-aged adults (18–64 years) and elderly (≥ 65 years). Younger and middle-aged adult populations were additionally stratified by gender, educational level using three categories, i.e. primary or lower secondary degree ('low'), higher secondary degree ('intermediate') and university or post-university degree ('high'), and overweight status using two categories, i.e. BMI < 25 and ≥ 25 kg/m².

As the information available consisted only of summarised data (i.e. mean and standard deviation of the energy-standardised dietary intake under study and sample size), analysis of variance test was performed to check whether there were differences in mean intake of food groups and nutrients between countries and within countries by population subgroups of age, gender, educational level and overweight status. Bonferroni post hoc test was used for multiple comparisons. A two sided *p* value below 0.0001 was considered as statistically significant. Statistical analyses were performed with SAS version 9.3 (SAS Institute Inc.).

Results

Baseline characteristics

Age and gender distribution were comparable between countries, with 80–90% of the population aged 18–64 years and 40–48% being men. Distribution of educational level varied markedly between countries; a low proportion of low-educated subjects in Denmark (15%) and a high proportion in France (46%); but proportion of the high-educated subjects was the lowest in Czech Republic (8%) and varied between 23–33% for Denmark, Italy and France. Approximately half of the Czech population (52%) was overweight, BMI ≥ 25 kg/m², whereas overweight in Denmark (44%), France (39%) and Italy (36%) was less prevalent.

Foods

Table 2 shows the energy-standardised intakes of food groups and general adherence to food-based dietary guidelines in four European adult populations, aged ≥ 18 years. Stratified intakes by age, gender, educational level and overweight status are shown in Table 3.

Foods to increase

Mean fruit and vegetable intake varied significantly between countries with lower intakes for Czech Republic (118 and 95 g/day, respectively) and higher intakes for Italy (199 and 239 g/day, respectively), and varied in the same direction between men and women within all four countries showing higher intakes for women. Higher fruit intake was also observed in all four countries for the elderly and for subjects with a higher educational level, but no differences by overweight status. Vegetable intake tended to be higher among elderly in Denmark and France, among higher educated subjects in Denmark and Czech Republic, and among overweight subjects in Italy and France. Mean intakes of legumes (6.5–16.7 g/day), and nuts and seeds (0.5–2.6 g/day) were generally low in all countries. Mean intake of dairy was higher in Denmark (302 g/day), while fish was higher in Italy (44.6 g/day) and France (34.4 g/day).

Foods to decrease

Mean intake of red and processed meat was generally high in all countries (84–94 g/day). Within-countries, red and processed meat intake was lower for the elderly and women in all four countries, and except in Italy for the higher educated subjects, and in Czech Republic and France for the non-overweight. Alcohol intake varied between countries

Table 2 Energy-standardised food group intakes and the adherence to their corresponding food-based dietary guidelines in four European populations, aged ≥ 18 years

	Denmark (n = 2025)			Czech Republic (n = 1869)			Italy (n = 2831)			France (n = 2624)		
	Mean	Median	%adh	Mean	Median	%adh	Mean	Median	%adh	Mean	Median	%adh
Foods to increase												
Fruit, g/day	174*	133	35%	118*	83	20%	199*	163	40%	140*	95	26%
Vegetables, g/day	147*	112	21%	95*	74	10%	239*	206	53%	187*	157	37%
Legumes, g/day	6.5	1.6	10%	7.5	0.0	12%	11.0	0.0	19%	16.5*	0.0	18%
Nuts and seeds, g/day	2.2	0.0	5%	2.6	0.0	7%	0.5*	0.0	1%	1.7	0.0	3%
Dairy products, g/day	302*	248	41%	134	94	12%	129	116	8%	199*	152	24%
Fish, g/day	18.0	5.5	28%	11.7	0.0	17%	44.6*	6.5	42%	34.3*	4.3	43%
Foods to decrease												
Red and processed meat, g/day	94	85	39%	88	82	42%	84	77	51%	93	82	43%
Cheese, g/day	29.3	24.3	44%	20.9*	13.2	63%	53*	47.2	28%	30.1	24.0	46%
Sweet beverages ^a , ml/day	224*	127	40%	108	0.0	63%	47.5*	0.0	76%	121	6.0	56%
Alcohol (ethanol), g/day	14.6*	7.3	56%	10.3	4.4	66%	8.2	0.1	67%	9.3	0.1	67%
Foods to replace												
Cereals, total, g/day	26.1*	16.9	–	48.2	32.5	–	46.6	38.3	–	38.8*	16.05	–
Cereals, whole grains, g/day	0.4	0.0	–	0.1	0.0	–	0.8	0.0	–	1.8	0.0	–
Pasta, total, g/day	5.2*	0.0	–	39.9*	13.6	–	52*	48.4	–	10.3*	0.0	–
Pasta, whole grains, g/day	–	–	–	0.0*	0.0	–	0.3*	0.0	–	9.8*	0.0	–
Bread, total, g/day	149*	140	–	122*	118	–	109*	103	–	98*	92	–
Bread, whole grains, g/day	52*	44.3	–	7.9*	0.0	–	41.4*	0.0	–	16.3*	0.0	–
Breakfast cereals, total, g/day	11.8*	0.6	–	2.9	0.0	–	1.5	0.0	–	5.3*	0.0	–
Breakfast cereals, whole grains, g/day	9.3*	0.0	–	1.9*	0.0	–	0.5*	0.0	–	3.4*	0.0	–
Red meat, g/day	66*	57.1	–	34.0*	28.4	–	58	53	–	58	45.6	–
Processed meat, g/day	27.3	19.4	–	54*	44.5	–	25.5	19.4	–	34.7*	22.6	–
White meat, g/day	21.3	1.6	–	22.5	0.0	–	23.5	0.0	–	31.5*	0.0	–
Butter and hard margarines, g/day	24.8*	22.7	–	17.6*	15.5	–	2.8*	0.0	–	16.3*	13.7	–
Soft margarine and oils, g/day	1.9*	0.0	–	15.0*	13.1	–	34.8*	34.0	–	11.2*	7.4	–

Intake of food groups are standardised to a 2000 kcal/day diet

%adherence represents a proxy for the percentage of the population that adhere to food-based dietary guidelines

^aSweet beverages instead of sugar-sweetened beverages due to a lack of detailed data on beverages

*Bonferroni $p < 0.0001$ test comparison for intake that was significantly different from all other three countries under study

Table 3 Energy-standardised food group intakes and the adherence to their corresponding food-based dietary guidelines in subgroups by age, gender, educational level, and overweight status: main findings

Cut-offs	Subgroups by age						Subgroups by gender ^a						p value						
	Younger and middle-aged adults			Elderly, ≥ 65 years			Men			Women				p value					
	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh							
Denmark	(n = 1739)	(n = 286)	(n = 777)	(n = 962)															
Fruit, g/day	≥ 200	171	126	(32.2; 251)	34%	197	159	(81; 281)	40%	0.011	120	74	(0.5; 172)	21%	222	187	(74; 324)	47%	<0.0001
Vegetables, g/day	≥ 200	151	114	(64; 189)	22%	119	98	(54; 167)	16%	<0.0001	117	95	(54; 146)	13%	185	141	(84; 231)	31%	<0.0001
Legumes, g/day	≥ 19	6.6	1.8	(0.0; 7.1)	10%	5.3	0.9	(0.0; 4.6)	10%	<0.0001	5.9	1.3	(0.0; 5.6)	8%	7.3	2.2	(0.0; 8.6)	11%	<0.0001
Red and processed meat, g/day	≤ 71	95	87	(52; 128)	38%	83	73	(41.5; 108)	48%	0.001	109	100	(66; 143)	29%	82	75	(43.3; 114)	47%	<0.0001
Alcohol, g/day	≤ 10	13.8	6.4	(0.0; 21.5)	58%	20.5	15.0	(1.7; 29.8)	40%	<0.0001	16.6	10.0	(0.0; 25.6)	50%	10.9	0.0	(0.0; 17.0)	66%	<0.0001
Czech Republic	(n = 1666)	(n = 203)	(n = 793)	(n = 873)															
Fruit, g/day	≥ 200	115	79	(10.0; 167)	19%	143	118	(38.7; 218)	28%	0.006	66	39	(0.7; 93)	6%	160	128	(51; 224)	31%	<0.0001
Vegetables, g/day	≥ 200	95	75	(39.3; 128)	10%	94	70	(39.4; 122)	8%	0.874	78	61	(35.0; 106)	5%	111	87	(46.0; 151)	14%	<0.0001
Legumes, g/day	≥ 19	7.6	0.0	(0.0; 2.2)	11%	6.7	0.0	(0.0; 4.2)	13%	0.591	6.1	0.0	(0.0; 1.7)	10%	9.0	0.0	(0.0; 2.6)	12%	0.012
Red and processed meat, g/day	≤ 71	89	81	(44.8; 125)	42%	83	79	(45.3; 118)	42%	0.253	108	103	(69; 142)	27%	71	64	(28.4; 103)	55%	<0.0001
Alcohol, g/day	≤ 10	10.7	5.1	(0.0; 17.0)	65%	7.4	0.0	(0.0; 9.4)	77%	0.002	15.8	12.5	(1.2; 23.5)	47%	6.1	0.0	(0.0; 8.6)	81%	<0.0001
Italy	(n = 2313)	(n = 518)	(n = 1068)	(n = 1245)															
Fruit, g/day	≥ 200	185	153	(67; 257)	37%	257	222	(125; 333)	54%	<0.0001	153	125	(50.4; 220)	28%	214	185	(88; 292)	45%	<0.0001
Vegetables, g/day	≥ 200	238	205	(134; 299)	52%	241	215	(149; 307)	55%	0.680	222	190	(126; 282)	47%	252	156	(145; 317)	56%	<0.0001
Legumes, g/day	≥ 19	10.7	0.0	(0.0; 2.9)	19%	12.4	0.0	(0.0; 0.0)	19%	0.194	10.1	0.0	(0.0; 3.9)	19%	11.3	27.1	(0.0; 2.3)	19%	0.265

Table 3 (continued)

Cut-offs	Subgroups by age						Subgroups by gender ^a																
	Younger and middle-aged adults			Elderly, ≥ 65 years			Men			Women													
	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh							
Red and pro-processed meat, g/day	85	77	(37.6; 120)	65%	75	68	(31.6; 111)	62%	0.015	88	81	(43.6; 122)	65%	82	74	(32.7; 119)	64%	<0.0001					
Alcohol, g/day	≤10	7.8	(0.0; 12.7)	70%	10.0	2.6	(0.0; 16.5)	60%	0.0002	11.3	6.8	(0.0; 18.9)	57%	4.8	8.4	(0.0; 7.0)	80%	<0.0001					
France	(n=2276)	129	77	(0.0; 198)	23%	209	174	(77; 309)	42%	<0.0001	103	65	(0.0; 154)	17%	148	103	(0.0; 219)	28%	<0.0001				
Fruit, g/day	≥200	182	152	(80; 248)	36%	219	196	(110; 293)	46%	<0.0001	152	128	(65; 204)	26%	202	173	(95; 272)	45%	<0.0001				
Vegetables, g/day	≥19	15.9	0.0	(0.0; 0.8)	17%	20.9	0.0	(0.0; 5.3)	20%	0.040	17.7	0.0	(0.0; 1.8)	19%	14.6	0.0	(0.0; 0.4)	16%	0.068				
Legumes, g/day	≤71	94	84	(40.7; 134)	43%	90	79	(37.8; 133)	45%	0.316	101	92	(49.8; 143)	38%	88	77	(33.9; 127)	47%	<0.0001				
Red and pro-processed meat, g/day	≤10	9.0	0.0	(0.0; 13.8)	69%	11.2	5.2	(0.0; 18.2)	56%	0.008	13.5	6.6	(0.0; 21.1)	57%	5.8	0.0	(0.0; 7.3)	81%	<0.0001				
Alcohol, g/day	≤10	9.0	0.0	(0.0; 13.8)	69%	11.2	5.2	(0.0; 18.2)	56%	0.008	13.5	6.6	(0.0; 21.1)	57%	5.8	0.0	(0.0; 7.3)	81%	<0.0001				
Subgroups by educational level ^a																							
	Low						Intermediate						High										
	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh							
Denmark	(n=248)	152	94	(0.0; 234)	29%	159	115	(30.4; 233)	32%	214	167	(64; 305)	42%	<0.0001	167	124	(33.1; 246)	34%	174	129	(23.5; 255)	33%	0.382
Fruit, g/day	≥200	126	96	(56; 152)	16%	150	118	(63; 185)	21%	184	137	(84; 238)	32%	<0.0001	154	118	(66; 191)	23%	146	108	(63; 182)	21%	0.072
Vegetables, g/day	≥19	6.1	0.4	(0.0; 6.7)	10%	6.5	1.6	(0.0; 6.8)	10%	7.7	2.8	(0.0; 7.8)	11%	<0.0001	6.4	1.9	(0.0; 6.9)	9%	6.9	1.5	(0.0; 7.4)	11%	0.055
Legumes, g/day	≤71	102	90	(58; 143)	39%	99	92	(58; 131)	33%	82	75	(44.5; 111)	46%	<0.0001	94	86	(52; 126)	38%	99	90	(54; 134)	37%	0.072
Red and pro-processed meat, g/day	≤71	102	90	(58; 143)	39%	99	92	(58; 131)	33%	82	75	(44.5; 111)	46%	<0.0001	94	86	(52; 126)	38%	99	90	(54; 134)	37%	0.072
Subgroups by overweight status ^a																							
	BMI < 25 kg/m ²						BMI ≥ 25 kg/m ²						p value										
	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh	Mean	Median	(P25; P75)	%adh							
Denmark	(n=972)	152	94	(0.0; 234)	29%	159	115	(30.4; 233)	32%	214	167	(64; 305)	42%	<0.0001	167	124	(33.1; 246)	34%	174	129	(23.5; 255)	33%	0.382
Fruit, g/day	≥200	126	96	(56; 152)	16%	150	118	(63; 185)	21%	184	137	(84; 238)	32%	<0.0001	154	118	(66; 191)	23%	146	108	(63; 182)	21%	0.072
Vegetables, g/day	≥19	6.1	0.4	(0.0; 6.7)	10%	6.5	1.6	(0.0; 6.8)	10%	7.7	2.8	(0.0; 7.8)	11%	<0.0001	6.4	1.9	(0.0; 6.9)	9%	6.9	1.5	(0.0; 7.4)	11%	0.055
Legumes, g/day	≤71	102	90	(58; 143)	39%	99	92	(58; 131)	33%	82	75	(44.5; 111)	46%	<0.0001	94	86	(52; 126)	38%	99	90	(54; 134)	37%	0.072
Red and pro-processed meat, g/day	≤71	102	90	(58; 143)	39%	99	92	(58; 131)	33%	82	75	(44.5; 111)	46%	<0.0001	94	86	(52; 126)	38%	99	90	(54; 134)	37%	0.072

Table 3 (continued)

	Subgroups by educational level ^a						Subgroup by overweight status ^a								
	Low			Intermediate			High			BMI < 25 kg/m ²			BMI ≥ 25 kg/m ²		
	Mean	Median	(P25; P75) %adh	Mean	Median	(P25; P75) %adh	Mean	Median	(P25; P75) %adh	Mean	Median	(P25; P75) %adh	Mean	Median	(P25; P75) %adh
Alcohol, g/day	13.2	6.3	(0.0; 21.4) 58%	13.7	6.0	(0.0; 20.6) 59%	15.0	8.8	(0.0; 24.5) 52%	13.2	6.2	(0.0; 20.5) 58%	14.5	6.7	(0.0; 23.4) 57%
Czech Republic	(n=345)			(n=1194)			(n=127)			(n=802)			(n=864)		
Fruit, g/day	89	61	(1.3; 141) 11%	122	82	(13.4; 173) 21%	121	96	(40.1; 179) 20%	112	79	(19.1; 165) 19%	118	79	(5.9; 168) 19%
Vegetables, g/day	90	71	(40.0; 123) 8%	94	74	(37.0; 126) 10%	120	85	(59; 160) 15%	96	77	(40.0; 126) 10%	95	73	(37.8; 128) 9%
Legumes, g/day	8.9	0.0	(0.0; 3.0) 12%	7.3	0.0	(0.0; 2.0) 11%	7.3	0.0	(0.0; 2.7) 11%	7.3	0.0	(0.0; 2.3) 11%	7.9	0.0	(0.0; 2.1) 11%
Red and processed meat, g/day	96	86	(47.4; 134) 42%	88	82	(44.3; 124) 41%	81	72	(43.4; 117) 48%	83	73	(40.0; 121) 48%	94	88	(50.2; 130) 37%
Alcohol, g/day	11.7	5.0	(0.0; 19.0) 61%	10.5	4.8	(0.0; 16.3) 66%	10.1	7.7	(0.0; 16.8) 61%	10.4	4.5	(0.0; 16.9) 65%	11.0	5.5	(0.0; 16.9) 64%
Italy	(n=692)			(n=985)			(n=507)			(n=1484)			(n=828)		
Fruit, g/day	182	155	(69; 260) 38%	183	149	(65; 250) 36%	206	169	(83; 282) 41%	185	155	(68; 249) 37%	187	150	(68; 272) 37%
Vegetables, g/day	242	206	(137; 296) 53%	238	205	(136; 300) 52%	232	202	(129; 287) 51%	229	200	(130; 288) 50%	254	213	(144; 323) 55%
Legumes, g/day	11.7	0.0	(0.0; 4.1) 22%	10.7	0.0	(0.0; 3.3) 19%	10.1	0.0	(0.0; 4.5) 17%	10.5	0.0	(0.0; 2.3) 19%	11.1	0.0	(0.0; 4.2) 19%
Red and processed meat, g/day	88	81	(41.0; 122) 65%	85	77	(37.5; 119) 65%	83	77	(35.9; 121) 65%	84	77	(36.8; 118) 65%	86	78	(39.2; 124) 64%
Alcohol, g/day	8.8	0.0	(0.0; 15.3) 66%	7.1	0.1	(0.0; 11.9) 72%	7.4	0.2	(0.0; 11.1) 74%	6.8	0.0	(0.0; 11.2) 73%	9.6	4.0	(0.0; 15.9) 62%
France	(n=1039)			(n=495)			(n=737)			(n=1379)			(n=871)		
Fruit, g/day	125	76	(0.0; 200) 24%	128	84	(0.0; 195) 21%	137	95	(13.9; 196) 23%	126	82	(0.0; 191) 22%	134	89	(0.0; 204) 24%
Vegetables, g/day	181	152	(77; 248) 36%	179	144	(74; 245) 33%	183	156	(87; 249) 37%	175	146	(75; 242) 33%	188	158	(85; 254) 39%

Table 3 (continued)

	Subgroups by educational level ^a						Subgroup by overweight status ^a										
	Low			Intermediate			High			BMI < 25 kg/m ²			BMI ≥ 25 kg/m ²				
	Mean	Median (P25; P75)	%adh	Mean	Median (P25; P75)	%adh	Mean	Median (P25; P75)	%adh	Mean	Median (P25; P75)	%adh	Mean	Median (P25; P75)	%adh		
Legumes, ≥19 g/day	19.5	0.0 (0.0; 1.3)	21%	13.2	0.0 (0.0; 0.4)	15%	12.5	0.0 (0.0; 0.5)	15%	0.0003	16.3	0.0 (0.0; 1.1)	19%	15.5	0.0 (0.0; 0.5)	16%	0.645
Red and processed meat, ≤71 g/day	102	91 (48.7; 144)	39%	90	79 (33.5; 129)	44%	84	74 (33.9; 123)	47%	<0.0001	89	78 (35.7; 127)	44%	101	91 (48.7; 145)	40%	0.0001
Alcohol, ≤10 g/day	8.3	0.0 (0.0; 11.8)	73%	9.4	0.2 (0.0; 15.1)	66%	9.6	0.2 (0.0; 15.5)	67%	0.135	8.0	0.0 (0.0; 12.1)	73%	10.6	0.1 (0.0; 16.9)	64%	<0.0001

Intake of food groups are standardised to a 2000 kcal/day diet

%adherence represents a proxy for the percentage of the population that adhere to food-based dietary guidelines

BMI Body Mass Index

^aYounger and middle-aged adults, aged 18–64 years, were stratified by gender, educational level and overweight status

^b*p* value for the overall comparisons between population subgroups

with lower intakes in Italy (8.2 g/day) and higher intakes for Denmark (14.6 g/day), and varied within countries in the same direction by gender and overweight status with lower intakes for women and the non-overweight. Alcohol intake also tended to be lower for the young and middle-aged adults, except in Czech Republic where intake is lower for the elderly. For the higher-educated subjects, alcohol intake tended to be lower in Czech Republic and Italy, but higher in Denmark and France.

Foods to replace

Mean intakes of whole grains from cereals, pasta and bread were low in all countries, illustrated by the fraction of whole grains on total grains of ≤15% with one exception for wholegrain pasta in France. Although mean intake of total breakfast cereals per day was very low, the whole grain variants were primarily eaten. Intake of white meat was much lower than red and processed meat, in particular red and processed meat contributed to 70–80% of total meat intake comprising mainly of red meat in Denmark, Italy and France, and of processed meat in Czech Republic. Intakes of butter and hard margarines were only slightly higher than intakes of soft margarines and vegetable oils, except for Denmark where butter and hard margarines were predominantly chosen as fat source, and for Italy where vegetable oils were dominating.

Nutrients

Table 4 shows the energy-standardised nutrient intakes, their corresponding proxy prevalence figures for inadequate intakes, and the NRD scores in four European adult populations, aged ≥18 years. Low intakes were observed for dietary fibre (15.8–19.4 g/day) and vitamin D (2.4–3.0 µg/day) in all countries, and for potassium (2288–2939 mg/day), and magnesium (268–285 mg/day), except in Denmark. Intake of vitamin E was lower in Denmark (6.7 mg/day), and folate in Czech Republic (212 µg/day). Mean intakes were high for protein (67.1–83.5 g/day), and iron (9.1–12.4 mg/day) in all countries analysed. Remaining nutrients, including calcium, zinc, vitamin A, C, B1, B2, and B12, showed varying intake levels between countries. Of the three nutrients to limit, a large penalty was obtained from saturated fatty acids (11.1–15.1 E%) in all countries, and from estimated sodium intake (2797–4244 mg/day) except in Italy. Based on the NRD scores, it is apparent that the nutrient density of the diet was highest in Italy (NRD9.3 of 537, and NRD15.3 of 1051), followed by Denmark (NRD9.3 of 416, and NRD15.3 of 896) and France, and the lowest in Czech Republic (NRD9.3 of 327 and NRD15.3 of 787). Within countries, nutrient density of the diet tended to be higher

Table 4 Energy-standardised nutrient intakes, prevalence of inadequate intake, and Nutrient Rich Diet scores in four European populations, aged ≥ 18 years

DRV	Denmark (n = 2025)			Czech Republic (n = 1869)			Italy (n = 2831)			France (n = 2624)						
	Mean	Median	% < DRV	Mean	Median	% < DRV	Mean	Median	% < DRV	Mean	Median	% < DRV				
Unstandardised energy intake, kcal/day	2264*	2155	(168.1; 2738)	–	2523*	2396	(1790; 3106)	–	2119*	2057	(1666; 2491)	–	1980*	1912	(1509; 2390)	–
Qualifying nutrients																
Protein, g/day	68.7	67.6	(59.7; 77.1)	16%	67.1	66.1	(59.1; 73.8)	12%	79.0*	77.8	(70.5; 86.1)	1%	83.5*	81.4	(70.9; 93.4)	2.4%
Protein, E%	–	13.9	(12.4; 15.2)	–	13.4	13.2	(11.8; 14.8)	–	15.6*	15.6	(14.1; 17.2)	–	16.7*	16.3	(14.2; 18.7)	–
Animal protein, g/day	44.8*	43.2	(35.6; 52.8)	–	38.8*	37.5	(30.1; 45.8)	–	48.6*	47.1	(38.9; 56.8)	–	–	–	–	–
Plant protein, g/day	20.3*	20.2	(16.9; 23.6)	–	23.9*	23.8	(20.1; 27.3)	–	30.3*	30.3	(26.5; 34)	–	–	–	–	–
Dietary fibre, g/day ^a	19.4*	18.6	(14.5; 23.2)	81%	15.8*	15.1	(12.7; 18.3)	96%	18.1*	17.0	(14.0; 21.0)	88%	16.6*	15.7	(12.3; 19.5)	91%
MUFA, g/day ^a	25.7*	25.5	(21.0; 30.0)	–	32.0*	31.8	(27.8; 36.4)	–	39.0*	38.7	(33.5; 44.1)	–	29.7*	28.9	(24.0; 34.2)	–
MUFA, E%	11.7*	11.6	(9.5; 13.6)	31%	14.4*	14.3	(12.5; 16.4)	8%	17.6*	17.4	(15.1; 19.9)	25%	13.4*	13.0	(10.8; 15.4)	23%
Calcium, mg/day	983*	928	(705; 1189)	30%	660*	593	(424; 805)	69%	742*	708	(539; 897)	57%	899*	842	(649; 1066)	38%
Iron, mg/day	M: 6; F: 7	9.1*	(7.7; 10.2)	8%	10.6*	10.1	(8.5; 12.1)	4%	11.1*	10.5	(9.0; 12.3)	2%	12.4*	11.2	(9.4; 13.8)	2%
Potassium, mg/day ^b	3500	3143*	(2514; 3658)	69%	2288*	2199	(1895; 2573)	96%	2938	2834	(2420; 3326)	81%	2879	2763	(2326; 3287)	82%
Magnesium, mg/day ^b	M: 350; F: 300	322*	(270; 365)	54%	285	274	(241; 315)	75%	268*	254	(219 299)	80%	282	263	(230 ; 309)	77%
Zinc, mg/day	M: 7.5; F: 6.2	9.5*	(8.1; 10.8)	10%	7.0*	6.7	(5.6; 8.0)	52%	11.0*	10.5	(9.1; 12.4)	3%	10.2*	9.6	(8.1; 11.8)	9%
Vitamin A, µg RE/day	M: 570; F: 490	1032*	(557; 1242)	23%	692*	450	(315; 631)	62%	854*	635	(467; 924)	34%	1200*	822	(552; 1279)	23%
Vitamin C, mg/day	M: 90; F: 80	102*	(57; 131)	50%	78*	63	(37; 103)	65%	126*	103	(66; 159)	38%	91*	76	(46; 119)	56%
Vitamin E, mg/day ^b	M: 13; F: 11	6.7*	(5.1; 7.7)	95%	11.7*	11.1	(8.4; 14.4)	56%	12.7*	11.8	(9.7; 14.1)	53%	10.6*	9.4	(6.9; 13.2)	66%
Vitamin D, µg/day ^b	15	3.0	(1.3; 2.7)	97%	2.9	2.1	(1.4; 3.2)	99%	2.4	1.5	(1.0; 2.4)	99%	2.6	1.7	(1.0; 3.0)	99%
Vitamin B1, mg/day	0.6	1.1	(0.9; 1.3)	3%	1.1	1.0	(0.9; 1.2)	2%	1.10	0.9	(0.8; 1.1)	53%	1.20	1.1	(0.9; 1.3)	0%
Vitamin B2, mg/day	M: 1.1; F: 0.9	1.47*	(1.13; 1.70)	20%	1.08*	0.99	(0.84; 1.20)	65%	1.40*	1.3	(1.1; 1.6)	16%	1.80*	1.7	(1.4; 2.1)	8%
Vitamin B12, µg/day ^b	4	4.7	(3.1; 5.6)	45%	4.4	3.4	(2.5; 4.8)	64%	6.1	4.1	(3.1; 5.8)	48%	5.6	4.0	(2.9; 5.8)	50%

Table 4 (continued)

DRV	Denmark (n=2025)				Czech Republic (n=1869)				Italy (n=2831)				France (n=2624)			
	Mean	Median	(P25; P75)	%<DRV	Mean	Median	(P25; P75)	%<DRV	Mean	Median	(P25; P75)	%<DRV	Mean	Median	(P25; P75)	%<DRV
Folate, µg DFE/d	293	268	(214; 334)	41%	212*	182	(146; 242)	76%	350*	305	(254; 380)	23%	278	253	(203; 322)	49%
Disqualifying nutrients	MRV			%> MRV				%> MRV				%> MRV				%> MRV
SFA, g/day	30.4	30.2	(25.0; 35.4)	–	30.6	30.4	(25.5; 35.1)	–	24.6*	24.2	(20.3; 28.3)	–	33.5*	33.4	(27.7; 39.1)	–
SFA, E%/day ^d	13.8	13.7	(11.3; 16.1)	86%	13.8	13.7	(11.5; 15.8)	80%	11.1*	10.9	(9.1; 12.7)	62%	15.1*	15.0	(12.5; 17.6)	91%
Added sugar, g/day	43.2*	36.4	(21.3; 57.2)	–	36.6	31.3	(18.8; 50.6)	–	38.6	35.2	(21.1; 52.5)	–	–	–	–	–
Added sugar, E% ^d	8.8*	7.4	(4.3; 11.6)	32%	7.3	6.3	(3.8; 10.1)	21%	7.7	7.0	(4.2; 10.5)	24%	–	–	–	–
Sodium, mg/day ^d	3012*	2919	(2484; 3439)	80%	4244*	4153	(3576; 4800)	98%	1703*	1648	(1245; 2076)	13%	2797*	2668	(2228; 3223)	85%
Nutrient Rich Diet Scores																
Sub-score NRD9	765	775	(710; 829)	–	715*	721	(643; 794)	–	781*	793	(730; 841)	–	759	767	(701; 826)	–
Sub-score NRD15	1245	1259	(1192; 1310)	–	1175*	1182	(1097; 263)	–	1295*	1310	(1246; 1356)	–	1250	1262	(1191; 1324)	–
Sub-score NRD9.3	349*	346	(300; 392)	–	388*	387	(347; 427)	–	244*	242	(215; 271)	–	–	–	–	–
Total score NRD9.3	416*	427	(334; 507)	–	327*	328	(256; 400)	–	537*	547	(482; 600)	–	–	–	–	–
Total score NRD15.3	896*	916	(823; 992)	–	787*	791	(704; 875)	–	1051*	1062	997; 1115	–	–	–	–	–

DRV dietary reference value, AR average requirement, AI adequate intake, RE retinol equivalents, DFE dietary folate equivalents, E% energy percentage, MUFA mono-unsaturated fatty acids, SFA saturated fatty acids, NRD Nutrient Rich Diet scores, including their sub-scores

Intakes of nutrients are standardised to a 2000 kcal/day diet

^a%<AR represents a proxy for the percentage of the population that have an inadequate intake, i.e. intake lower than the dietary reference value

^bNutrients where AR cannot be set, hence AI is defined

^cCannot be computed

^dPercentages shown for SFA, added sugar and sodium reflect the proportion of the population that have an excessive intake, i.e. intake higher than the reference value (Maximum Recommend Value)

*Bonferroni $p < 0.0001$ test comparison for intake that was significantly different from all other three countries under study

for women in all four countries and for the higher-educated subject, except in Italy (Table 5).

Discussion

In this study, we found that dietary intakes varied markedly across the four European countries, irrespective of energy intake. Within countries, food intakes also varied markedly by socio-economic factors such as age, gender, and educational level, but less pronounced by anthropometric factors such as overweight status. However, the set of food-based dietary guideline was not met by a large part of the population and/or population subgroup by age, gender, educational level or overweight status.

When describing food group intakes, mean daily intakes of fruit and vegetables, sweet beverages, and alcohol varied most between countries, showing higher intakes of fruit and vegetables, and lower intakes of sweet beverages and alcohol in Italy. In addition, we observed in Italy and France a similar vegetable intake among the different levels of education, whereas in Denmark and Czech Republic higher intake of vegetables was observed among higher-educated subjects; which is in line with previous studies conducted in European populations [40–42]. This region-dependent tendency might be attributed to the long-standing cultural tradition of using vegetables in the Mediterranean diet, as consumed in Italy and France, and is often easily recognisable by all layers of the population. However, a comparison of population subgroups within-countries is often closely related to dietary preferences, beliefs and practices of that particular consumer group. Higher intake of fish, nuts and seeds along with lower intake of red and processed meat are, for example, generally seen among women and higher-educated subjects, which might be driven by their health considerations and awareness of climate change [43].

When describing nutrient intakes summarised by the NRD9.3 and 15.3, the higher scores were observed for Italy, which is mainly attributed to their lower penalty score, i.e. NRDX.3, for the disqualifying nutrients of SFA and sodium. Because of the interrelation between food groups and nutrients intake, our results on variation in nutrient intakes can be partly reflected by our results on variation in food group intake. Low penalty score in Italy is likely to be in correspondence with its lower intakes for important sources of SFA intake such as butter and hard margarines, red and processed meat, and dairy products; however, with the estimates of sodium intake, caution must be applied, as they are very likely to be underestimated due to difficulties in quantifying sodium content in recipes and discretionary salt intake [44]. Moreover, when focussing on qualifying nutrients, higher sub-scores NRD9 and NRD15 were also observed for Italy, but intake

for calcium, potassium and magnesium was lower when compared with Denmark; related to intake of dairy products and whole-grain products. It could, thus, be argued whether these summary estimates could be used solely to describe nutrient intakes, as they do not point out specific inadequate nutrient intakes.

In the context of the SUSFANS project, we prefer to describe dietary intakes in terms of foods rather than nutrients, since foods are the constituents of a dietary pattern and the common denominator for linking dietary intakes with health, environment, affordability, consumer's preferences, etc. Diet-associated environmental impact, in particular, has been attracting a lot of interest, as current food production and consumption patterns have been recognised as a major human-induced driver of climate change [45]. Some European countries have, therefore, developed guidelines for diets that are both healthy and environmentally-friendly [46–49]. Such recommendations mostly emphasise the reduction of greenhouse gas emissions through propagating a shift towards plant-based foods. However, given European dietary intakes, there is still much progress to be made in this respect, simply showed by a percentage of around 35% for the intake of plant protein as opposed to total protein for the countries we studied. Moreover, predominant food groups contributing to animal and plant protein intake have been associated with regional and cultural traditions around dietary habits. Meat intake is regarded as the most important contributor to animal protein in European diets, but with differences related to the amount and types of meat consumed, as also denoted by previous studies [50, 51]. With regard to plant protein, cereals and cereal products have been identified as the main contributor to plant protein in European diets [52], while joint contributions from vegetables, legumes and fruit varied between countries, as observed in the present study.

The present study provides further support for the application of individual-level dietary data to address the food-climate connection. Often diet-associated environmental impact was quantified using food availability data related to food production, but not to food consumption as such. Using individual-level reported dietary data might, therefore, be regarded as a useful tool in the connection between health and environment with foods as their common denominator. Cross-country comparison of individual-level dietary data is, however, challenged by the dietary surveys conducted with different survey characteristics and data collection methods that may influence the comparability of the results. First, sampling procedures used in the surveys reported in this study varied in terms of recruitment methods, household and individual representativeness, number of subjects per household and weighting factors used; however, they all aimed at including a nationally representative sample of at least all age-sex categories. It still remains a possibility that those

Table 5 Nutrient density of the diet, using Nutrient Rich Diet scores 9.3 and 15.3, in four European populations in subgroups by age, gender, educational level and overweight status

	Subgroups by age				Subgroups by gender ^a				<i>p</i> value			
	Younger and middle-aged adults		Elderly, ≥ 65 years		Men		Women					
	Mean	Median	(P25; P75)	Mean	Median	(P25; P75)	Mean	Median		(P25; P75)		
Denmark	(<i>n</i> = 1739)		(<i>n</i> = 286)		(<i>n</i> = 777)		(<i>n</i> = 965)					
Sub-score NRD9	764	774	(708; 829)	772	787	(721; 833)	733	796	(679; 786)	808	(758; 853)	<0.0001
Sub-score NRD15	1243	1256	(1191; 1308)	1256	1275	(1198; 1325)	1215	1271	(1162; 1280)	1284	(1226; 1328)	<0.0001
Sub-score NRDx.3	351	348	(301; 395)	333	336	(291; 382)	355	346	(309; 400)	339	(297; 388)	0.011
Total score NRD9.3	413	424	(327; 505)	439	424	(328; 505)	376	450	(295; 456)	465	(388; 537)	<0.0001
Total score NRD15.3	892	913	(817; 988)	923	940	(847; 1010)	860	925	(780; 944)	944	(859; 1021)	<0.0001
Czech Republic	(<i>n</i> = 1666)		(<i>n</i> = 203)		(<i>n</i> = 793)		(<i>n</i> = 873)					
Sub-score NRD9	714	720	(641; 793)	729	728	(666; 807)	659	763	(597; 719)	777	(713; 821)	<0.0001
Sub-score NRD15	1174	1182	(1092; 1261)	1185	1181	(1114; 1269)	1119	1223	(1039; 1197)	1235	(1157; 1297)	<0.0001
Sub-score NRDx.3	387	385	(345; 427)	396	395	(360; 430)	375	398	(333; 417)	397	(358; 436)	<0.0001
Total score NRD9.3	327	327	(253; 400)	333	342	(270; 401)	284	366	(216; 349)	373	(298; 440)	<0.0001
Total score NRD15.3	787	790	(703; 876)	789	792	(711; 873)	744	826	(665; 821)	836	(751; 910)	<0.0001
Italy	(<i>n</i> = 2313)		(<i>n</i> = 518)		(<i>n</i> = 1068)		(<i>n</i> = 1245)					
Sub-score NRD9	777	790	(725; 837)	796	805	(759; 852)	747	803	(692; 806)	814	(764; 856)	<0.0001
Sub-score NRD15	1293	1307	(1240; 1350)	1305	1321	(1272; 1360)	1264	1317	(1210; 1330)	1329	(1278; 1367)	<0.0001
Sub-score NRDx.3	245	243	(215; 271)	242	240	(213; 269)	242	247	(212; 271)	245	(219; 272)	0.004
Total score NRD9.3	533	541	(476; 598)	554	563	(509; 609)	505	556	(443; 572)	565	(509; 614)	<0.0001
Total score NRD15.3	1048	1059	(991; 1115)	1064	1075	(1021; 1122)	1022	1070	(959; 1091)	1079	(1024; 1127)	<0.0001
France	(<i>n</i> = 2276)		(<i>n</i> = 348)		(<i>n</i> = 936)		(<i>n</i> = 1340)					
Sub-score NRD9	754	762	(696; 821)	785	787	(743; 841)	717	788	(668; 775)	799	(743; 846)	<0.0001
Sub-score NRD15	1244	1256	(1182; 1319)	1278	1289	(1222; 1346)	1208	1278	(1147; 1284)	1289	(1228; 1346)	<0.0001

Table 5 (continued)

	Subgroups by educational level ^a						Subgroup by overweight status ^a						<i>p</i> value			
	Low			Intermediate			High			BMI < 25 kg/m ²				BMI ≥ 25 kg/m ²		
	Mean	Median (P25; P75)		Mean	Median (P25; P75)		Mean	Median (P25; P75)		Mean	Median (P25; P75)			Mean	Median (P25; P75)	
Denmark	(<i>n</i> = 248)			(<i>n</i> = 943)			(<i>n</i> = 548)			(<i>n</i> = 972)			(<i>n</i> = 739)			
Sub-score NRD9	746	754 (690; 814)	760	767	791	803	791	791	769	779	779	759	766	766	766	
Sub-score NRD15	1221	1236 (1165; 1293)	1242	1254	1271	1282	1271	1271	1250	1261	1261	1237	1249	1249	1249	
Sub-score NRD9.3	356	356 (305; 404)	356	350	334	334	334	334	351	349	349	351	347	347	347	
Total score NRD9.3	390	404 (292; 498)	405	414	456	459	456	456	408	418	418	408	418	418	418	
Total score NRD15.3	865	893 (767; 978)	887	905	937	942	937	937	887	908	908	887	907	907	907	
Czech Republic	(<i>n</i> = 345)			(<i>n</i> = 1194)			(<i>n</i> = 127)		(<i>n</i> = 802)			(<i>n</i> = 864)				
Sub-score NRD9	695	684 (624; 780)	716	722	740	744	740	740	719	725	725	709	713	713	713	
Sub-score NRD15	1153	1149 (1060; 1252)	1175	1181	1217	1238	1217	1217	1175	1186	1186	1172	1178	1178	1178	
Sub-score NRD9.3	378	378 (339; 421)	390	387	384	381	384	384	389	390	390	385	382	382	382	
Total score NRD9.3	317	307 (237; 387)	327	327	356	360	356	356	330	329	329	324	323	323	323	
Total score NRD15.3	775	775 (681; 862)	785	789	833	847	833	833	786	791	791	787	789	789	789	
Italy	(<i>n</i> = 692)			(<i>n</i> = 985)			(<i>n</i> = 507)		(<i>n</i> = 1484)			(<i>n</i> = 828)				
Sub-score NRD9	774	788 (718; 835)	776	789	788	801	788	788	779	792	792	775	788	788	788	
Sub-score NRD15	1291	1309 (1234; 1355)	1292	1304	1300	1316	1300	1300	1294	1308	1308	1291	1307	1307	1307	
Sub-score NRD9.3	240	240 (211; 267)	246	243	249	246	249	249	248	245	245	240	237	237	237	
Total score NRD9.3	534	545 (478; 603)	530	536	539	550	539	539	531	539	539	535	545	545	545	
Total score NRD15.3	1051	1065 (992; 1118)	1046	1056	1051	1064	1051	1051	1046	1058	1058	1051	1064	1064	1064	
France	(<i>n</i> = 1039)			(<i>n</i> = 495)			(<i>n</i> = 737)		(<i>n</i> = 1379)			(<i>n</i> = 871)				
Sub-score NRD9	749	760 (681; 822)	756	763	761	764	761	761	753	760	760	758	766	766	766	

Table 5 (continued)

Sub-score NRD15	Subgroups by educational level ^a				Subgroup by overweight status ^a				<i>p</i> value			
	Low		Intermediate		High		BMI < 25 kg/m ²			BMI ≥ 25 kg/m ²		
	Mean	Median (P25; P75)	Mean	Median (P25; P75)	Mean	Median (P25; P75)	Mean	Median (P25; P75)		Mean	Median (P25; P75)	
	1237	1252 (1166; 1319)	1247	1250 (1194; 1314)	1254	162 (1190; 1326)	0.002	1242	1256 (1177; 1316)	1249	1258 (1191; 1329)	0.110

BMI Body Mass Index, *NRD* Nutrient Rich Diet scores, including their sub-scores

For France, sub-score NRD9.3 and 15.5 cannot be computed due to a lack of data on sugars

^aYounger and middle-aged adults, aged 18–64 years, were stratified by gender, educational level and overweight status

^b*p* value for the overall comparisons between population subgroups

who have agreed to participate form a group with a greater interest in health, hence more optimistic results.

Second, methods of dietary assessment used in the surveys reported were conducted differently, with regard to the methods used and in the manner in which the assessment was carried out. Replicates of 24-h recall as applied in Czech Republic showed a higher mean energy intake compared to diet records as applied in Denmark, Italy and France. This might be explained by factors related to the methods themselves, such as reliance on memory and portion size estimations [53–55], and/or characteristics of the populations. Standardising intake data to a 2000 kcal/day diet had, therefore, the largest impact on results of Czech Republic; lowering its mean dietary intakes under the assumption that energy intake is positively correlated with food group and nutrient intake. Standardisation for energy is one of the more practical ways of reducing part of the extraneous variation in dietary estimates [56], and enables to study the relative contribution of food groups and nutrients intake to the total diet, regardless of energy intake. In the European Food CONsumption VALidation project, it has been suggested to adjust for BMI instead when analysing and interpreting dietary data of nutritional monitoring surveys to reduce mean bias at population level [57]. Given that stratified analyses by overweight status showed no relevant differences in dietary intakes within a country, it is questionable whether BMI-adjusted values should be the main exposure of interest in the present study describing the heterogeneity of European diets.

Another important factor in estimating dietary intakes consistently is the number of days included in the dietary assessment to enable comparison between countries across Europe. In this study, dietary data were, therefore, standardised for the number of days, but have not been corrected for time-interval between the two selected record/recall days, hence not corrected for within-subject day-to-day variability. Correcting for within-subject day-to-day variability would have resulted in comparable means for dietary intakes compared to unadjusted data, though with a shrinkage of intake distributions which in turn would have decreased the percentage of the population above and below a cut-off point [58]. However, relying on consecutive days, including days spaced over a week time-interval, is likely to underestimate the within-subject day-to-day variation [59] because of the interdependence of days that captures some of the day-to-day variation in the between-subject variation [60, 61]. Thus, this day-interdependence would have resulted in a shrinkage of the observed intake distribution that is too much toward the group mean, hence an under-estimation of true percentage of the population above and below a cut-off when statistically correcting intake distributions. In addition, the use of country-specific food composition databases might affect the number of subjects whose intake was below the DRV. In

particular, when using different food composition databases, potential systematic errors in estimating nutrient intake would be different between countries, and in all probability alternate with magnitude and direction. With increasing globalisation, however, the foods and mixed dishes available in different countries are not all grown/produced/prepared in the same manner and, therefore, using a country-specific composition database is likely to reflect nutrient intake more accurately.

Exclusion of under-reporters would have increased the prevalence of adherence to the food-based dietary guidelines and decreased the prevalence of inadequate nutrient intakes, and inclusion of supplementation use would have decreased the prevalence of nutrient inadequacy even further. The present study did estimate the percentage of under- and over-reporters (Online Resource 1), but did not estimate intakes excluding them, because some of the misreporters may truly be consuming a low- or a high-energy diet. Over the past decades, dietary supplementation use has increased in Europe with a clear north–south gradient [62], showing a high number of users in Denmark (Online Resource 1). Hence, it is likely that in countries with higher level of supplementation use, dietary supplementation might have contributed to improved total nutrient intakes, with its impact dependent on the supplementation formulation, the frequency of use, and the level of micronutrient intakes of those taking supplements. However, our interest is on nutrient intakes from foods only to find nutritional gaps that are most in need to improve the healthiness of dietary intake.

In conclusion, there is considerable variation in food and nutrient intakes across European countries. The present study indicated that the intake of food groups showed larger deviations from food-based dietary guidelines for the overall population and population subgroups of the countries we studied. In addition, results suggested inadequate nutrient intakes from foods for dietary fibre and vitamin D in all countries, and for potassium, magnesium, vitamin E and folate in specific regions. Individual-level dietary data in different European population and population subgroups are, therefore, needed for balancing diets for European citizen.

Moreover, individual-level dietary data from national surveys serve as a practical tool for describing the healthiness of diet in terms of foods and nutrients, but dietary data harmonisation remains challenging. Using a common food classification system is a first step in the alignment of surveys and necessary to enable cross-country comparisons for food group intakes. However, further steps, such as standardisation for energy, number of days, etc., are needed for harmonisation of dietary data. Besides the healthiness of dietary intake, these dietary surveys might also be important in shaping optimised diets where other factors, such as environmental impact, affordability and consumer preferences are incorporated. We aim, therefore, to support

further engagement of key stakeholders from the food supply chain and policy-makers in the next stages for the design of SHARP diets.

Author contributions JMG and PvtV initiated the topic of the paper. MD, LM, LD, AT, CD, SF, and ET were responsible for the data collection and data analysis. EM, AK and were responsible for data interpretation. EM drafted the manuscript, which was reviewed by all authors for intellectual content. All authors read and approved the final submission of the paper.

Funding Financial support for this original contribution was obtained from funding from the European Union's H2020 Programme under Grant Agreement number 633692 (SUSFANS: Metrics, models and foresight for European sustainable food and nutrition security) and from the Top Consortia for Knowledge and Innovation of the Dutch Ministry of Economic Affairs.

Compliance with ethical standards

Conflict of interest The authors have no conflicts of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. GBD 2016 Risk Factors Collaborators (2017) Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 390:1345–1422. [https://doi.org/10.1016/S0140-6736\(17\)32366-8](https://doi.org/10.1016/S0140-6736(17)32366-8)
2. Traill WB, Mazzocchi M, Shankar B, Hallam D (2014) Importance of government policies and other influences in transforming global diets. *Nutr Rev* 72:591–604
3. Global Panel on Agriculture and Food Systems for Nutrition (2016) Food systems and diets: Facing the challenges of the 21st century. Global Panel on Agriculture and Food Systems for Nutrition, London
4. Schmidhuber J, Traill WB (2006) The changing structure of diets in the European Union in relation to healthy eating guidelines. *Public Health Nutr* 9:584–595
5. Balanza R, García-Lorda P, Pérez-Rodrigo C, Aranceta J, Bonet MB, Salas-Salvadó J (2007) Trends in food availability determined by the Food and Agriculture Organization's food balance sheets in Mediterranean Europe in comparison with other European areas. *Public Health Nutr* 10:168–176
6. Gerbens-Leenes P, Nonhebel S, Krol M (2010) Food consumption patterns and economic growth. Increasing affluence and the use of natural resources. *Appetite* 55:597–608
7. Mozaffarian D, Ludwig DS (2010) Dietary guidelines in the 21st century—a time for food. *JAMA* 304:681–682
8. Kromhout D, Spaaij C, de Goede J, Weggemans R (2016) The 2015 Dutch food-based dietary guidelines. *Eur J C Nutr* 70(8):869–878
9. World Health Organisation (WHO) (2003) Food based dietary guidelines in the WHO European region. WHO, Copenhagen:

10. Rutten M, Achterbosch TJ, de Boer IJ, Cuaresma JC, Geleijnse JM, Havlik P, Heckelet T, Ingram J, Leip A, Marette S (2016) Metrics, models and foresight for European sustainable food and nutrition security: the vision of the SUSFANS project. *Agric Syst*. <https://doi.org/10.1016/j.agry.2016.10.014>
11. Pedersen A, Fagt S, Groth MV, Christensen T, Biloft-Jensen A, Matthiessen J, Andersen NL, Kørup K, Hartkopp H, Ygil K, Hinsch HJ, Saxholt E, Trolle E (2009) Danskernes kostvaner 2003–2008. In: DTU Fødevareinstituttet
12. Ruprich JDM, Rehurkova I, Slamenikova E, Resova D (2006) Individual food consumption—the national study SISPO4. CHFCH National Institute of Public Health, Prague
13. Leclercq CAD, Piccinelli R, Sette S, Le Donne C, Turrini A (2009) The Italian national food consumption survey INRAN-SCAI 2005–06: main results in terms of food consumption. *Publ Health Nutr* 12(12):2504–2532
14. Agence Française de Sécurité Sanitaire des Aliments (AFSSA) (2009) Report of the 2006/2007 Individual and National Study on Food Consumption 2 (INCA 2). In: Synthèse de l'étude individuelle nationale des consommations alimentaires 2 (INCA 2), 2006–2007, pp 1–44
15. Matthiessen J, Stockmarr A, Biloft-Jensen A, Fagt S, Zhang H, Groth MV (2014) Trends in overweight and obesity in Danish children and adolescents: 2000–2008-exploring changes according to parental education. *Scand J Public Health* 42:385–392. <https://doi.org/10.1177/1403494813520356>
16. European Food Safety Authority (2015) The food classification and description system FoodEx2 (revision 2). *EFSA Suppl Publ* 804:90
17. EFSA (European Food Safety Authority) (2011) Use of the EFSA comprehensive European food consumption database in exposure assessment. *EFSA J* 9:2097
18. Møller ASE, Christensen AT, Hartkopp H (2005) Fødevaredata-banken version 6.0. Fødevareinformatik, Afdeling for Ernæring, Denmark
19. Saxholt E, Christensen AT, Møller A, Hartkopp HB, Hess Ygil H, Hels OH (2008) Fødevaredata-banken, version 7. In: Fødevareinformatik, Afdeling for Ernæring, Fødevareinstituttet, Danmarks Tekniske Universitet
20. Czech Centre for Food Composition Database (2016) Czech food composition database version 6.16. Institute of Agricultural Economics and Information, Prague
21. Food Research Institute (2016) Slovak food composition data bank. Department of Risk Assessment Food Composition Data Bank and Consumer's Survey VUP Food Research Institute, Bratislava
22. Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione (INRAN) (2016) Banca Dati di Composizione degli Alimenti. Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione, Roma
23. Feinberg M (1995) Répertoire général des aliments (General Inventory of Foods). In: FJ-CLC (ed) Institut national de la recherche agronomique. Technique & Documentation—Lavoisier, Paris
24. Ireland J, dCL, Oseredczuk M et al (2008) French food composition table, version 2008. In: French Food Safety Agency (AFSSA)
25. Goldberg G, Black A, Jebb S, Cole T, Murgatroyd P, Coward W, Prentice A (1991) Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr* 45:569–581
26. Black AE (2000) Critical evaluation of energy intake using the Goldberg cut-off for energy intake: basal metabolic rate. A practical guide to its calculation, use and limitations. *Int J Obes Relat Metab Disord* 24:1119
27. Ministry of Food Agriculture and Fisheries (2013) The official dietary guidelines (Danish: De officielle kostråd). Ministry of Food, Agriculture and Fisheries Glostrup, Denmark
28. Czech Society for Nutrition (2012) Nutrition recommendations for Czech Republic (Czech: Výživová doporučení pro obyvatelstvo České republiky). Czech Society for Nutrition, Prague
29. Italian National Research Institute on Food and Nutrition (INRAN; CRA-NUT) (2003) Guidelines for healthy Italian food habits, 2003 (Italian: Linee guida per una sana alimentazione italiana. Revisione 2003). Italian National Research Institute on Food and Nutrition (INRAN; CRA-NUT), Rome
30. Programme National Nutrition Santé (PNNS) (2016) La Santé vient en mangeant Le guide alimentaire pour tous. ANSES Agence Nationale de Sécurité Sanitaire de l'alimentation, de l'environnement et du travail, Maisons-Alfort Cedex
31. Van Kernebeek HRJ, Oosting SJ, Feskens EJM, Gerber PJ, De Boer IJM (2014) The effect of nutritional quality on comparing environmental impacts of human diets. *J Clean Prod* 73:88–99. <https://doi.org/10.1016/j.jclepro.2013.11.028>
32. Roos E, Karlsson H, Witthoft C, Sundberg C (2015) Evaluating the sustainability of diets—combining environmental and nutritional aspects. *Environ Sci Policy* 47:157–166. <https://doi.org/10.1016/j.envsci.2014.12.001>
33. Drewnowski A (2009) Defining nutrient density: development and validation of the nutrient rich foods index. *J Am Coll Nutr* 28:421 s–426 s
34. Fulgoni VL 3rd, Keast DR, Drewnowski A (2009) Development and validation of the nutrient-rich foods index: a tool to measure nutritional quality of foods. *J Nutr* 139:1549–1554. <https://doi.org/10.3945/jn.108.101360>
35. EFSA (European Food Safety Authority) (2010) Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2010. Scientific opinion on principles for deriving and applying dietary reference values. *EFSA J* 8:1458
36. World Health Organisation (WHO) (2012) Guideline: sodium intake for adults and children. WHO, Geneva
37. World Health Organisation (WHO) (2015) Guideline: sugars intake for adults and children. WHO, Geneva
38. Food and Agriculture Organisation (FAO) (2010) Fats and fatty acids in human nutrition. Report of an expert consultation. *FAO Food Nutr Pap* 91:1–166
39. Institute of Medicine (IOM) (2000) Dietary reference intakes: applications in dietary assessment. National Academy, Washington DC
40. De Irala-Estevez J, Groth M, Johansson L, Oltersdorf U (2000) A systematic review of socio-economic differences in food habits in Europe: consumption of fruit and vegetables. *Eur J Clin Nutr* 54:706
41. Prättälä R, Hakala S, Roskam A-JR, Roos E, Helmert U, Klumbiene J, Van Oyen H, Regidor E, Kunst AE (2009) Association between educational level and vegetable use in nine European countries. *Public Health Nutr* 12:2174–2182
42. Roos E, Talala K, Laaksonen M, Helakorpi S, Rahkonen O, Uutela A, Prättälä R (2008) Trends of socioeconomic differences in daily vegetable consumption, 1979–2002. *Eur J Clin Nutr* 62:823–833
43. de Boer J, Schosler H, Aiking H (2014) “Meatless days” or “less but better”? Exploring strategies to adapt Western meat consumption to health and sustainability challenges. *Appetite* 76:120–128. <https://doi.org/10.1016/j.appet.2014.02.002>
44. McLean RM (2014) Measuring population sodium intake: a review of methods. *Nutrients* 6:4651–4662
45. Tukker AHG, Guinée J, Heijungs R, de Koning A, van Oers L et al (2006) Environmental Impact of Products (EIPRO) Analysis of the life cycle environmental impacts related to the final consumption of the EU25. In: European Commission Technical Report EUR 22284 EN. IPTS/ESTO, European Commission Joint Research Centre Brussels

46. German Nutrition Society (2013) Ten guidelines for wholesome eating and drinking from the German Nutrition Society (German: Vollwertig essen und trinken nach den 10 Regeln der DGE). Deutsche Gesellschaft für Ernährung e.V., Bonn
47. The Swedish National Food Agency (Livsmedelsverket) (2017) Find your way to eat greener, not too much and to be active! (Hitta ditt sätt att äta grönare, lagom mycket och röra på dig!). Livsmedelsverket, Uppsala
48. Health Council of the Netherlands (2011) Guidelines for a healthy diet: the ecological perspective. Health Council of the Netherlands, The Hague
49. Macdiarmid J, Kyle J, Horgan G, Loe J, Fyfe C, Johnstone A, McNeill G (2011) Livewell: a balance of healthy and sustainable food choices. In: WWF-UK
50. Linseisen J, Kesse E, Slimani N, Bueno-De-Mesquita H, Ocké M, Skeie G, Kumle M, Iraeta MD, Gómez PM, Janzon L (2002) Meat consumption in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts: results from 24-hour dietary recalls. *Public Health Nutr* 5:1243–1258
51. Kushi LH, Lenart EB, Willett WC (1995) Health implications of Mediterranean diets in light of contemporary knowledge. 2. Meat, wine, fats, and oils. *Am J Clin Nutr* 61:1416S–1427S
52. Halkjaer J, Olsen A, Bjerregaard L, Deharveng G, Tjønneland A, Welch A, Crowe F, Wirfält E, Hellstrom V, Niravong M (2009) Intake of total, animal and plant proteins, and their food sources in 10 countries in the European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr* 63:S16–S36
53. Bingham S, Gill C, Welch A, Day K, Cassidy A, Khaw K, Sneyd M, Key T, Roe L, Day N (1994) Comparison of dietary assessment methods in nutritional epidemiology: weighed records v. 24 h recalls, food-frequency questionnaires and estimated-diet records. *Br J Nutr* 72:619–643
54. Holmes B, Dick K, Nelson M (2008) A comparison of four dietary assessment methods in materially deprived households in England. *Public Health Nutr* 11:444–456
55. De Keyzer W, Huybrechts I, De Vriendt V, Vandevijvere S, Slimani N, Van Oyen H, De Henauw S (2011) Repeated 24-hour recalls versus dietary records for estimating nutrient intakes in a national food consumption survey. *Food Nutr Res* 55:7307
56. Willett WC, Howe GR, Kushi LH (1997) Adjustment for total energy intake in epidemiologic studies. *Am J Clin Nutr* 65:1220S–1228S
57. Crispim SP, Geelen A, De Vries JH, Freisling H, Souverein OW, Hulshof PJ, Ocke MC, Boshuizen H, Andersen LF, Ruprich J (2012) Bias in protein and potassium intake collected with 24-h recalls (EPIC-Soft) is rather comparable across European populations. *Eur J Nutr* 51:997–1010
58. Dodd KW, Guenther PM, Freedman LS, Subar AF, Kipnis V, Midthune D, Tooze JA, Krebs-Smith SM (2006) Statistical methods for estimating usual intake of nutrients and foods: a review of the theory. *J Am Diet Assoc* 106:1640–1650
59. Larkin FA, Metzner HL, Guire KE (1991) Comparison of 3 consecutive-day and 3 random-day records of dietary-intake. *J Am Diet Assoc* 91:1538–1542
60. Tarasuk V, Beaton GH (1991) The nature and individuality of within-subject variation in energy-intake. *Am J Clin Nutr* 54:464–470
61. Ellozy M (1983) Dietary variability and its impact on nutritional epidemiology. *J Chron Dis* 36:237–249
62. Skeie G, Braaten T, Hjartaker A, Lentjes M, Amiano P, Jakszyn P, Pala V, Palanca A, Niekerk E, Verhagen H (2009) Use of dietary supplements in the European Prospective Investigation into Cancer and Nutrition calibration study. *Eur J Clin Nutr* 63:S226–S238