



## Hydrodynamic functionality of the lorica in choanoflagellates

**Walther, Jens Honore; Asadzadeh, Seyed Saeed; Nielsen, Lasse Tor; Andersen, Anders; Dölger, Julia; Kjørboe, Thomas; Larsen, Poul Scheel**

*Publication date:*  
2018

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*

Walther, J. H., Asadzadeh, S. S., Nielsen, L. T., Andersen, A., Dölger, J., Kjørboe, T., & Larsen, P. S. (2018). *Hydrodynamic functionality of the lorica in choanoflagellates*. Abstract from 71st Annual Meeting of the APS Division of Fluid Dynamics , Atlanta, United States.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

DFD18-000639

# Hydrodynamic functionality of the lorica in choanoflagellates

Jens Honore Walther<sup>1</sup>, Sayed Saeed Asadzadeh<sup>1</sup>, Lasse Tor Nielsen<sup>1</sup>, Anders Andersen<sup>1</sup>, Julia Dolger<sup>1</sup>, Thomas Kjørboe<sup>1</sup>, Poul Scheel Larsen<sup>1</sup>

<sup>1</sup>Technical University of Denmark


## Body:

Choanoflagellates are unicellular microswimmers that are ubiquitous in aquatic habitats. They have a single flagellum that creates a flow toward the collar, the filtration apparatus composed of closely spaced filter strands. Loricated choanoflagellates have evolved a basket-like “skeleton” around the cell, the lorica, the function of which remains unknown. Here, we use Computational Fluid Dynamics (CFD) to explore the possible hydrodynamic function of the lorica by studying the choanoflagellate *Diaphaoneca grandis*, with and without its lorica. We study the flow rate, the flow recirculation, and the resulting clearance rate for the capture of motile and non-motile prey by the freely swimming choanoflagellate. We find no support for several previous hypotheses regarding the effects of the lorica. Rather, our simulations suggest that the main function of the lorica is to enhance the capture efficiency, but this happens at the cost of lower encounter rate with motile prey.

## Funding Acknowledgment:

We gratefully acknowledge funding from the Villum Foundation for SSA and JHW through research grant (9278), for AA, JD and TK through the Centre for Ocean Life, and from The Danish Council For Independent Research for LTN through research grant (7014-00033B).

## Special Instructions:



Close