



Mechanisms of action involved in chemically induced effects on male reproductive health

Schwartz, Camilla Victoria Lindgren; Christiansen, Sofie; Vinggaard, Anne Marie; Svingen, Terje

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Schwartz, C. V. L., Christiansen, S., Vinggaard, A. M., & Svingen, T. (2017). *Mechanisms of action involved in chemically induced effects on male reproductive health*. Poster session presented at 3rd ReproYoung Conference, Båstad, Sweden.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Mechanisms of action involved in chemically induced effects on male reproductive health

Schwartz C.L.^{1,*}; Christiansen S.¹; Vinggaard A.M.¹; and Svingen T.¹.

¹ Division of Diet, Disease Prevention and Toxicology, Technical University of Denmark, Søborg, Denmark.

*cavi@food.dtu.dk

Background


Fetal exposure to anti-androgenic compounds can adversely affect development of the reproductive system and in turn, reproductive function later in life. A shorter anogenital distance (AGD) in newborn animals and humans is a noninvasive biomarker of compromised male reproductive development and health. Although this morphometric effect is thought to be primarily caused by disrupted androgen signaling, the molecular mechanisms underpinning the effect are not well understood. This hinders the development of predictive, non-animal test methods for chemicals.

Aim

To characterize the molecular mechanisms affected when fetal perineal development is disrupted by xenobiotics.


1. Finasteride is an AR antagonist in vitro

Finasteride is a known 5 α -reductase inhibitor. Using an androgen receptor (AR) gene reporter assay, we also find Finasteride to be a strong AR antagonist in vitro.





2. In vivo study design

Finasteride (10 mg/kg bw/day) was used to induce shorter AGD in male offspring. Pregnant Sprague Dawley rats were exposed orally during GD7-21. Fetal AGD was measured on GD21 using a stereomicroscope. Blood and amniotic fluids and reproductive tissues were collected at GD17 and GD21 for further analyses.



3. Finasteride exposure shortens male AGD

Exposed male fetuses had significantly shorter AGD (~37%) compared to controls at GD21. Results are in agreement with previous findings for Finasteride, and confirms its utility for studying the underlying molecular mechanisms.



Results are shown as millimetre \pm SEM, n=5-6 litters/group. Statistical significance indicated by ***p < 0.0001. Statistical analyses was adjusted using litter as an independent, random and nested factor. Body weight was used as a covariate. ANOVA, followed by Dunnett's post hoc test in SAS. Female control AGDmm=1.81 mm

4. Molecular profiling in perineal development

Serial sagittal section of GD17 male fetus. Eosin staining (left) for identification of tissues of interest, including perineum. The mesenchymal cell marker SALL1 is expressed in the perineum (right; red). Together with antibodies such as MyoG this will be used to investigate cell type specific expression of the AR and its associated signalling pathways within the perineum, both in normal and disrupted development.

