



Controlling oxidation in skin care products with novel seaweed antioxidants

Hermund, Ditte Baun; Sivasubramaniam, Niruja; Yee Heung, Shuk; Neerup, R.; Klinder, Louise Mellisa ; Holdt, Susan Løvstad; Thomsen, Birgitte Raagaard; Jacobsen, Charlotte

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Hermund, D. B. (Author), Sivasubramaniam, N. (Author), Yee Heung, S. (Author), Neerup, R. (Author), Klinder, L. M. (Author), Holdt, S. L. (Author), ... Jacobsen, C. (Author). (2018). Controlling oxidation in skin care products with novel seaweed antioxidants. Sound/Visual production (digital)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Controlling oxidation in skin care products with novel seaweed antioxidants

Ditte B. Hermund, Niruja Sivasubramaniam, Shuk Yee Heung, Randi Neerup, Birgitte R. Thomsen & Charlotte Jacobsen

AOCS, 2018



Bioactive compounds

Fucus vesiculosus



VS

Saccharina latissima



Bioactive compounds	<i>Fucus</i>	<i>Saccharina</i>
Polysaccharides	62-66%	38-61%
Protein	1.4-17%	3-21%
α -Tocopherol	38-73 mg/kg dw	0.1 mg/kg dw
Pigments (fucoxanthin)	340 mg/kg dw	-
Phenolic compounds	0.4-12.2%	0.2-5.3%

Source: Holdt & Kraan, 2011


Antioxidant activity?

Studies (S): Extraction of antioxidant from brown algae

Tasks


- Extraction of antioxidants from S1) *Fucus vesiculosus* (Danish) and S2) *Saccharina latissima* (Faroes)
- Determine the antioxidant composition of the extracts (major compounds)
- Evaluate the *in vitro* antioxidant properties of the extracts
- Evaluate the antioxidant efficacy of seaweed extracts in skin care model systems (facial cream)

Extraction process (Solid Liquid Extraction)



Extract overview

Study	Seaweed	Solvent	Code
S1	<i>Fucus vesiculosus</i>	Water	FWE
		80% ethanol	F80EE
S2	<i>Saccharina latissima</i>	Water	SWE
		50% ethanol	S50EE
		70% ethanol	S70EE



S. latissima, 70 and 50 % ethanol extracts

Extract screening


Total phenolic content and antioxidant properties (mean±sd)

Seaweed	Solvent	Code	TPC	Metal chelating ability		DPPH radical scavenging capacity		Reducing power	
			TPC (mg GAE/L)	EC50 (mL/mL)	I _{max} (%) [*]	EC50 (mL/mL)	I _{max} (%) [*]	EC0.5 (mL/mL)	I _{max} (Abs) [*]
<i>Fucus vesiculosus</i>	Water	FWE	867.2±34.9	0.12±0.03	90.7±3.5	4.2*10 ⁻³ ±0.2	89.5±3.1	0.4±0.1	1.7±0.2
	80% ethanol	F80EE	830.3±19.8	0.09±0.01	88.3±5.3	3.7*10 ⁻³ ±0.1	90.8±3.7	0.4±0.0	1.8±0.1
<i>Saccharina latissima</i>	Water	SWE	160.9±0.3	0.5±0.0	61.6±1.9	0.43	82.6±3.8	ND	0.4±0.0
	50% ethanol	S50EE	141.5±0.3	ND	29.1±7.5	0.34	82.6±3.8	ND	0.7±0.0
	70% ethanol	S70EE	111.8±0.0	ND	51.5±7.6	0.42	76.3±4.7	ND	0.5±0.0

*Undiluted extract solution

Metal chelating ability

F. vesiculosus vs *S. latissima* (water extracts)



Chemical composition

Pigments

Pigments (mg/100 g dw)	<i>Fucus vesiculosus</i>		<i>Saccharina latissima</i>		
	FWE	F80EE	SWE	S50EE	S70EE
Chlorophyl c3	nd	nd	nd	0.3±0.0	35.4±16.6
Chlorophyl c2	nd	6.8±0.7	nd	0.4±0.2	11.7±0.9
19-But- Fucoxanthin	nd	20.2±2.4	nd	nd	6.5±4.2
Fucoxanthin	nd	nd	nd	0.1±0.0	13.6±0.8
Lutein	nd	0.8±0.1	0.2	nd	nd
β-carotene	nd	0.6±0.4	3.4±0.1	nd	nd

Storage trials



S1

F. vesiculosus added to facial cream formulation


Storage trial S1:

- 0.5 and 1% extract added (water or 80% (v/v) ethanol extract)
 - Samples: WE0.5, WE1, EE0.5 or EE1 (+REF without added extract)
- Storage for 42 days (dark, room temp)
- Determine oxidative stability: i.e fatty acid composition, peroxide value, secondary oxidation products, and tocopherols
- Almond oil and waxes used for the formulation:
 - 18:1: 54 %
 - 18:2: 18 %

S1

F. vesiculosus added to facial cream formulation

Results: Peroxide value



- EE0.5, EE1 and WE1, had a significantly higher PV than REF at day 0
- Maximum PV at day 28 for samples with extracts
- Oxidation rate at day 28:
 - REF > EE0.5, EE1 and WE1

S1

F. vesiculosus added facial cream formulations

Results: Secondary volatile oxidation products

Oxidation rate					
	REF	EE0.5	EE1	WE0.5	WE1
Heptanal	4.92%	-2.12%	1.82%	5.26%	1.04%
Hexanal	15.56%	7.81%	6.82%	15.76%	11.47%
2-Pentylfuran	86.28%	60.01%	72.42%	73.55%	68.25%

- Generally EE0.5 and EE1 had a lower oxidation rate compared with REF and cream added FWE
- Antioxidation effect
 - EE1 > EE0.5 > WE1 > WE0.5 ≈ REF
 - EE1 showed the lowest oxidation rate for hexanal

S1

F. vesiculosus added facial cream formulation

Total Tocopherol Content			
	Day 0	Day 42	Percentage Change
unit	µg toc/g oil	µg toc/g oil	(%)
REF	5937 ± 307	5072 ± 383	-14.58 %
EE0.5	5155 ± 120	5292 ± 93.5	2.67 %
EE1	5379 ± 167	5440 ± 48.3	1.13 %
WE0.5	5432 ± 95.6	4950 ± 88.7	-8.87 %
WE1	6138 ± 9.66	5264 ± 82.9	-14.14 %

- REF and WE1: significant decrease in TTC between day 0 and 42
- No tocopherol consumption when ethanol extract was added
 - Phenolic compounds used before tocopherols?
 - A synergistic effect between tocopherol and phenolic compounds or pigments?
 - Regeneration of tocopherol?

S2

S. Latissima added to commercial cream


Storage trial S2:

- 0.2 and 2% extract added (water, 50% or 70% (v/v) ethanol extract)
 - Samples: WE0.2, WE2, 50EE0.2, 50EE2, 70EE0.2 or 70EE2 (+CON without added extract)
- Abricot oil used in the creams:
 - 45 % 18:1 n-9
 - 15 % 18:2 n-6
- Storage for 70 days (dark, room temp)
- Determine oxidative stability
 - Fatty acid composition
 - Peroxide value
 - Secondary oxidation products
 - Tocopherol consumption
- Microbial contamination of WE2 (not included)

S2


S. Latissima added to commercial cream

Results: Peroxide value and secondary oxidation products (e.g. hexanal)



Preliminary results

- No antioxidant protection of unsaturated fatty acids compared to control
- Prooxidant activity towards formation of primary oxidation products?
- No clear development in secondary oxidation products
- No consumption of α -tocopherol during storage (not shown)



The cream is stable

Conclusion

- Antioxidant was extracted from both types of seaweed with water and aqueous ethanol solutions using SLE.
- *F. vesiculosus* extracts was higher in TPC and *in vitro* antioxidant capacity compared with *S. latissima*
- Aqueous ethanol solutions extracted fucoxanthin from the seaweed
- Application of *F. vesiculosus* extracts improved the oxidative stability of the facial cream when stored in darkness → decreased formation of primary and secondary oxidation products
- No clear antioxidant activity of *S. latissima* in facial cream (cream itself was stable)
 - Accelerated oxidation studies are needed

Acknowledgement



Mellisa[®]
Organic skincare



DTU Food
National Food Institute

Susan L. Holdt



Innovation Fund Denmark



DANISH
TECHNOLOGICAL
INSTITUTE



Thank you!