



Fast Water Transport in CNTs: length dependence and entrane/exit effects

Walther, Jens Honore; Koumoutsakos, Petros

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Walther, J. H., & Koumoutsakos, P. (2011). Fast Water Transport in CNTs: length dependence and entrane/exit effects. Abstract from 64th Annual Meeting of the American Physical Society's Division of Fluid Dynamics , Baltimore, MD, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract Submitted
for the DFD11 Meeting of
The American Physical Society

Sorting Category: 35. (C)

Fast Water Transport in CNTs: length dependence and entrance/exit effects JENS HONORE WALTHER, TU Denmark, PETROS KOUMOUTSAKOS, ETH Zurich — Superfast water transport in carbon nanotube (CNT) membranes has been reported in experimental studies. We use Molecular Dynamics simulations to elucidate the mechanisms of water entry, exit and transport in 2 nm -diameter hydrophobic CNTs embedded in a hydrophilic membrane matrix. We demonstrate, for the first time, that under imposed pressures of the order of 1 bar, water entry into the CNT cavity and exit from the CNT end, can occur only on pre-wetted membranes. We conduct large scale simulations for up to 500 nm long CNTs and observe a previously unseen dependence of the flow enhancement rates on the CNT length. We relate the present findings to past computational and experimental studies, we discuss previous continuum assessments for this flow and propose underlying physical mechanisms.

Prefer Oral Session
 Prefer Poster Session

Petros Koumoutsakos
petros@ethz.ch
ETH Zurich

Date submitted: 05 Aug 2011

Electronic form version 1.4